
IB ALL HL P2 REVISION BOOK

HIGHER LEVEL

1165marksPaper 22019 to 2015 inclusive (DRAFT)

Missing 17w, 2019sTZ1, 2019sTZ2


All Higher Level topics have been combined with their respective Standard Level equivalent, so in HL topic 2 revision resources there will be topics 2 and 12.



Contents

Patrick Brannac	www.SmashingScience.org	Page 1 of 314	
Topic Chem 2		2	3
Topic Chem 1			5
Standard and Higher Level compo	nents compared		3
All topics ranked according to their	r impact on your final grade using exam papers	from 1999 to 2016	3
Missing 17w, 2019sTZ1, 2019sTZ2			1


Topic Chem 3	28
Topic Chem 4	40
Topic Chem 5	67
Topic Chem 6	89
Topic Chem 7	111
Topic Chem 8	
Topic Chem 9	
Topic Chem 10	
Topic Chem 11	
Mark Scheme	219
Topic Chem 1	219
Topic Chem 2	
Topic Chem 3	
Topic Chem 4	233
Topic Chem 5	245
Topic Chem 6	
Topic Chem 7	
Topic Chem 8	267
Topic Chem 9	
•	
Topic Chem 11	302



All topics ranked according to their impact on your final grade using exam papers from 1999 to 2016


Topic	Option	10	N.ofSci	8	4	5	9	6	1	3	7	2	11
Rank	1	2	3	4	5	6	7	8	9	10	11	12	14
% of IB Grade	16.6	8.4	7.4	6.5	6.3	6.3	5.8	5.0	4.1	4.0	3.1	2.5	1.8
Paper 1		14.0		12.8	11.5	12.2	9.6	8.0	8.4	9.3	3.7	4.7	3.4
Paper 2 A		11.1		9.2	9.3	12.3	9.5	11.9	11.8	6.3	6.1	6.0	3.2
Paper 2 B		20.0		12.6	13.2	9.1	12.2	6.9	1.6	5.8	6.9	2.9	2.9
Paper 2 ALL		15.6		10.9	11.2	10.7	10.8	9.4	6.7	6.1	6.5	4.5	3.1
Paper 3	69.2		30.8										

Standard and Higher Level components compared



Essentially, IA has the exact same weight, the Option in HL is almost 50% more important than in SL but Topic 10 is more important in SL than HL. All other topics contribute almost equally to a SL and HL grade.





ALL PAPERS Topic Number

The dark blue bars are where your final IB grade will be from:

- 1. Your IA is the single most important part of your IB HL, more important than even the Option. Imagine how much time in class, at home and in revision you have or will give to topics 9, 10 and 11. Your IA, on average, will be worth more to your final grade than all those combined.
- 2. The Option is the most important topic for your IB grade compared to the everything else
- 3. Topic 10, Organic Chemistry, is by far the most important topic for papers 1 and 2.

(a)	/i\		
	(i)	Deduce the equation for the decomposition of guanidinium nitrate.	[1]
	(ii)	Calculate the total number of moles of gas produced from the decomposition of 10.0 g of guanidinium nitrate.	[1]
	(iii)	Calculate the pressure, in kPa, of this gas in a 10.0 dm ³ air bag at 127°C, assuming no gas escapes.	[1]
	(iv)	Suggest why water vapour deviates significantly from ideal behaviour when the gases are cooled, while nitrogen does not.	[2]
		SMASHINGILL	
(b)			[2]
		(iii)	(iii) Calculate the pressure, in kPa, of this gas in a 10.0 dm³ air bag at 127°C, assuming no gas escapes. (iv) Suggest why water vapour deviates significantly from ideal behaviour when the gases are cooled, while nitrogen does not.

Topic Chem 1 Q# 1/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q8./ www.SmashingScience.org:o)

Topic Chem 1 Q# 2/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org:o)

- 6. Copper forms two chlorides, copper(I) chloride and copper(II) chloride.
 - (ii) Copper(II) chloride is used as a catalyst in the production of chlorine from hydrogen chloride.

$$4HCl(g) + O_2(g) \rightarrow 2Cl_2(g) + 2H_2O(g)$$

(b) Solid copper(II) chloride absorbs moisture from the atmosphere to form a hydrate of formula CuCl₂•xH₂O.

A student heated a sample of hydrated copper(II) chloride, in order to determine the value of x. The following results were obtained:

Mass of crucible = 16.221 g
Initial mass of crucible and hydrated copper(II) chloride = 18.360 g
Final mass of crucible and anhydrous copper(II) chloride = 17.917 g

Determine the value of x. [3]

Topic Chem 1 Q# 3/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q5/ www.SmashingScience.org:o)

This reaction is used in the manufacture of sulfuric acid.

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$
 $K_c = 280$ at 1000 K

(a) State why this equilibrium reaction is considered homogeneous. [1]

SMASHINGILLI



Topic Chem 1 Q# 4/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q2/ www.SmashingScience.org:o)

An organic compound containing carbon, hydrogen and oxygen has 62.02% carbon and

	10.4	3% hydrogen by mass.	
·	(a)	Determine the empirical formula of the compound, showing your working.	[3]
	* *	* ***** ***** ***** ***** ***** ***** ****	
	\$\$\$\$\$\$\$\$		
	5.7.5		
		-	
Горіс 1.	3.2	1 Q# 5/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o) 6g of iron powder are added to 80.0 cm ³ of 0.200 mol dm ⁻³ copper(II) sulfate solution. The owing reaction occurs:	
		$Fe(s) + CuSO_4(aq) \rightarrow FeSO_4(aq) + Cu(s)$	
0	(a)	(i) Determine the limiting reactant showing your working.	[2]
	• •		
	* *		
#I		(ii) The mass of copper obtained experimentally was 0.872g. Calculate the percentage yield of copper.	[2]
	*11.50		



A student determined the percentage of the active ingredient magnesium hydroxide, Mg(OH)2, in a 1.24g antacid tablet. The antacid tablet was added to 50.00 cm3 of 0.100 mol dm-3 sulfuric acid, which was in excess. (a) Calculate the amount, in mol, of H₂SO₄. [1] Formulate the equation for the reaction of H₂SO₄ with Mg(OH)₂. (b) [1] The excess sulfuric acid required 20.80 cm³ of 0.1133 mol dm⁻³ NaOH for neutralization. (c) Calculate the amount of excess acid present. [1] Calculate the amount of H₂SO₄ that reacted with Mg(OH)₂. (d) [1]

Topic Chem 1 Q# 6/ IB Chem/2018/s/TZ 2/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o)



	(e)	Determine the mass of Mg(OH) ₂ in the antacid tablet.	[2]
	7.7.7		
	(f)	Calculate the percentage by mass of magnesium hydroxide in the 1.24g antacid tablet to three significant figures.	[1]

Горіс 1.		1 Q# 7/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q1/ www.SmashingScience.org :o) a, (H ₂ N) ₂ CO, is excreted by mammals and can be used as a fertilizer.	
	(a)	(i) Calculate the percentage by mass of nitrogen in urea to two decimal places using section 6 of the data booklet.	[2]
	(a)		[1]
	(a)	section 6 of the data booklet. (ii) Suggest how the percentage of nitrogen affects the cost of transport of fertilizers	[1]
	(a)	section 6 of the data booklet. (ii) Suggest how the percentage of nitrogen affects the cost of transport of fertilizers	[1]



(b) The structural formula of urea is shown.

(c) Urea can be made by reacting potassium cyanate, KNCO, with ammonium chloride, NH₄Cl.

$$KNCO(aq) + NH_4Cl(aq) \rightarrow (H_2N)_2CO(aq) + KCl(aq)$$

Determine the maximum mass of urea that could be formed from 50.0 cm³ of 0.100 mol dm⁻³ potassium cyanate solution.

[2]

			-		*	 					+		to L1	- 1	 		 Rif		-	-	r s	-	•		-									*					-			 		ne.						÷	7.		1.7	+	*	17:1
 ٠	٠	•		•	•		-	*	٠	+	*	•	+50	* 000	 	-01	 			0	(*) ·	*		*	*	-	•	•			*	-	•	*	-	*	*	•	*:	•	+ -		-	æ	•	*		*		٠	*	٠			٠	+
×					•			÷	+	*	*	*	483	*:0	 				- 14	100		2		*					24		×-	-		4	-			÷		-			-			4		-	Q.		ş		-			
 ×	٠	•		٠	•			+	*	×	**	*	800	*6	 						*0.00		•				• 15						*	+	-			*	*	*		 	-		 ٠	*		×	+	•	*	(*)	×	٠	٠	*
			-					-			*	-0		*10			2000	 						+		. ,		632							-					-			-	20			-				¥					
	• •	•••				 																																																		

(f) The combustion of urea produces water, carbon dioxide and nitrogen.

Formulate a balanced equation for the react	ion.
---	------

[2]

	A. I. (C. A

(g) Calculate the maximum volume of CO₂, in cm³, produced at STP by the combustion of 0.600 g of urea, using sections 2 and 6 of the data booklet.

[1]



Topic	Chem 1 Q# 8/	IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q2/ www.SmashingScience.org :o)	
2.		acidic sample of a waste solution containing Sn ²⁺ (aq) reacted completely with solution to form Sn ⁴⁺ (aq).	
	(iii)	13.239 g of K ₂ Cr ₂ O ₇ (s) were dissolved in distilled water to form a 0.100 dm ³ solution. Calculate its molar concentration.	I

_	(iv)	$10.0\mathrm{cm^3}$ of the waste sample required $13.24\mathrm{cm^3}$ of the $\mathrm{K_2Cr_2O_7}$ solution. Calculate the molar concentration of $\mathrm{Sn^{2+}}(\mathrm{aq})$ in the waste sample.	I
	*****	**************************	

		many oxides of silver with the formula Ag_xO_y . All of them decompose into their when heated strongly. After heating 3.760 g of a silver oxide 3.275 g of silver remained. Determine the empirical formula of Ag_xO_y .	[2
e e			
	(ii)	Suggest why the final mass of solid obtained by heating 3.760 g of Ag _x O _y may be greater than 3.275 g giving one design improvement for your proposed suggestion. Ignore any possible errors in the weighing procedure.	[2
		suggestion. Ignore any possible errors in the weighing procedure.	
		suggestion. Ignore any possible errors in the weighing procedure.	



	(b)	Nat	turally occurring silver is composed of two stable isotopes, ¹⁰⁷ Ag and ¹⁰⁹ Ag.	
			e relative atomic mass of silver is 107.87. Show that isotope ¹⁰⁷ Ag is re abundant.	[1]

A	(d)		ine the model of electron configuration deduced from the hydrogen line emission ctrum (Bohr's model).	[2]
	2/2/2			

opic	Chem	1 Q # 1	L1/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q6/ www.SmashingScience.org :o)	
	(c)	(i)	One possible product, X , of the reaction of ethane with chlorine has the following composition by mass:	
			carbon: 24.27%, hydrogen: 4.08%, chlorine: 71.65%	
			Determine the empirical formula of the product.	[2]
			SMASHINGLLI	

Topic Chem 1 Q# 10/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q1/ www.SmashingScience.org :o)



Topic Chem 1 Q# 12/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q5 DATA Booklet T22/www.SmashingScience.org:o)

5				THE RESERVE OF THE PROPERTY OF	March 1997 Annie 1997 December 1997 Annie 19	
1	h۱	Hydrazine is used to	remove ovviden	from water used	to generate steam	or hot water
٦	11/	Tryurazine is used to	ICHIOVE ONYGEN	HOITI Water useu	to generate steam	of flot water.

$$N_2H_4(aq) + O_2(aq) \rightarrow N_2(g) + 2H_2O(l)$$

The concentration of dissolved oxygen in a sample of water is $8.0 \times 10^{-3} \, \text{g} \, \text{dm}^{-3}$.

(i)	Calculate, showing your working, the mass of hydrazine needed to remove all the dissolved oxygen from 1000 dm ³ of the sample.	[

Calculate the volume, in dm³, of nitrogen formed under SATP conditions.

Topic Chem 1 Q# 13/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q2d

(ii)

- Titanium and vanadium are consecutive elements in the first transition metal series.
 - (h) TiCl₄ reacts with water and the resulting titanium(IV) oxide can be used as a smoke screen.

(1)	Formulate an equation for this reaction.	
	1/ 1-11114-1 1 5 1	
	**** ***** ***** ** **** ** **** **** ****	rancia and endicate and

(ii)	Suggest one disadvantage of using this smoke in an enclosed space.	

Topic Chem 1 Q# 14/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/zzzz

The concentration of a solution of a weak acid, such as ethanedioic acid, can be determined by titration with a standard solution of sodium hydroxide, NaOH(aq).



(C)	5.00 g of an impure sample of hydrated ethanedioic acid, (COOH), •2H,O, was dissolved
	in water to make 1.00 dm3 of solution. 25.0 cm3 samples of this solution were titrated
	against a 0.100 mol dm ⁻³ solution of sodium hydroxide using a suitable indicator.

$$(\mathsf{COOH})_2(\mathsf{aq}) + 2\mathsf{NaOH}(\mathsf{aq}) \rightarrow (\mathsf{COONa})_2(\mathsf{aq}) + 2\mathsf{H}_2\mathsf{O}(\mathsf{I})$$

	A STATE OF THE STA	
1	The mean value of the titre was 14.0 cm ³ .	
(ii	i) Calculate the amount, in mol, of NaOH in 14.0 cm ³ of 0.100 mol dm ⁻³ solution.	[1]
* * * * * *		
(ii	ii) Calculate the amount, in mol, of ethanedioic acid in each 25.0 cm ³ sample.	[1]
/ */* */*		
(i	v) Determine the percentage purity of the hydrated ethanedioic acid sample.	[3]

* * * * * *		

Topic Chem 1 Q# 15/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/Q7/ www.SmashingScience.org :o)

- (b) Bute, a painkiller used on horses, has caused widespread concern recently because analytical tests showed that it entered the food chain through horse meat labelled as beef. The drug is suspected of causing cancer.
 - (i) Analysis of a sample of bute carried out in a food safety laboratory gave the following elemental percentage compositions by mass:

Element	Percentage
С	73.99
Н	6.55
N	9.09
О	Remainder

	irical formula of bute, showing your working.	

	135000 1207	
The molar mass. M	f, of bute, is 308.37 g mol ⁻¹ . Calculate the molecular	formula.
1	1 42	



Topic Chem 1 Q# 16/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/

- Two IB students carried out a project on the chemistry of bleach.
 - (a) The bleach contained a solution of sodium hypochlorite, NaClO(aq). The students determined experimentally the concentration of hypochlorite ions, ClO, in the bleach:

Experimental procedure:

- The bleach solution was first diluted by adding 25.00 cm³ of the bleach to a 250 cm³ volumetric flask. The solution was filled to the graduation mark with deionized water.
- 25.00 cm³ of this solution was then reacted with excess iodide in acid.

$$ClO^{-}(aq) + 2I^{-}(aq) + 2H^{+}(aq) \rightarrow Cl^{-}(aq) + I_{2}(aq) + H_{2}O(1)$$

 The iodine formed was titrated with 0.100 mol dm⁻³ sodium thiosulfate solution, Na₂S₂O₃(aq), using starch indicator.

$$I_2(aq) + 2S_2O_3^{2-}(aq) \rightarrow 2\Gamma(aq) + S_4O_6^{2-}(aq)$$

The following data were recorded for the titration:

	First titre	Second titre	Third titre
Final burette reading of $0.100 \text{mol dm}^{-3} \text{Na}_2 \text{S}_2 \text{O}_3 \text{(aq)}$ (in cm ³ ± 0.05)	23.95	46.00	22.15
Initial burette reading of $0.100 \mathrm{mol}\mathrm{dm}^{-3}\mathrm{Na_2S_2O_3(aq)}$ (in cm ³ \pm 0.05)	0.00	23.95	0.00

c me vomme.	ш сш,	or o'roo moran	1 11a23203(aq)	required to	react
			11 10 11 1		[1]
			iodine to reach the end point.		te the volume, in cm ³ , of 0.100 mol dm ⁻³ Na ₂ S ₂ O ₃ (aq) required to iodine to reach the end point.



(11)	Calculate the amount, in mol, of Na ₂ S ₂ O ₃ (aq) that reacts with the iodine.	[1]
iii)	Calculate the concentration, in mol dm ⁻³ , of hypochlorite ions in the diluted bleach solution.	[1]
v)	Calculate the concentration, in mol dm ⁻³ , of hypochlorite ions in the undiluted bleach solution.	[1]
v)		
pic	Chem 1 Q# 17/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/	[1]
pic	bleach solution.	[1] PH ₃ .
	Chem 1 Q# 17/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/ Phosphine (IUPAC name phosphane) is a hydride of phosphorus, with the formula (b) Phosphine is usually prepared by heating white phosphorus, one of the allotre of phosphorus, with concentrated aqueous sodium hydroxide. The equation is	[1] PH ₃ .



	$P_4(s) + 3OH^-(aq) + 3H_2O(l) \rightarrow PH_3(g) + 3H_2PO_2^-(aq)$	
(i)	Calculate the amount, in mol, of white phosphorus used.	[1
(ii)	This phosphorus was reacted with 100.0 cm³ of 5.00 mol dm⁻³ aqueous sodium hydroxide. Deduce, showing your working, which was the limiting reagent.	[1]

* * * * * * * * * *		
(iii)	Determine the excess amount, in mol, of the other reagent.	[1]

(iv)	Determine the volume of phosphine, measured in cm³ at standard temperature and pressure, that was produced.	[1]

(c) 2.478g of white phosphorus was used to make phosphine according to the equation:

(d) Impurities cause phosphine to ignite spontaneously in air to form an oxide of phosphorus and water.



1 Mole of Phosphine was completely combusted in air./ www.SmashingScience.org:0)

(iii)) The molar mass of the oxide is ap Determine the molecular formula	

The oxide formed in the reaction with air contains 43.6 % phosphorus by mass.

Topic Chem 1 Q# 18/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/

1. A student used the technique of titration to determine the concentration of ascorbic acid (C₆H₈O₆) in a sample of orange juice. Excess potassium iodide, KI(aq), was added to acidified orange juice. The resulting solution was titrated with potassium iodate, KIO₃(aq), in the presence of starch as an indicator. The end-point of the titration was shown by a blue-black colour.

Step 1
$$IO_3^-(aq) + 5I^-(aq) + 6H^+(aq) \rightarrow 3I_2(aq) + 3H_2O(l)$$

lodine is only slightly soluble in water; but in the presence of excess iodide ions, $I^-(aq)$, it forms the soluble tri-iodide ion, $I_3^-(aq)$.

Step 2
$$I_2(aq) + I^-(aq) \rightleftharpoons I_3^-(aq)$$

Ascorbic acid reacts with tri-lodide ions as follows.

Step 3
$$C_6H_8O_6(aq) + I_3^-(aq) \rightarrow C_6H_6O_6(aq) + 2H^+(aq) + 3I^-(aq)$$



(b)	Calculate the mass, in g, of potassium iodate 0.250 dm ³ of a 2.00 × 10 ⁻³ mol dm ⁻³ solution.	k, KIO ₃ (s), which	h was required	d to prepare	[2]
(C)	The concentration of KIO ₃ used in the titratio The titration produced the following results.	n was 2.00 × 1	0 ⁻³ moldm ⁻³ .		
		Titration 1	Titration 2	Titration 3	
	Final volume of KIO ₃ (± 0.05 cm ³)	7.10	14.40	21.60	
	Initial volume of KIO ₃ (± 0.05 cm ³)	0.00	7.10	14.40	
	Volume added of KIO ₃ (± 0.10 cm ³)	7.10	7.30	7.20	
	Mean volume added of KIO ₃ (± 0.10 cm ³)		7.20		
	(iii) Determine the amount, in mol, of KIO ₃	(aq), in the me	an volume.		[1]
(d)	Determine the amount, in mol, of ascorbic acorange juice.	cid, C ₆ H ₈ O ₆ (aq)), in the sample		[2]
(e)	Calculate the mass, in g, of ascorbic acid, C, acidified orange juice.	₆ H ₈ O ₆ (aq), pres	sent in the san	nple of	[1]



Topic Chem 1 Q# 19/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/

 Ethanedioic acid is a diprotic acid. A student determined the value of x in the formula of hydrated ethanedioic acid, HOOC-COOH•xH₂O, by titrating a known mass of the acid with a 0.100 mol dm⁻³ solution of NaOH (aq).

0.795 g of ethanedioic acid was dissolved in distilled water and made up to a total volume of 250 cm³ in a volumetric flask.

25 cm³ of this ethanedioic acid solution was pipetted into a flask and titrated against aqueous sodium hydroxide using phenolphthalein as an indicator.

The titration was then repeated twice to obtain the results below.

Volume of 0.100 mol dm ⁻³ NaOH / cm ³	Titration 1	Titration 2	Titration 3
Final burette reading (± 0.05)	13.00	25.70	38.20
Initial burette reading (± 0.05)	0.00	13.00	25.70
Volume added	La Carlo		

1)	C	a														_																				*	ir	1 (CI	n	0,	1	n	t	tr	a	ti	10	15	5 2	2	a	n	d	3		a	n	d	ti	h	eı	n			
		_	_	_	_	_	_	_	_	-	_	_	_	_	_	_	_	_		_	_	_	_	_	_	_	_	_	_									_	_		_					_	_	_	_	_	_	_		_	_	_	_	_	H	_	_	_	_	_	_	_
	7			•	-	•					-					10			7		0,					*		-			87.			9.5	N/I						*0		1		7.5	•	1/3			*	1	36													*:	
																										*		17												-						+ 0	0.3			٠	+	*)				÷		+			+					
										4											- 34									2		10	8	V.	5			-	·	+		i i	*		+					+		410						4								
											,					0.5									,		100	70	+								-			-	*	100		(10)						*	٠	•										٠	÷			




(i)	The equation for the reaction taking place in the titration is:	
	$\mbox{HOOC-COOH}(\mbox{aq}) + 2\mbox{NaOH}(\mbox{aq}) \rightarrow \mbox{NaOOC-COONa}(\mbox{aq}) + 2\mbox{H}_2\mbox{O}(\mbox{l})$	
	Determine the amount, in mol, of ethanedioic acid that reacts with the average volume of NaOH (aq).	[1]
(ii)	Determine the amount, in mol, of ethanedioic acid present in 250 cm ³ of the original solution.	[1]
(iii)	Determine the molar mass of hydrated ethanedioic acid.	[1]
(iv)	Determine the value of x in the formula HOOC-COOH•xH ₂ O.	[2]

(b)

Topic Chem 2 Q# 20/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q4/ www.SmashingScience.org:o)

(c) Sketch a graph to show the relative values of the successive ionization energies of boron.





Topic Chem 2 Q# 21/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q4/ www.SmashingScience.org :o)

(c) The first six ionization energies, in kJ mol⁻¹, of an element are given below.

IE,	IE ₂	IE ₃	IE ₄	IE ₅	IE ₆
578	1816	2744	11576	14 829	18375

Explain the large increase in ionization energy from IE₃ to IE₄.

ro	7
1 2	
_	-

	en noncontra na manda per territa ben c



Topic	Chem 2 Q# 22/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/
4.	Magnesium is a group 2 metal which exists as a number of isotopes and forms many

(a) S		mbol notation, ^A X, for		
	* * * * * * * * * * * * * * * * * * * *		* * * * * * * * * * * * * * * * * * * *	
(b) N	Mass spectroscopic	analysis of a sample	of magnesium gave the fo	llowing results:
		5	% abundance	
		Mg-24	78.60	
		Mg-25	10.11	
	Calculate the relative places.	Mg-26	11.29 this sample of magnesium	to two decimal
p	places.	Mg-26 e atomic mass, A _r , of t		
	olaces.	e atomic mass, A _r , of t	this sample of magnesium	
	olaces.	e atomic mass, A _r , of t	this sample of magnesium	
	olaces.	e atomic mass, A _r , of t	this sample of magnesium	
	olaces.	e atomic mass, A _r , of the	this sample of magnesium	
	olaces.	e atomic mass, A _r , of the	this sample of magnesium	
	olaces.	Mg-26 e atomic mass, A _r , of the second sec	this sample of magnesium	




Patrick Brannac

	•••			
		(ii)	State what can be determined from the frequency of the convergence limit.	
	hem		23/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/	
(d)	(i)	St	ate the condensed electron configuration of sulfur.	[1]
	(ii)	De	educe the orbital diagram of sulfur, showing all the orbitals present in the diagra	m. [1]
			The state of the s	
. 0	One	of the	24/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/ e main constituents of acid deposition is sulfuric acid, H ₂ SO ₄ . This acid is foulfur dioxide pollutant, SO ₂ .	ormed
(e)			eference to section 9 of the data booklet, explain the difference between tradius and the ionic radius of nitrogen.	the [1]
	***	• • • •		
	* .*		******	



(e) Eight successive ionisation energies of vanadium are shown in the graph below:





(i) State the sub-levels from which each of the first four electrons are lost.	L
First: Second: Third: Fourth:	
(ii) Outline why there is an increase in ionization energy from electron 3 to electron 5.	[1
 (iii) Explain why there is a large increase in the ionization energy between electrons 5 and 6.	[3
(iv) Vanadium is comprised almost entirely of ⁵¹ V. State the number of neutrons an atom of ⁵¹ V has in its nucleus.	[1
Chem 2 Q# 26/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q7/ www.SmashingScience.org :c5 refers to the periodic table The two most abundant isotopes of bromine have the mass numbers 79 and 81. Calculate the relative abundance of ⁷⁹ Br using table 5 of the data booklet, assuming the abundance of the other isotopes is negligible. [2]))



6.	Copper forms two chlorides, copper(I) chloride and copper(II) chloride.	
Fig	(a) (i) State the electron configuration of the Cu ⁺ ion.	[1]
	(f) Dilute copper(II) chloride solution is light blue, while copper(I) chloride solution is colourle	SS.
55	 Describe how the blue colour is produced in the Cu(II) solution. Refer to section 17 of the data booklet. 	[3

	(ii) Deduce why the Cu(I) solution is colourless.	[1
Topic	Chem 3 Q# 28/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q4/ www.SmashingScience.org :o)	
Topic 4.	Chem 3 Q# 28/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q4/ www.SmashingScience.org :o) Properties of elements and their compounds can be related to the position of the elements in the periodic table.	
	Properties of elements and their compounds can be related to the position of the elements in	[2
	Properties of elements and their compounds can be related to the position of the elements in the periodic table.	[2
	Properties of elements and their compounds can be related to the position of the elements in the periodic table.	[2
	Properties of elements and their compounds can be related to the position of the elements in the periodic table.	[2

Topic Chem 3 Q# 27/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org:o)




<u>, </u>	(b) Explain why the radius of the sodium ion, Na ⁺ , is smaller than the radius of the oxide ion, O ²⁺ .	[2]
		; ;
Горі 3 .	ic Chem 3 Q# 29/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q3/ www.SmashingScience.org :o) Bromine can form the bromate(V) ion, BrO ₃ ⁻ .	
	(a) (i) State the electron configuration of a bromine atom.	[1]
	(ii) Sketch the orbital diagram of the valence shell of a bromine atom (ground state) on the energy axis provided. Use boxes to represent orbitals and arrows to represent electrons.	[1]
	Energy	
	(e) State and explain the magnetic property of iron(II) and iron(III) ions.	[2]

Topic Chem 3 Q# 30/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q2/ www.SmashingScience.org :o)


(c) When calcium compounds are introduced into a gas flame a red colour is seen; sodium compounds give a yellow flame. Outline the source of the colours and why they are different.

[2]




(e) Sketch a graph of the first six ionization energies of calcium.

[2]



Topic Chem 3 Q# 31/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q1/ www.SmashingScience.org :o)

(b) The structural formula of urea is shown.



(h) Describe the bond formation when urea acts as a ligand in a transition metal complex ion.

[2]

				ú	4			2					270	1						_		žV.	į.	-		0 0					0.2	Į.	2	_		2 :		4	2	20	4	2 (-	ú	J	ů,		
	*			10				-	nid.		•							57	Ť.S		•	10		15.0	*:					•			2	: T	•			0.00			•	***	•						
			•			•	•		·	*					٠				•			• (•		*			•			•		•		•			•	٠			•	•		 -				•	
					-	٠				- 1		*																											٠										
						•		-				+				*	•	*	٠					*				*	*				*							*	*				*				٠

Topic Chem 3 Q# 32/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q1 Data Booklet T22/ www.SmashingScience.org :o)

- There are many oxides of silver with the formula Ag_xO_y. All of them decompose into their elements when heated strongly.
 - (c) (i) Some oxides of period 3, such as Na₂O and P₄O₁₀, react with water. A spatula measure of each oxide was added to a separate 100 cm³ flask containing distilled water and a few drops of bromothymol blue indicator. The indicator is listed in section 22 of the data booklet.

Deduce the colour of the resulting solution and the chemical formula of the product formed after reaction with water for each oxide.


[3]

Flask containing	Colour of solution	Product formula
Na ₂ O	*****	
P ₄ O ₁₀		

Topic Chem 3 Q# 33/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q2d

- Titanium and vanadium are consecutive elements in the first transition metal series.
 - (iii) Sketch a graph of the first six successive ionization energies of vanadium on the axes provided.

[1]





(e)	Describe, in terms of the electrons involved, how the bond between a ligand and a central metal ion is formed.

(f)	Outline why transition metals form coloured compounds.
(f)	Outline why transition metals form coloured compounds.
(f)	Outline why transition metals form coloured compounds.
(f)	Outline why transition metals form coloured compounds.
(f)	Outline why transition metals form coloured compounds.
(f)	Outline why transition metals form coloured compounds.
(f)	

Topic Chem 3 Q# 34/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q2/ www.SmashingScience.org :o)

2. Titanium and vanadium are consecutive elements in the first transition metal series.



(b) Titanium exists as several isotopes. The mass spectrum of a sample of titanium gave the following data:

Mass number	% abundance
46	7.98
47	7.32
48	73.99
49	5.46
50	5.25

 	 	8	1 1 1	1 1/1			1121	2-1-2					12.0	2 2 2				 	
	 							7.7.5		 		• • •	1.7		*.*:			 10.57.5	
 	 									 					• •		*, **), *	 	
] 54 454			* * 1	K (# .4		 	
 	 					107 1S1		***		 								 	
		-			1	7			-		-	1							

Protons:		 	 	
Neutrons:				
	* * * * * * * *	 	 	***************************************
Electrons:				

(d) (i)	State the full electron configuration of the 48 Ti ²⁺ ion.	[1
*******	**** ***** ** *** ** *** ** ** ** ** **	

equati	on for the react	ion of magnesium	oxide with water.		
(f) Descr	be the trend in	acid-base properti	es of the oxides of p	period 3, sodium to	chlorine.
	ition to magness	eium ovide magne	sium forms another	compound when b	urned in
		ula of this compou			
air. S Chem 3 Q# 30	uggest the form	/w/TZ0/Paper 2 Sec			
air. S Chem 3 Q# 30	uggest the form	/w/TZ0/Paper 2 Sec	ind.		
air. S Chem 3 Q# 30	uggest the form	/w/TZ0/Paper 2 Sec	ind.		
air. S Chem 3 Q# 30	uggest the form	/w/TZ0/Paper 2 Sec	ind.		
air. S Chem 3 Q# 30	uggest the form	/w/TZ0/Paper 2 Sec	ind.		
air. S Chem 3 Q# 36 the Lewis stru	of IB Chem/2016 acture for the eth	/w/TZ0/Paper 2 Sec	ind.		
air. S Chem 3 Q# 30 the Lewis stru	of IB Chem/2016 acture for the eth	/w/TZ0/Paper 2 Sec nandioate ion:	ind.		
air. S Chem 3 Q# 30 the Lewis stru	of IB Chem/2016 acture for the eth	/w/TZ0/Paper 2 Sec nandioate ion:	ind.		
chem 3 Q# 30 the Lewis stru	of IB Chem/2016 acture for the eth	/w/TZ0/Paper 2 Sec nandioate ion:	ind.		

Topic Chem 3 Q# 37/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/

4. Tin(II) chloride is a white solid that is commonly used as a reducing agent.



(c)	Outline, giving the full electron configuration of the vanadium atom, what is meant by the term transition metal.	[2]
2 264		
* * *		
(d)	In an aqueous solution of vanadium(III) chloride, the vanadium exists as $[V(H_2O)_6]^{3+}$, $[VCl(H_2O)_5]^{2+}$ or $[VCl_2(H_2O)_4]^+$ depending on the concentration of chloride ions in the solution.	
	(i) Describe how Cl ⁻ and H ₂ O bond to the vanadium ion.	[1]
	(ii) Outline what would happen to the wavelength at which the vanadium complex ions would absorb light as the water molecules are gradually replaced by chloride ions, using section 15 of the data booklet.	[2]

12.5		

	130000	



(i)	State the acid-base character of the oxides of the period 3 elements Na to Ar. [2]	

(ii)	State balanced equations to illustrate the acid-base character of sodium oxide and sulfur trioxide. [2]	
	Sodium oxide:	

	Sulfur trioxide:	

	1, 44 (1) (1) (1)	
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/	
Chen (a)		[
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/	[1
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/	[
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/ (i) Define the term electronegativity.	[
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/ (i) Define the term electronegativity.	[1
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/ (i) Define the term electronegativity.	[1
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/ (i) Define the term electronegativity.	[]
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/ (i) Define the term electronegativity.	
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/ (i) Define the term electronegativity.	
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/ (i) Define the term electronegativity.	
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/ (i) Define the term electronegativity.	
	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/ (i) Define the term electronegativity.	
	(i) Define the term electronegativity. (ii) Suggest why the noble gases are generally not assigned electronegativity values. Outline one reason why the sodium ion, Na*, has a smaller radius than the sodium	[1
(a)	n 3 Q# 39/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/ (i) Define the term electronegativity. (ii) Suggest why the noble gases are generally not assigned electronegativity values.	[1
(a)	(i) Define the term electronegativity. (ii) Suggest why the noble gases are generally not assigned electronegativity values. Outline one reason why the sodium ion, Na*, has a smaller radius than the sodium	
(a)	(i) Define the term electronegativity. (ii) Suggest why the noble gases are generally not assigned electronegativity values. Outline one reason why the sodium ion, Na*, has a smaller radius than the sodium	[1
(a)	(i) Define the term electronegativity. (ii) Suggest why the noble gases are generally not assigned electronegativity values. Outline one reason why the sodium ion, Na*, has a smaller radius than the sodium	[1

(a)	State the full electron configurations of copper, Cu, and copper (II) ion, Cu ²⁺ .	
(c)	Explain why complexes of Zn ²⁺ (aq) are colourless whereas complexes of Cu ²⁺ (aq) are coloured.	

	3 Q# 41/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/Q9/ www.SmashingScience.org	_
(b)	Identify the acid-base character of the oxides of each of the elements from sodium to chlorine in period 3.	
	(i) Identify the acid-base character of the oxides of each of the elements from	-
	(i) Identify the acid-base character of the oxides of each of the elements from	
	(i) Identify the acid-base character of the oxides of each of the elements from	
	(i) Identify the acid-base character of the oxides of each of the elements from	
	(i) Identify the acid-base character of the oxides of each of the elements from	
	(i) Identify the acid-base character of the oxides of each of the elements from	
(b)	(i) Identify the acid-base character of the oxides of each of the elements from sodium to chlorine in period 3. State the equations for the separate reactions of sodium oxide and phosphorus(V)	
(b)	(i) Identify the acid-base character of the oxides of each of the elements from sodium to chlorine in period 3. State the equations for the separate reactions of sodium oxide and phosphorus(V)	

Topic Chem 3 Q# 42/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/Q8/ www.SmashingScience.org :o)

(c) Chromium forms the complex ion $[Cr(NH_3)_4Cl_2]^+$.



		43/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/	
(a)	(i)	Draw an orbital diagram (using the arrow-in-box notation) showing the electrons in the 4s and 3d sub-levels in chromium metal.	
	(ii)	Outline the nature of the metallic bonding present in chromium.	
	(iii)	Explain why chromium metal is malleable.	
		Water 11	_

(iv) Draw the structures of two possible isomers of this complex ion.



[2]

	Deduce the oxidation number of chromium in this complex.	I

	.,,	
(ii)	Describe the nature of the ligand-chromium ion bonds in terms of acid-base theory.	I

(iii)	Explain why [Cr(NH ₃) ₄ Cl ₂] ⁺ is coloured.	I

	W COLOR W TO A SECOND	

(c) Chromium forms the complex ion $[Cr(NH_3)_4Cl_2]^+$.



Topic Chem 3 Q# 45/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/

This question is about the compounds of some period 3 elements.

(a)		the equations for the reactions of sodium oxide with water and phosphorus(V) with water.	[2]
		6/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/ metal that has been used by humans for thousands of years.	
(a)	State t	the full electron configuration of ⁶⁵ Cu.	[1]

(b)		one difference in the physical properties of the isotopes ⁶³ Cu and ⁶⁵ Cu and n why their chemical properties are the same.	[2]
	Physic	al:	
	Chemi	cal:	
		# 47/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org :o) ms two chlorides, copper(I) chloride and copper(II) chloride.	
	(iii)	When excess ammonia is added to copper(II) chloride solution, the dark blue complex ion, [Cu(NH ₃) ₄ (H ₂ O) ₂] ²⁺ , forms.	е
		State the molecular geometry of this complex ion, and the bond angles within	n it.
Mo	olecular	geometry:	
:		nas tende de mas formes de mes dem to dem to dem do des do dem des de dem todo de tendeno.	
Во	ond angl	es:	



Topic Chem 4 Q# 48/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q2./ www.SmashingScience.org :o)

	ions show steps in the formation and decomposition of ozone in the stratosphere, hich absorb ultraviolet light.	
Step 1	$O_2 \rightarrow 2O^{\bullet}$	
Step 2	$O \cdot + O_2 \rightarrow O_3$	
Step 3	$O_3 \rightarrow O^{\bullet} + O_2$	
Step 4	$O \cdot + O_3 \rightarrow 2O_2$	
(a) Drav	v the Lewis structures of oxygen, O ₂ , and ozone, O ₃ .	[2
	ine why both bonds in the ozone molecule are the same length and predict the	
bono	I length in the ozone molecule. Refer to section 10 of the data booklet.	[
Reason:		
Length:		
Lengui.		
* * * * * * *		
(c) Pred	lict the bond angle in the ozone molecule.	[
	* ** *** *** ** *** *** *** *** *** **	



		cuss how the different bond strengths between the oxygen atoms in O ₂ and O ₃ in ozone layer affect radiation reaching the Earth's surface.	
* * *			
9 K.A			
(e)	(i)	Identify the steps which absorb ultraviolet light.	
	(ii)	Determine, showing your working, the wavelength, in m, of ultraviolet light absorbed by a single molecule in one of these steps. Use sections 1, 2 and 11 of the data booklet.	
10.00			
. 122			
(f)	070	ne depletion is catalysed by nitrogen monoxide, NO, which is produced in aircraft	
(.)		motor vehicle engines, and has the following Lewis structure.	
		$\dot{N}=0$	
		w how nitrogen monoxide catalyses the decomposition of ozone, including ations in your answer.	
1. V.			

Topic Chem 4 Q# 49/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q3/ www.SmashingScience.org :o)
3. Bromine can form the bromate(V) ion, BrO₃⁻.

(b) (i) Draw two Lewis (electron dot) structures for BrO ₃ ⁻ .	[2]
Structure I – follows octet rule:	
Structure II – does not follow octet rule:	
 (ii) Determine the preferred Lewis structure based on the formal charge on the bromine atom, giving your reasons. 	[2]
	7
(c) Predict, using the VSEPR theory, the geometry of the BrO ₃ ion and the O-Br-O bond angles.	[3]
Geometry:	
Reason:	
O-Br-O angle:	

Topic Chem 4 Q# 50/ IB Chem/2018/s/TZ 2/Paper 2/Higher Level/Q7/ www.SmashingScience.org :o)

Some physical properties of molecular substances result from the different types of forces between their molecules. (a) Explain why the hydrides of group 16 elements (H₂O, H₂S, H₂Se and H₂Te) are (i) polar molecules. [2] The graph shows the boiling points of the hydrides of group 16 elements. (ii) H_2O **3oiling point** ■H₂Te H₂Se 2 3 5 Period Explain the increase in the boiling point from H₂S to H₂Te. [2] Lewis structures show electron domains and are used to predict molecular geometry. Deduce the electron domain geometry and the molecular geometry for the NH₂ ion. [2] Electron domain geometry: Molecular geometry:

(c) Resonance structures e Lewis structure.	exist when a molecule can be repre	sented by more than one
(i) Carbon dioxide ca	an be represented by at least two re	esonance structures, I and II.
	T II	
	_r; :o _ r;	
(1)	=c==0 :o ≡ c0: (2) (1) (2)	
	nal charge on each oxygen atom in	the two structures. [2]
Structure	1	II
O atom labelled (1)		
O atom labelled (2)		***************
(ii) Deduce, giving a	reason, the more likely structure.	[1]
	n the ozone layer causes the dissonding, the molecule that requires a lo	

(e) Carbon and silicon are	elements in group 14.	
Explain why CO ₂ is a ga	as but SiO ₂ is a solid at room tempe	erature. [2]
************		*** ***** ***** *****
		SSEL PRINCIPLE AND A SERVICE AND SERVICE
	***********	****************

Topic Chem 4 Q# 51/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q3/ www.SmashingScience.org :o)

(d) One possible Lewis structure for benzene is shown.

	State one piece of physical evidence that this structure is incorrect .	[1
(3)	c Chem 4 Q# 52/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q2/ www.SmashingScience.org :o)	
2.	Calcium carbide, CaC ₂ , is an ionic solid.	
	(a) Describe the nature of ionic bonding.	[1
1	(d) (i) Suggest two reasons why solid calcium has a greater density than solid potassium.	[2]
1	(ii) Outline why solid calcium is a good conductor of electricity.	[1]

(g)	(i)	Describe h	ow sigma	(0)	and n	$i(\pi)$	honds are	formed
19/	11/	Describe II	OW Sigina	(0)	and p	1 (11)	DOINGS GIC	william.

sig	m	na	(3)	C.																																																															
														-										*		4		33.					*									8		4		+		٠	¥			•			-	*		٠	ě	í.								
				•				٠		•		•		•		•						•					•	•			•	•				•	•	•		•	7				4		٠				٠			38				•						4	•		•	
	/_																																																																			
pi ((π):																																																																		
• •	•		ं	*		ō	•	-	•	-	-	ं	•	ē		ं	•	*	7	-	•	•	Ť	Ţ.	*	i,			•	7	•	*	-	÷	7	Ţ.	-	57	÷	7		33	3	† ·	Ť	•	•		Ť	æ	- (*		7			Œ.	÷		c		3	•	•	•			•	÷
																																												war.																								

(ii) Deduce the number of σ and π bonds in a molecule of ethyne.

.

[1]


[2]

pi (π):

sigma (σ):

Topic Chem 4 Q# 53/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q1/ www.SmashingScience.org :0)

(b) The structural formula of urea is shown.



Predict the electron domain and molecular geometries at the nitrogen and carbon atoms, applying the VSEPR theory.

	Electron domain geometry	Molecular geometry
Nitrogen		
Carbon		trigonal planar

[3]



(e)	(i)	Suggest one reason why urea is a solid and ammonia a gas at room temperature.	[1]
	(ii)	Sketch two different hydrogen bonding interactions between ammonia and water.	[2]
(i)			[1
55.5			
	((((((((((((((((((((i) The	(ii) Sketch two different hydrogen bonding interactions between ammonia and water. (i) The C–N bonds in urea are shorter than might be expected for a single C–N bond. Suggest, in terms of electrons, how this could occur.



Topic Chem 4 Q# 54/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q4/ www.SmashingScience.org :o)

- Bonds can be formed in many ways.
 - (a) The landing module for the Apollo mission used rocket fuel made from a mixture of hydrazine, N₂H₄, and dinitrogen tetraoxide, N₂O₄.

$$2N_2H_4(l) + N_2O_4(l) \rightarrow 3N_2(g) + 4H_2O(g)$$

(i)	State and explain the difference in bond strength between the nitrogen atoms in a hydrazine and nitrogen molecule.	[2

(ii)	State why hydrazine has a higher boiling point than dinitrogen tetraoxide.	[1

7.67.6	ne has a higher boilin			
	 ********	to the temperature the	extend to the extended	

(b) (ı)	Discuss the bonding in the resonance structures of ozone.	[3
	A STATE OF LAND	

(ii)	Deduce one resonance structure of ozone and the corresponding formal charges on each oxygen atom.	[2



Topic Chem 4 Q# 55/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q3/ www.SmashingScience.org :o)

(b)	Deduce the Lewis (electron dot) structure, molecular geometry and the bond angles of PCl ₃ .	
Lew	vis structure:	
Mole	ecular geometry:	
Bon	nd angles:	
	nd angles: 14 Q# 56/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q1c (ii) Explain the electrical conductivity of molten Na ₂ O and P ₄ O ₁₀ .	
	14 Q# 56/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q1c	
	14 Q# 56/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q1c	
	14 Q# 56/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q1c	
	14 Q# 56/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q1c	
	14 Q# 56/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q1c	_



Topic Chem 4 Q# 57/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q8/ www.SmashingScience.org :o)

There is concern about damage done to the ozone layer in the stratosphere by jet-propelled aircraft. (a) Formulate two equations to show how nitrogen(II) oxide, NO, catalyses the destruction of ozone. [2] (b) Suggest why the loss of ozone is an international environmental concern. [2] Topic Chem 4 Q# 58/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q5/ www.SmashingScience.org :0) 5. Two hydrides of nitrogen are ammonia and hydrazine, N₂H₄. One derivative of ammonia is methanamine whose molecular structure is shown below. Estimate the H-N-H bond angle in methanamine using VSEPR theory. [1] State the electron domain geometry around the nitrogen atom and its hybridization in (b)

[2]

methanamine.

Topic Chem 4 Q# 59/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q2/ www.SmashingScience.org :o) Titanium and vanadium are consecutive elements in the first transition metal series. Describe the bonding in metals. [2] d (ii) Suggest why the melting point of vanadium is higher than that of titanium. [1] (iv) Explain why an aluminium-titanium alloy is harder than pure aluminium. [2] (g) (i) State the type of bonding in potassium chloride which melts at 1043 K. [1] A chloride of titanium, TiCl₄, melts at 248 K. Suggest why the melting point is so (11) much lower than that of KCl. [1]



	(h)		D	e	SC	:ri	b	е	th	ıe	. 5	st	rı	IC	tı	ur	е	é	ar	10	11	00	or	no	11	n	g	ir	1	SC	oli	Id	r	na	ıg	n	es	SIL	ur	n	0	XI	de	е.											
							0.5			*:													50																		10				-			**		 		 		 	
	250	*	70				-	To	*						-				ON:	•	-														*						(In	• (•	-	5.50	•		•		•	205.	

Topic Chem 4 Q# 61/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience.org:o)

(c) 5.00 g of an impure sample of hydrated ethanedioic acid, (COOH)₂•2H₂O, was dissolved in water to make 1.00 dm³ of solution. 25.0 cm³ samples of this solution were titrated against a 0.100 mol dm⁻³ solution of sodium hydroxide using a suitable indicator.

$$(COOH)_2(aq) + 2NaOH(aq) \rightarrow (COONa)_2(aq) + 2H_2O(l)$$

(d) Draw the Lewis (electron dot) structure of the ethanedioate ion, "OOCCOO".

[1]




(e) Outline why all the C–O bond lengths in the ethanedioate ion are the same length and suggest a value for them. Use section 10 of the data booklet.

[2]



Topic Chem 4 Q# 62/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/Not with Q5 (a)

Propane and propene are members of different homologous series.





(b) (i)	Draw diagrams to show how sigma (σ) and patoms.	oi (π) bonds are formed between [2]
Sigma (σ)		
Pi (π):		
(ii)	State the number of sigma (σ) and pi (π) bor	nds in propane and propene. [2
	Number of sigma (σ) bonds	Number of pi (π) bonds
Propane		
Propene		
1. Ethane-1,	63/ IB Chem/2016/w/TZ0/Paper 2 Section A/High 2-diol, HOCH₂CH₂OH, has a wide variety of us d heat transfer in a solar cell. rmula CH2CH2	
(d) Expl ethe	ain why the boiling point of ethane-1,2-diol is sine.	significantly greater than that of [2]



Topic Chem 4 Q# 64/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/

- Ozone, dinitrogen monoxide, CFCs, sulfur hexafluoride and methane are all examples of greenhouse gases.
 - (a) (i) Draw one valid Lewis (electron dot) structure for each molecule of the greenhouse gases listed below. [2]

	Lewis (electron dot) structure
Ozone	
Sulfur hexafluoride	

(ii) Deduce the name of the electron domain geometry and the molecular geometry for each molecule listed below. [2]

*	Electron domain geometry	Molecular geometry
Ozone		
Sulfur hexafluoride		



(111)	Identify which molecule(s) given in part (a) (i) has/have an extended octet of electrons.	[1]
(iv)	State the bond angles for each species in part (a) (ii).	[1]
	Ozone:	
	Sulfur hexafluoride:	
(v)	Draw all the resonance structures of ozone. Lone pairs should be shown.	[1]
	THEAT	



(b) Nitrous oxide can be represented by different Lewis (electron dot):	(b)	Nitrous oxide	can be represented	by different Lewis	(electron dot)	structures
---	-----	---------------	--------------------	--------------------	----------------	------------

(i)	Deduce the formal charge (FC) of the nitrogen and oxygen atoms in three of these
	Lewis (electron dot) structures, A, B and C, represented below.

[2]

LHS: atom on left-hand side; RHS: atom on right-hand side.

	Lewis (electron dot) structure	FC of O on LHS	FC of central N	FC of N on RHS
A	:ö==n==n:			
В	:ö—n <u>≡</u> n:			
C	:o=n-ii:			

(ii) FC can be useful in electron book-keeping, but electronegativity values are ignored when FCS are assigned.

Based on the assignment of FCS of the atoms in part (i), deduce which Lewis (electron dot) structure of N_2O (A, B or C) is expected to be the preferred structure. Explore why another factor needs to be considered.

[2]

				•			 140	4014									-	-				• * * *	-	-		-	201				-		0	 -	 			4	•
				•						•	. 5	3.1	* *																				*	 *	 		* *		
	 4				- 2	2			-					4	4	4				-	4		0	345		30			-							-			-
* *			OH:	+.			 .+:	- ,	100						*	#id			-					*	-	-	- 1	. 30	100	*:-	0	 -	-	 -	 00				+ -



(d) (i) The concentration of ozone in the upper atmosphere is maintained by the following three reactions, I, II and III.


I
$$O_2 \xrightarrow{hv} 2O_{\bullet}$$

II
$$O_2 + O \cdot \rightarrow O_3$$

III
$$O_3 \xrightarrow{hv} O_2 + O_4$$

Explain by reference to the bonding in O_2 and O_3 , which of the reactions, I or III, needs more energy.

[3]



(ii) Using dichlorodifluoromethane, CCl₂F₂, as an example, outline the reactions in which ozone depletion occurs in the upper atmosphere. Formulate an equation for each step in this process and explain the initial step by reference to the bonds in [5] CCl₂F₂.



Topic Chem 4 Q# 65/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q3/ www.SmashingScience.org:o)

(c) One of the intermediates in the reaction between nitrogen monoxide and hydrogen is dinitrogen monoxide, N₂O. This can be represented by the resonance structures below:

	(i)	Analyse the bonding in dinitrogen monoxide in terms of $\sigma\text{-bonds}$ and $\pi\text{-bonds}.$	[3]

	• • • • • • • • • • • • • • • • • • • •		

	(ii)	State what is meant by resonance.	[1]

Topic		66/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/ e (IUPAC name phosphane) is a hydride of phosphorus, with the formula PH ₃ .	
		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	F43
=0	(a) (i)	Draw a Lewis (electron dot) structure of phosphine.	[1]
		MAL NOT 1	
	(ii)	State the hybridization of the phosphorus atom in phosphine.	[1]



	giving a brief re	eason.			
				,	
			ar mass than ammon boiling point than am		
Chem -	4 Q# 67/ IB CHEM/2015	5/w/TZ0/Paper 2 Sect	ion B/Higher Level/Q6	/ www.SmashingScie	nce.org
Chem (i)	4 Q# 67/ IB CHEM/2015 Deduce the type of h N ₂ H ₄ .		(70)		(5%) }
	Deduce the type of h		(70)		(S)
	Deduce the type of h	nybridization shown	by the nitrogen atom	is in NF ₄ ⁺ , N ₂ H ₂ and	(S)
	Deduce the type of h	NF ₄ *	N₂H₂	is in NF ₄ ⁺ , N ₂ H ₂ and	0.70
(i)	Deduce the type of h N ₂ H ₄ . Hybridization	NF ₄ *	N₂H₂	is in NF ₄ ⁺ , N ₂ H ₂ and	[3
(i)	Deduce the type of h N ₂ H ₄ . Hybridization	NF ₄ *	N₂H₂	is in NF ₄ ⁺ , N ₂ H ₂ and	[3
(i)	Deduce the type of h N ₂ H ₄ . Hybridization	NF ₄ *	N₂H₂	is in NF ₄ ⁺ , N ₂ H ₂ and	[3
(i)	Deduce the type of h N ₂ H ₄ . Hybridization	NF ₄ *	N₂H₂	is in NF ₄ ⁺ , N ₂ H ₂ and	[3
(i)	Deduce the type of h N ₂ H ₄ . Hybridization	NF ₄ *	N₂H₂	is in NF ₄ ⁺ , N ₂ H ₂ and	[3
(i)	Deduce the type of h N ₂ H ₄ . Hybridization	NF ₄ *	N₂H₂	is in NF ₄ ⁺ , N ₂ H ₂ and	[3

(iii)	Draw the Lewis (electron dot) structures of SF ₄ and SF ₆ . Use the valence shell electron pair repulsion (VSEPR) theory to predict the name of the shape of each molecule.	
(i)	List the following compounds in order of increasing boiling point: CH ₃ CH ₀ , CH ₃ CH ₂ CH ₃ , CH ₃ COOH, CH ₃ CH ₂ OH.	
(ii)	Explain the order of boiling points in the compounds listed in part (d) (i), in terms of intermolecular forces.	




(b)	Explain why the melting points of the group 7 whereas the melting points of the group 7	o 1 metals (Li → Cs) decrease down the g elements (F → I) increase down the group

	n 4 Q# 69/ IB CHEM/2015/s/TZ2/Paper 2 Sections sider the molecules PBr ₃ and SF ₄ . Deduce the Lewis (electron dot) structure	
	134	
(ii)	Predict the shapes of the two molecules, g and the F–S–F bond angles in SF ₄ .	giving the Br–P–Br bond angle in PBr₃ SF₄
(ii)	and the F-S-F bond angles in SF ₄ .	-32 / \-
(ii)	and the F–S–F bond angles in SF ₄ . PBr ₃	SF ₄
(ii)	and the F–S–F bond angles in SF ₄ . PBr ₃	SF ₄
(ii)	PBr ₃ Shape:	SF₄ Shape:
(ii)	and the F–S–F bond angles in SF ₄ . PBr ₃	SF ₄
(ii)	PBr ₃ Shape:	SF₄ Shape:
(ii)	PBr ₃ Shape:	SF₄ Shape:

[3]

(iii)	Explain why both PBr ₃ and SF ₄ are polar.	[2]

(d) The structure of cis-but-2-ene-1,4-dioic acid is shown below.



 Describe the covalent bond between carbon and hydrogen in the molecule above and how it is formed.



[2]



	Dea	luce the hybridization of the oxygen atoms labelled α and β .	[1]
	α:		

	β:		
	444		
(iii)	Des	cribe sigma (σ) and pi (π) bonds between atoms.	[2]
	σbo	ond:	
		N	
	πbo	ond:	
	..*		

(iv)	lder	ntify the number of sigma (σ) and pi (π) bonds present in a molecule of but-2-ene-1,4-dioic acid.	[1]
(iv)	Ider cis-l	ntify the number of sigma (σ) and pi (π) bonds present in a molecule of but-2-ene-1,4-dioic acid.	[1]
(iv)	Ider	htify the number of sigma (σ) and pi (π) bonds present in a molecule of but-2-ene-1,4-dioic acid.	[1]
(iv)	Ider cis-I	but-2-ene-1,4-dioic acid.	[1]
	cis-l	but-2-ene-1,4-dioic acid.	
Chem -	cis-l	but-2-ene-1,4-dioic acid. 70/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/Q8/ www.SmashingScience.	
Chem -	cis-l	70/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/Q8/ www.SmashingScience. State the name of Cr ₂ O ₃ .	
Chem -	4 Q# (i)	70/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/Q8/ www.SmashingScience. State the name of Cr ₂ O ₃ .	
Chem -	4 Q# (i)	70/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/Q8/ www.SmashingScience. State the name of Cr ₂ O ₃ .	



(iii)	Suggest why solid Cr ₂ O ₃ de	oes not conduct electricity.	[1]
	4 Q# 71/ IB CHEM/2015/s/TZ1/cribe the formation of σ and π		/Q7/ www.SmashingScience.org :o [2]
	4 Q# 72/ IB CHEM/2015/s/TZ1/ tify three allotropes of carbon		/Q5/ www.SmashingScience.org :o 6. [4]
f) SCl ₂ a	4 Q# 73/ IB CHEM/2015/s/TZ1/ and SClF₅ are two sulfur chlor tion states. Predict the name cules.	ride type compounds with sul	fur having different
Molecule	Shape	Bond angle	Polarity
SCl ₂	***************		
SCIF ₅			



	(b)	(i)		Iting point of phosphorus(V) oxide is I eir bonding and structure.	ower than that of sodium [2

		(ii)	THE RESERVE OF THE PROPERTY OF	osphorus(V) oxide and sodium oxide of ates. Complete the boxes with "yes" of	그 가장 집에 가장 아니는 아니는 생각이 하는 것이 없는 사람들이 되었다. 그 사람들이 되었다면 하는데 하는데 하는데 되었다면 하는데
		_		Phosphorus(V) oxide	Sodium oxide
			Solid state		
			Molten state		
	Chem (c)		75/ IB Chem/2015/s/T cribe the bonding in s	Z1/Paper 2 Section A/Higher Level/ Q4/ olid copper.	www.SmashingScience.org :o)
		***			**************
		***			*****
opic (Etha	nedic	oic acid is a diprotic ac	Z1/Paper 2 Section A/Higher Level/ cid. A student determined the value of OC-COOH•xH ₂ O, by titrating a known OH(ag).	
c)				cular force in solid ethanedioic acid.	[1]
	/ */*				****



(d)	Deduce the Lewis (electron dot) structure of ethanedioic acid, HOOC-COOH. [1]	
(e)	Predict and explain the difference in carbon-oxygen bond lengths in ethanedioic acid and its conjugate base, "OOC-COO". [3]	
opic	 C Chem 5 Q# 77/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org:o) Copper forms two chlorides, copper(I) chloride and copper(II) chloride. (ii) Copper(II) chloride is used as a catalyst in the production of chlorine from hydrogen chloride. 4HCl(g) + O₂(g) → 2Cl₂(g) + 2H₂O(g) Calculate the standard enthalpy change, ΔH^o, in kJ, for this reaction, using 	
	section 12 of the data booklet.	[2]
	(d) Determine the enthalpy of solution of copper(II) chloride, using data from sections 18 and 20 of the data booklet.	
	The enthalpy of hydration of the copper(II) ion is -2161 kJ mol ⁻¹ .	[2]

Topic Chem 5 Q# 78/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q4./ www.SmashingScience.org:o)

 Propene is an important starting material for many products. The following shows some compounds which can be made from propene, C₃H₆.

Propene
$$(C_3H_6) \rightarrow C_3H_7Cl \rightarrow C_3H_8O \rightarrow C_3H_6O$$

(b) An experiment was carried out to determine the order of reaction between one of the isomers of C₃H₇Cl and aqueous sodium hydroxide. The following results were obtained.

Experiment	[C ₃ H ₇ CI] / mol dm ⁻³	[OHT] / mol dm ⁻³	Initial rate / mol dm ⁻³ s ⁻¹
1	0.05	0.10	3.1 × 10 ⁻⁴
2	0.10	0.20	1.3 × 10 ⁻³
3	0.15	0.10	9.2 × 10 ⁻⁴

2	(c)	(i)	Write an equation for the complete combustion of the compound C ₃ H ₈ O formed in (a)(iv).	[1]
		(ii)	Determine the enthalpy of combustion of this compound, in kJ mol ⁻¹ , using data from section 11 of the data booklet.	[3]



Topic Chem 5 Q# 79/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q7/ www.SmashingScience.org :o)

Consider the following Hess's law cycle:

$$CH_2 = CH_2(g) + H_2(g) \xrightarrow{Step 1} CH_3CH_3(g) + \frac{7}{2}O_2(g)$$
Step 2
 $+\frac{7}{2}O_2(g)$
Step 3

 $2CO_{2}(g) + 3H_{2}O(l)$

(b)	Calculate the standard enthalpy change, ΔH^{Θ} , of step 2 using section 13 of the
	data booklet.

[1]

 	 	 	 		::	 		5.7	-	 	 	 	*:*	 	::0	 	•	: :		 11 15	 	
 	 		 	+:+		 	* *				 	 	٠.,	 					٠.	• •	 	

(c)	Determine the standar	d enthalpy change, ΔH° , of step 1.	[1]
-----	-----------------------	--	-----

8				-					-		-	10	-		-	20		-		4	2.5	-	-	-	40		-		-	-	-					 277		2		2	-			. 13				 - 1				-	,
																-/-	*1																		 	 									Ι.							 	
e	-	٠		-				- 0	3	-		+6	(0)	-	-	*3		-	-	+	23	-			*	+	-	-	*			55		- 0			- 1	+	*3	•		-		6 9		0.0	•		*	*		-	٠

Suggest one reason why the calculated value of ΔH^{Θ} using Hess's Law in part (c) can (d) be considered accurate and one reason why it can be considered approximate.

[2]

Accur	rate:									
* * * *				 	 	* * * * *	 	 	 	
	· · · · · ·		• • •	 	 		 	 	 	
Appro	oximate) :								
	· · · · · ·			 	 		 	 	 	

Topic Chem 5 Q# 80/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q5/ www.SmashingScience.org :o)

5. This reaction is used in the manufacture of sulfuric acid.

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$
 $K_c = 280 \text{ at } 1000 \text{ K}$

$$K_c = 280 \text{ at } 1000 \text{ K}$$



_	(b)	Predict, giving your reason, the sign of the standard entropy change of the forward reaction.	[1]
	(c)	Calculate the standard Gibbs free energy change, ΔG^{\oplus} , in kJ, for this reaction at 1000 K. Use sections 1 and 2 of the data booklet.	[2]
	(d)	Predict, giving your reasons, whether the forward reaction is endothermic or exothermic. Use your answers to (b) and (c).	[2]
opic 4.	Prop	Q# 81/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q4/ www.SmashingScience.org :o) erties of elements and their compounds can be related to the position of the elements in periodic table.	
	(d)	Predict, giving your reasons, whether Mn ²⁺ or Fe ²⁺ is likely to have a more exothermic enthalpy of hydration.	[2]

Topic Chem 5 Q# 82/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q1/ www.SmashingScience.org :o)

3.26g of iron powder are added to 80.0 cm3 of 0.200 mol dm3 copper(II) sulfate solution. The 1. following reaction occurs:

$$Fe(s) + CuSO_4(aq) \rightarrow FeSO_4(aq) + Cu(s)$$



	(b)	(i)	The reaction was carried out in a calorimeter. The maximum temperature rise of the solution was 7.5 °C.	
			Calculate the enthalpy change, ΔH , of the reaction, in kJ, assuming that all the heat released was absorbed by the solution. Use sections 1 and 2 of the data booklet.	[2]
3				
	* */*			

<u> </u>		(ii)	State another assumption you made in (b)(i).	[1]
	• • •			
	# TS#5	* * * *		
L Tonic	Chem	5 O# 8	3/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q3/ www.SmashingScience.org :o)	
3.			tion is about ethene, C_2H_4 , and ethyne, C_2H_2 .	
	(c)	(i)	Under certain conditions, ethyne can be converted to benzene.	
			Determine the standard enthalpy change, ΔH^{Θ} , for the reaction stated, using section 11 of the data booklet.	[2]
			$3C_2H_2(g) \rightarrow C_6H_6(g)$	
	20,000			
1				

(ii)	Determine the standard enthalpy change, ΔH^{\oplus} , for the following similar reaction, using $\Delta H_{\rm f}$ values in section 12 of the data booklet.	[2]
	$3C_2H_2(g) \rightarrow C_6H_6(l)$	
* * * * * * *		
202222233		
* * * * * * * *		
(iii)	Explain, giving two reasons, the difference in the values for (c)(i) and (ii). If you did not obtain answers, use -475 kJ for (i) and -600 kJ for (ii).	[2]

(iv)	Calculate the standard entropy change, ΔS^9 , in J K ⁻¹ , for the reaction in (ii) using section 12 of the data booklet.	[1]
To the side of the		
	SMASHHULLI	
(v)	Determine, showing your working, the spontaneity of the reaction in (ii) at 25°C.	[3]
4444444		

* * * * * * * *		

Topic Chem 5 Q# 84/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q9/ www.SmashingScience.org :o)

9.	The Bomba	ardier beetle sprays a mixture of hydroquinone and hydrogen peroxide to fight off
	predators.	The reaction equation to produce the spray can be written as:

$$C_6H_4(OH)_2(aq) + H_2O_2(aq) \rightarrow C_6H_4O_2(aq) + 2H_2O(l)$$

hydroquinone quinone

(a)	(i)	Calculate the enthalpy	change, in kJ	for the spray reaction, us	ing the data below.	[2]
-----	-----	------------------------	---------------	----------------------------	---------------------	-----

$$C_8H_4(OH)_2(aq) \rightarrow C_8H_4O_2(aq) + H_2(g)$$

$$\Delta H^{\odot} = +177.0 \text{ kJ}$$

$$2H_2O(l) + O_2(g) \rightarrow 2H_2O_2(aq)$$

$$\Delta H^{\Theta} = +189.2 \text{ kJ}$$

$$H_2O(l) \rightarrow H_2(g) + \frac{1}{2}O_2(g)$$

$$\Delta H^{\odot} = +285.5 \,\text{kJ}$$

(ii) The energy released by the reaction of one mole of hydrogen peroxide with hydroquinone is used to heat 850 cm³ of water initially at 21.8°C. Determine the highest temperature reached by the water.

Specific heat capacity of water = 4.18 kJ kg⁻¹ K⁻¹.

(If you did not obtain an answer to part (i), use a value of 200.0 kJ for the energy released, although this is not the correct answer.)

***	**********



 (b) (i) Hydrogenation of propene produces propane. Calculate the standard entropy change, ΔS^Θ, for the hydrogenation of propene.

	_		_		_
-1	r	4	_	ı.	1
1	•			,	1
		- 5	g		1
			•		

Formula	Se / J K ⁻¹ mol ⁻¹
H ₂ (g)	+131
C ₃ H ₆ (g)	+267
C ₃ H ₈ (g)	+270

(ii)	The standard enthalpy change, ΔH^{\oplus} , for the hydrogenation of propene is $-124.4\mathrm{kJmol^{-1}}$. Predict the temperature above which the hydrogenation reaction is not spontaneous.

Topic Chem 5 Q# 85/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org :0)

4. Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.



(c) The reaction for the formation of liquid tetracarbonylnickel is shown below:

$$Ni(s) + 4CO(g) \rightarrow Ni(CO)_4(l)$$

Calculate the standard entropy change, ΔS^o, of the reaction, in J K⁻¹, using the values given.

[2]

	S ^e / J K ⁻¹ mol ⁻¹
Ni(s)	29.9
CO(g)	197.6
Ni(CO) ₄ (l)	313.4

21.00		,		•		•	•	 e Plet		*))	*	*:::	_			#0		•	((·		*	•	(*)		*	•		0.00	*	*) :*		*(0)		*.		*(1)	•		-		*	*00	-	-			•	
	009						. ,	 - 4	-				1			*	-						20					-	1			2	•31						-	-					 			٠	-
			•	٠		٠	•			*				•	č					•	7	-	(40)	•	*				1.0				•		*	÷	•			5,000	٠					•		•	•
					 	 (6)			*	400			-						 				**				 						*:03	 		*	*		 				# C		 -00-	 	116		

(ii) Calculate a value for ΔH^o in kJ.

[1]

	ΔH ^e _f / kJ mol ⁻¹
CO(g)	-110.5
Ni(CO) ₄ (l)	-633.0



		(iii)	Use your answers to (c)(i) and (c)(ii), to determine the temperature, in °C, at whethe decomposition of liquid tetracarbonylnickel to nickel and carbon monoxide becomes favourable.	ich
			(If you did not get answers to (c)(i) and (c)(ii), use $-500\mathrm{JK^{-1}}$ and $-200\mathrm{kJ}$ respectively but these are not the correct answers.)	[3]
		****	• • • • • • • • • • • • • • • • • • • •	***

				100
	(d)	Suc	gest why experiments involving tetracarbonylnickel are very hazardous.	[1]
	(u)	Sug	gest why experiments involving terracarbonymicker are very nazardous.	ניו
opi	c Chem	5 Q#	86/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/	
1.			2-diol, HOCH ₂ CH ₂ OH, has a wide variety of uses including the removal of ice from d heat transfer in a solar cell.	
	(a)	Etha	ne-1,2-diol can be formed according to the following reaction.	
			$2CO(g) + 3H_2(g) \Longrightarrow HOCH_2CH_2OH(g)$	
		(iii)	Calculate the enthalpy change, ΔH^{Θ} , in kJ, for this reaction using section 11 of the data booklet. The bond enthalpy of the carbon–oxygen bond in CO(g) is	[3]
			1077 kJ mol⁻¹.	[0]

	* * *			
	* * *			
2				



(b) (i) Calculate ΔA ⁻¹ , In kJ, for this similar reaction below using ΔA ⁻¹ data from section 12 of the data booklet. ΔH ⁰ of HOCH ₂ CH ₂ OH (l) is −454.8 kJ mol ⁻¹ .	[1]
$2CO(g) + 3H_2(g) \rightleftharpoons HOCH_2CH_2OH(l)$	
(ii) Deduce why the answers to (a)(iii) and (b)(i) differ.	[1]
(iii) ΔS^{\oplus} for the reaction in (b)(i) is -620.1 J K ⁻¹ . Comment on the decrease in entropy.	[1]
 (iv) Calculate the value of ΔG^e, in kJ, for this reaction at 298 K using your answer to (b)(i). (If you did not obtain an answer to (b)(i), use −244.0 kJ, but this is not the correct value.) 	[2]
 (v) Comment on the statement that the reaction becomes less spontaneous as temperature is increased. 	[1]

Topic Chem 5 Q# 87,	/ IB Chem/2016/s/TZoSP/Pape	r 2 Section A/Higher Level/Q6/	www.SmashingScience.org:o)
---------------------	-----------------------------	--------------------------------	----------------------------

Ozone in the atmosphere can be formed from the combustion of methane. State the equation for this combustion reaction. [1] (ii) Calculate the standard enthalpy change for the reaction, ΔH^Θ, in kJmol⁻¹, using the thermodynamic data in section 12 of the data booklet and the information [1] given below. $O_3(g)$ $\Delta H_f^{\Theta} = +142.3 \,\mathrm{kJ}\,\mathrm{mol}^{-1}$ (iii) State why the standard enthalpy change of formation, ΔH_f^{Θ} , for oxygen is not given. [1]

(c)

(iv) Calculate the standard entropy change for the reaction, ΔS^Θ, in JK⁻¹mol⁻¹, using the thermodynamic data in section 12 of the data booklet and the information given below.

[1]

O ₂ (g)	O ₃ (g)
$S^{\Theta} = +205.0 \mathrm{J K^{-1} mol^{-1}}$	$S^{\Theta} = +237.6 \text{JK}^{-1} \text{mol}^{-1}$

.....

(v) Deduce the standard change in Gibbs Free Energy, ΔG^Θ, in kJmol⁻¹, for this reaction at 298 K.

[1]

(vi) Deduce, giving a reason, whether the reaction is spontaneous or non-spontaneous at this temperature.

[1]


Topic Chem 5 Q# 88/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/

- One of the main constituents of acid deposition is sulfuric acid, H₂SO₄. This acid is formed from the sulfur dioxide pollutant, SO₂.
 - (b) Consider the following equilibrium between the two oxides of sulfur, sulfur dioxide and sulfur trioxide:

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$
 $\Delta H = -198 \text{ kJ}$



(c) Sketch the potential energy profile for the forward reaction in part (b) to show the effect of a catalyst on the activation energy, $E_{\rm act}$.



Topic Chem 5 Q# 89/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/

- 5. Many automobile manufacturers are developing vehicles that use hydrogen as a fuel.
 - (a) Suggest why such vehicles are considered to cause less harm to the environment than those with internal combustion engines.

[1]




(b) Hydrogen can react with ethene to form ethane.

$$H_2(g) + C_2H_4(g) \rightarrow C_2H_6(g)$$


Using average bond enthalpies at 298 K from section 11 of the data booklet, calculate the change in enthalpy, ΔH , in kJ mol⁻¹, for this reaction.

[3]



Topic Chem 5 Q# 90/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q3/ www.SmashingScience.org:o)

(b) The enthalpy change for the reaction between nitrogen monoxide and hydrogen is -664 kJ and its activation energy is 63 kJ.





Topic Chem 5 Q# 92/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/

4. Tin(II) chloride is a white solid that is commonly used as a reducing agent.

	(i)	State why you would expect tin(II) chloride to have a similar lattice enthalpy to strontium chloride, using section 9 of the data booklet.	[1
	(ii)	Calculate the molar enthalpy change when strontium chloride is dissolved in water, using sections 18 and 20 of the data booklet.	[2

cl	041	22 / ID Character / 774 / Day 22 Carrier A / IV about 1	
Phosg	jene	O3/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/ COCl ₂ , is usually produced by the reaction between carbon monoxide and	
cniorir	ne ac	cording to the equation:	
cniorir	ne ac		
CNIONI	ne ad	ccording to the equation: $CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$	
	ne ad	cording to the equation:	[2
		coording to the equation: $CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$ The standard enthalpy change of formation of phosgene, ΔH_r^{Θ} , is $-220.1 \text{kJ} \text{mo} \Gamma^1$. Determine the standard enthalpy change, ΔH_r^{Θ} , for the forward reaction of the	[2
		coording to the equation: $CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$ The standard enthalpy change of formation of phosgene, ΔH_r^{Θ} , is $-220.1 \text{kJ} \text{mo} \Gamma^1$. Determine the standard enthalpy change, ΔH_r^{Θ} , for the forward reaction of the	[2
		coording to the equation: $CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$ The standard enthalpy change of formation of phosgene, ΔH_r^{Θ} , is $-220.1\mathrm{kJmor^1}$. Determine the standard enthalpy change, ΔH_r^{Θ} , for the forward reaction of the equilibrium, in kJ, using section 12 of the data booklet.	[2
		coording to the equation: $CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$ The standard enthalpy change of formation of phosgene, ΔH_r^{Θ} , is $-220.1 \text{kJ} \text{mo} \Gamma^1$. Determine the standard enthalpy change, ΔH_r^{Θ} , for the forward reaction of the	[2
		coording to the equation: $CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$ The standard enthalpy change of formation of phosgene, ΔH_r^{Θ} , is $-220.1\mathrm{kJmor^1}$. Determine the standard enthalpy change, ΔH_r^{Θ} , for the forward reaction of the equilibrium, in kJ, using section 12 of the data booklet.	[2



(iv) Calculate the standard entropy change, ΔS[⊕], in JK⁻¹, for the forward reaction at 25°C, using your answers to (a) (ii) and (a) (iii).
 (If you did not obtain an answer to (a) (ii) and/or (a) (iii) use values of +20.0 kJ and -120.0 kJ respectively, although these are not the correct answers.)

[2]

1049	0.00			5.5	0.50		119	2117	221.0		200	100			2.5		00110								000	15:35		51.17	223		312	
	50																															
	 •							-										•	•						•					•	-	
							1			1		* .												 •		0.0	-	7. 5				
***				5.37		*								1			0.15			* *							-	-	× .		-	3.5
****		30				13.0					100	* *		1.7		7.5					7.7	2.5	-		 * *			37			30	12.

Topic Chem 5 Q# 94/ IB CHEM/2015/w/TZ0/Paper 2 Section B/Higher Level/

- Iron (III) oxide is the main source of iron but the decomposition of Fe₂O₃(s) into its elements
 is extremely difficult due to a large positive value of ΔG^Θ.
 - (a) Consider the following reactions:

Fe₂O₃(s)
$$\rightarrow$$
 2Fe(s) + $\frac{3}{2}$ O₂(g) $\Delta G^{\oplus} = +742 \text{ kJ mol}^{-1}$
CO(g) + $\frac{1}{2}$ O₂(g) \rightarrow CO₂(g) $\Delta G^{\oplus} = -257 \text{ kJ mol}^{-1}$

Suggest, with a reason, whether it is possible to produce iron by reacting Fe₂O₃ with CO. [2]

* * * * * * * * * * * * * * * * * * *	**********

(b) The thermite reaction is one of the most exothermic reactions.

$$Fe_2O_3(s) + 2Al(s) \rightarrow 2Fe(l) + Al_2O_3(s)$$
 $\Delta H^{\Theta} = -825.2 \text{ kJ}$

Species	S [®] / JK ⁻¹ mol ⁻¹	∆G°/kJmol
Al(s)	+28.3	0
Al ₂ O ₃ (S)	+50.9	-1582
Fe(l)	+34.8	+10.0
Fe ₂ O ₃ (s)	+87.5	-742



	 (i) Calculate the standard free energy change of for the standard free energy change of for 	ange, ΔG^{Θ} , in kJ mo Γ^1 , by using values of rmation, ΔG^{Θ}_{f} , from the table above.	[2
			Ī

(ii)	Calculate the standard entropy change, $\Delta S'$ standard entropy, S^{Θ} , from the table.	e, in JK ⁻¹ mol ⁻¹ , by using values of [1]	
/iiii	Calculate the standard free energy change	AC® for the reaction using AU® and	
(iii)	Calculate the standard free energy change, ΔS ^e values at 25°C.	[2]	

hem	******************************		rg
Prop	******************************	A/Higher Level/Table 11/ www.SmashingScience.or	g
Prop	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section ana. C ₃ H ₈ (g), undergoes complete combustion	A/Higher Level/Table 11/ www.SmashingScience.or	rg [1]
Prop wate	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section of pane, C ₃ H ₈ (g), undergoes complete combustions, H ₂ O(g).	A/Higher Level/Table 11/ www.SmashingScience.or	
Prop wate	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section of pane, C ₃ H ₈ (g), undergoes complete combustions, H ₂ O(g).	A/Higher Level/Table 11/ www.SmashingScience.or	
Prop wate	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section Apane, C ₃ H ₈ (g), undergoes complete combustions, H ₂ O(g). State an equation for the complete combustions of the complete combustions of the complete combustions.	A/Higher Level/Table 11/ www.SmashingScience.or on to form carbon dioxide, $CO_2(g)$, and on of propane, $C_3H_8(g)$.	[1]
Prop wate (a)	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section Apane, C ₃ H ₈ (g), undergoes complete combustions, H ₂ O(g). State an equation for the complete combustions	A/Higher Level/Table 11/ www.SmashingScience.or on to form carbon dioxide, $CO_2(g)$, and on of propane, $C_3H_8(g)$.	
Prop wate (a)	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section Apane, C ₃ H ₈ (g), undergoes complete combustions, H ₂ O(g). State an equation for the complete combustions of the complete combustions of the complete combustions.	A/Higher Level/Table 11/ www.SmashingScience.or on to form carbon dioxide, $CO_2(g)$, and on of propane, $C_3H_8(g)$.	[1]
Prop wate (a)	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section Apane, C ₃ H ₈ (g), undergoes complete combustions, H ₂ O(g). State an equation for the complete combustions of the complete combustions of the complete combustions.	A/Higher Level/Table 11/ www.SmashingScience.or on to form carbon dioxide, $CO_2(g)$, and on of propane, $C_3H_8(g)$.	[1]
Prop wate (a)	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section Apane, C ₃ H ₈ (g), undergoes complete combustions, H ₂ O(g). State an equation for the complete combustions of the complete combustions of the complete combustions.	A/Higher Level/Table 11/ www.SmashingScience.or on to form carbon dioxide, $CO_2(g)$, and on of propane, $C_3H_8(g)$.	[1]
Prop wate (a)	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section Apane, C ₃ H ₈ (g), undergoes complete combustions, H ₂ O(g). State an equation for the complete combustions of the complete combustions of the complete combustions.	A/Higher Level/Table 11/ www.SmashingScience.or on to form carbon dioxide, $CO_2(g)$, and on of propane, $C_3H_8(g)$.	[1]
Prop wate (a)	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section Apane, C ₃ H ₈ (g), undergoes complete combustions, H ₂ O(g). State an equation for the complete combustions of the complete combustions of the complete combustions.	A/Higher Level/Table 11/ www.SmashingScience.or on to form carbon dioxide, $CO_2(g)$, and on of propane, $C_3H_8(g)$.	[1]
Prop wate (a)	5 Q# 95/ IB Chem/2015/w/TZ0/Paper 2 Section Apane, C ₃ H ₈ (g), undergoes complete combustions, H ₂ O(g). State an equation for the complete combustions of the complete combustions of the complete combustions.	A/Higher Level/Table 11/ www.SmashingScience.or on to form carbon dioxide, $CO_2(g)$, and on of propane, $C_3H_8(g)$.	[1]

Topic Chem 5 Q# 96/ IB Chem/2015/s/TZ2/Paper 2 Section A/Higher Level/

Carbon monoxide reacts with hydrogen to produce methanol.

$$CO(g) + 2H_2(g) \rightarrow CH_3OH(l)$$

Substance	ΔH ^O _f / kJ mol ⁻¹	ΔG ^O _f / kJ mol ⁻¹	S ^O / J mol ⁻¹ K ⁻¹
CO (g)	-110.5	- 137.2	+ 197.6
CH₃OH(l)	-239.0	- 166.0	+ 126.8


(a)	Calculate the standard enthalpy change, ΔH° , in kJ mol ⁻¹ , for the reaction.	[1]
(b)	Calculate the standard free energy change, ΔG^{\ominus} , in kJ mol ⁻¹ , for the reaction $(\Delta G_{f}^{\ominus}(H_{2}) = 0 \text{ kJ mol}^{-1})$.	[1]
(c)	Using the values obtained in parts (a) and (b), calculate the standard entropy change, ΔS^{Θ} , in Jmol ⁻¹ K ⁻¹ , for the reaction at 298 K.	[1]
(d)	Determine the absolute entropy, S [⊕] , in Jmol ⁻¹ K ⁻¹ , for H ₂ (g) at 298 K.	[2]



Topic Chem 5 Q# 97/ IB Chem/2015/s/TZ2/Paper 2 Section A/Higher Level/

 A student carried out an experiment to determine the concentration of a hydrochloric acid solution and the enthalpy change of the reaction between aqueous sodium hydroxide and this acid by thermometric titration.

She added 5.0 cm³ portions of hydrochloric acid to 25.0 cm³ of 1.00 mol dm⁻³ sodium hydroxide solution in a glass beaker until the total volume of acid added was 50.0 cm³, measuring the temperature of the mixture each time. Her results are plotted in the graph below.



The initial temperature of both solutions was the same.

(a)	(i)	By drawing appropriate lines, determine the volume of hydrochloric acid required	
	**	to completely neutralize the 25.0 cm³ of sodium hydroxide solution.	[2]



	Determine the concentration of the hydrochloric acid, including units.	[2
(i)	Determine the change in temperature, ΔT .	[1]
(ii)	Calculate the enthalpy change, in kJ mol ⁻¹ , for the reaction of hydrochloric acid and sodium hydroxide solution.	[3]
(iii	The accepted theoretical value from the literature of this enthalpy change is -58 kJ mol ⁻¹ . Calculate the percentage error correct to two significant figures.	[1]
(iii		[1]
(iii		[1]
	–58 kJ mol ⁻¹ . Calculate the percentage error correct to two significant figures. Suggest the major source of error in the experimental procedure and an	
	–58 kJ mol ⁻¹ . Calculate the percentage error correct to two significant figures. Suggest the major source of error in the experimental procedure and an	
	–58 kJ mol ⁻¹ . Calculate the percentage error correct to two significant figures. Suggest the major source of error in the experimental procedure and an	



Na	halpy	Born-Haber cycle and tables 7, 10 and 13 of the data booklet. The standard change of atomization (standard enthalpy change of sublimation), ΔH_{at}^{Θ} , of $+108 \mathrm{kJ}\mathrm{mol}^{-1}$. [4]	ı
	* *		1
* *			
*			
* *	1 2 2 2 2		
* *			
+ +			
	11]
c Chem (b)	15 Q# (i)	99/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q5/ www.SmashingScience.org: Define the term average bond enthalpy.	
(0)	(1)	Define the term average bond entitlalpy.	[2]
		AND STATES	
	(ii)		[4]
	(ii)	Ethanol can be used as a fuel. Determine the enthalpy of combustion of ethanol at 298 K, in kJ mol ⁻¹ , using the values in table 10 of the data booklet, assuming all	[4
	(ii)	Ethanol can be used as a fuel. Determine the enthalpy of combustion of ethanol at 298 K, in kJ mol ⁻¹ , using the values in table 10 of the data booklet, assuming all	[4
	(ii)	Ethanol can be used as a fuel. Determine the enthalpy of combustion of ethanol at 298 K, in kJ mol ⁻¹ , using the values in table 10 of the data booklet, assuming all	[4]
	(ii)	Ethanol can be used as a fuel. Determine the enthalpy of combustion of ethanol at 298 K, in kJ mol ⁻¹ , using the values in table 10 of the data booklet, assuming all reactants and products are gaseous.	[4]
	(ii)	Ethanol can be used as a fuel. Determine the enthalpy of combustion of ethanol at 298 K, in kJ mol ⁻¹ , using the values in table 10 of the data booklet, assuming all	[4]
	(ii)	Ethanol can be used as a fuel. Determine the enthalpy of combustion of ethanol at 298 K, in kJ mol ⁻¹ , using the values in table 10 of the data booklet, assuming all reactants and products are gaseous.	[4]
	(ii)	Ethanol can be used as a fuel. Determine the enthalpy of combustion of ethanol at 298 K, in kJ mol ⁻¹ , using the values in table 10 of the data booklet, assuming all reactants and products are gaseous.	[4]
	(ii)	Ethanol can be used as a fuel. Determine the enthalpy of combustion of ethanol at 298 K, in kJ mol ⁻¹ , using the values in table 10 of the data booklet, assuming all reactants and products are gaseous.	[4]
	(ii)	Ethanol can be used as a fuel. Determine the enthalpy of combustion of ethanol at 298 K, in kJ mol ⁻¹ , using the values in table 10 of the data booklet, assuming all reactants and products are gaseous.	[4]

(f)	(i)	The standard enthalpy change of combustion, ΔH_c^{\ominus} , of propanoic acid is
		-1527 kJ mol ⁻¹ . Determine the standard enthalpy change of formation of
		propanoic acid, in kJ mol-1, using this information and data from table 12 of the
		data booklet.

			 		-8					2 .																		 									
* *	4	*	•	*		*	*	•	*	•		(A)		-			•	-		*5	 *	*	*	W. (*				*	*				*	
																																			•		
		•																								•							 				
			 ٠	+											~					*10	 *	**		40)								*					
															3				٠		 -								٠								
-:	*	•		÷	7.0		0	*		-	7	7			. 7	7	30			*	2.5	200	•		-	200			-	 7		+		to	7.0		
																	- 1			*																	

(ii) Deduce, giving a reason, the sign of the standard entropy change of the system for the formation of propanoic acid from its elements.

[2]



Topic Chem 6 Q# 100/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org:o)

- 6. Copper forms two chlorides, copper(I) chloride and copper(II) chloride.
 - (ii) Copper(II) chloride is used as a catalyst in the production of chlorine from hydrogen chloride.


$$4HCl(g) + O_2(g) \rightarrow 2Cl_2(g) + 2H_2O(g)$$



(iii) The diagram shows the Maxwell–Boltzmann distribution and potential energy profile for the reaction without a catalyst.

Annotate both charts to show the activation energy for the catalysed reaction, using the label $E_{\rm a\,(cat)}$.

[2]



(iv) Explain how the catalyst increases the rate of the reaction.

[2]

Topic Chem 6 Q# 101/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q4./ www.SmashingScience.org:o)

 Propene is an important starting material for many products. The following shows some compounds which can be made from propene, C₃H₆.


Propene (C_3H_8) $\rightarrow C_3H_7Cl \rightarrow C_3H_8O \rightarrow C_3H_8O$



(b) An experiment was carried out to determine the order of reaction between one of the isomers of C₃H₇Cl and aqueous sodium hydroxide. The following results were obtained.

Experiment	[C ₃ H ₇ CI] / mol dm ⁻³	[OHT] / mol dm ⁻³	Initial rate / mol dm ⁻³ s ⁻¹
1	0.05	0.10	3.1 × 10 ⁻⁴
2	0.10	0.20	1.3 × 10 ⁻³
3	0.15	0.10	9.2 × 10 ⁻⁴

(i)	Determine	the rate expression	from the results,	explaining your method.	[3]
-----	-----------	---------------------	-------------------	-------------------------	-----



Topic Chem 6 Q# 102/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q10/ www.SmashingScience.org:o)

10. The following mechanism is proposed for a reaction:

$$A+B \rightarrow C+D$$
 slow step
 $D+B \rightarrow A+E$ fast step

(a) Classify substances B and D as reactant, product, catalyst, or intermediate, based on the proposed mechanism.

B: D:

(b) Deduce the rate expression. [1]

(c) Calculate the initial rate of reaction for experiment 2, if measured under the same conditions.

[1]

Experiment	[A] / moldm ⁻³	[B] / mol dm ⁻³	Initial rate / moldm ⁻³ s ⁻¹
1	0.200	0.200	1.20
2	0.300	0.200	


Topic Chem 6 Q# 103/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o)

 3.26 g of iron powder are added to 80.0 cm³ of 0.200 mol dm⁻³ copper(II) sulfate solution. The following reaction occurs:

$$Fe(s) + CuSO_4(aq) \rightarrow FeSO_4(aq) + Cu(s)$$

(c) (i) Sketch a graph of the concentration of iron(II) sulfate, FeSO₄, against time as the reaction proceeds.

[2]



(ii) Outline how the initial rate of reaction can be determined from the graph in part (c)(i).

[2]

* ** ****	 	
* * * * * * * * * * * * * * * * * * * *	 	

(iii) Explain, using the collision theory, why replacing the iron powder with a piece of iron of the same mass slows down the rate of the reaction.

Calcium carbonate reacts with hydrochloric acid.


$$CaCO_3(s) + 2HCl(aq) \rightarrow CaCl_2(aq) + H_2O(l) + CO_2(g)$$

(a) Outline two ways in which the progress of the reaction can be monitored. No practical details are required.

[2]

* * *	• •	+ - +	 •	• •		•		*	٠		6 24	٠		*	+	*	• •	e) (e	•	*	•	٠	*		•	•		٠		*		*		 . *	•	 	*	*	•	 		*		 •	+	
			 -	1-2		S.		-					-		-				- 2				-		- 1	÷	 +	+		-			_	 -				*		 		-	-			
	• •			• •		 •	• •			•	•				•					•		•		• •		•	 	•	• •	*3	٠.		•				•	*				•			٠	
			 •	• •)÷									٠	•	•	•		٠	*		 ٠			•			•		*		•		 *		 	٠			 	•			 	٠	

(b) The results of a series of experiments in which the concentration of HCl was varied are shown below.





(i) Suggest why point D is so far o	ut of line assuming human error is not the cause. [1	1]
(ii) Draw the best fit line for the rea	action excluding point D. [1	1]
(iii) Suggest the relationship that po of the acid and the rate of reach	oints A, B and C show between the concentration [1	1]
(iv) Deduce the rate expression for	the reaction.	1]
(v) Calculate the rate constant of the	he reaction, stating its units. [2	2]
(c) Predict from your line of best fit the ra	ate of reaction when the concentration of HCl is	1]




(d) De	escribe h	ow the a	activation	energy	of this	reaction	could	be determined	I.
--------	-----------	----------	------------	--------	---------	----------	-------	---------------	----

Topic Chem 6 Q# 105/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q5/ www.SmashingScience.org:o)

5. (a) Magnesium reacts with sulfuric acid:

$$Mg(s) + H_2SO_4(aq) \rightarrow MgSO_4(aq) + H_2(g)$$

The graph shows the results of an experiment using excess magnesium ribbon and dilute sulfuric acid.



Outline why the rate of the reaction decreases with time.

(ii) Sketch, on the same graph, the expected results if the experiment were repeated using powdered magnesium, keeping its mass and all other variables unchanged. [1]

[1]

[3]

(b) Nitrogen dioxide and carbon monoxide react according to the following equation:

$$NO_2(g) + CO(g) \rightleftharpoons NO(g) + CO_2(g)$$

$$\Delta H = -226 \text{ kJ}$$

Experimental data shows the reaction is second order with respect to NO_2 and zero order with respect to CO.

(i) State the rate expression for the reaction.

[1]

(ii) The following mechanism is proposed for the reaction.

Step I
$$NO_2(g) + NO_2(g) \rightarrow NO(g) + NO_3(g)$$

Step II
$$NO_3(g) + CO(g) \rightarrow NO_2(g) + CO_2(g)$$

Identify the rate determining step giving your reason.


[1]

(iii) State one method that can be used to measure the rate for this reaction.

[1]

(iv) Calculate the activation energy for the reverse reaction.

[1]




Reaction coordinate

(V	Sketch the relationship between the rate of reaction and the concentration of NC	
	Ĩ	
	Rate	
	[NO ₂]	
	he Arrhenius equation, $k = Ae^{-\frac{E_a}{RT}}$, gives the relationship between the rate constant nd temperature.	
	tate how temperature affects activation energy.	
222234		* * ·
(d) S	tate the equation for the reaction of NO ₂ in the atmosphere to produce cid deposition.	2.2
(d) S	tate the equation for the reaction of NO_2 in the atmosphere to produce	
(d) S	tate the equation for the reaction of NO_2 in the atmosphere to produce	
(d) Si ad	tate the equation for the reaction of NO_2 in the atmosphere to produce	
(d) Si ad	tate the equation for the reaction of NO ₂ in the atmosphere to produce cid deposition. 106/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o) of the acid-catalysed iodination of propanone can be followed by measuring how the	**
(d) Si ad	tate the equation for the reaction of NO ₂ in the atmosphere to produce cid deposition. 106/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o) of the acid-catalysed iodination of propanone can be followed by measuring how the ation of iodine changes with time.	
(d) Si ad	tate the equation for the reaction of NO_2 in the atmosphere to produce cid deposition. 106/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o) of the acid-catalysed iodination of propanone can be followed by measuring how the ation of iodine changes with time. $I_2(aq) + CH_3COCH_3(aq) \rightarrow CH_3COCH_2I(aq) + H^+(aq) + I^-(aq)$	
(d) Si ad	tate the equation for the reaction of NO_2 in the atmosphere to produce cid deposition. 106/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o) of the acid-catalysed iodination of propanone can be followed by measuring how the ation of iodine changes with time. $I_2(aq) + CH_3COCH_3(aq) \rightarrow CH_3COCH_2I(aq) + H^+(aq) + I^-(aq)$ and form of the rate equation is:	

(ii) A student produced these results with [H⁺] = 0.15 mol dm⁻³. Propanone and acid were in excess and iodine was the limiting reagent.

Determine the relative rate of reaction when $[H^{+}] = 0.15 \,\text{mol dm}^{-3}$.

[2]




(b) The student then carried out the experiment at other acid concentrations with all other conditions remaining unchanged.

[H ⁺] / mol dm ⁻³	Relative rate of reaction
0.05	0.0025
0.10	0.0051
0.20	0.0100

Determine the relationship between the rate of reaction and the concentration of acid and the order of reaction with respect to hydrogen ions.

Re	ela	itic	on	sł	ip):																																																						
					•					ं	(30)	•					•	-		•				•	÷			•		+	÷			+		3	•	٠			•		•	٠	٠	-	÷		•		•			•						
505				-	•	91.0	505		15		ž.		•	***		*		-10	*	-	÷	-	-	*	7	-	-		7.	-	Ť	50		•			12	-	ti	-	-53	-	 2.5		-		-	-	-	•	•	-	•	35	-	-	•	+	-	
Or	rde	er	of	re	ea	iC	tic	ıc	n	V	/i	th	ı	e	s	p	e	C	t	to)	[]	1	†]:																																				

(c) When the concentration of iodine is varied, while keeping the concentrations of acid and propanone constant, the following graphs are obtained.




Deduce, giving your reason, the order of reaction with respect to iodine.


Order with respect to iodine	
Reason:	



(d) When the reaction is carried out in the absence of acid the following graph is obtained.



Discuss the shape of the graph between A and B.





Topic Chem 6 Q# 107/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/

Sodium thiosulfate solution reacts with dilute hydrochloric acid to form a precipitate of sulfur at room temperature.

$$Na_2S_2O_3(aq) + 2HCl(aq) \rightarrow S(s) + SO_2(g) + 2NaCl(aq) + X$$

(a) Identify the formula and state symbol of X.

	• •																					


(b) Suggest why the experiment should be carried out in a fume hood or in a well-ventilated laboratory.

[1]

[1]



(c) The precipitate of sulfur makes the mixture cloudy, so a mark underneath the reaction mixture becomes invisible with time.



10.0 cm³ of 2.00 mol dm⁻³ hydrochloric acid was added to a 50.0 cm³ solution of sodium thiosulfate at temperature, T₁. Students measured the time taken for the mark to be no longer visible to the naked eye. The experiment was repeated at different concentrations of sodium thiosulfate.


Experiment	[Na ₂ S ₂ O ₃ (aq)] / mol dm ⁻³	Time, t, for mark to disappear / s ± 1 s	$\frac{1}{t}^* / 10^{-3} s^{-1}$
1	0.150	23	43.5
2	0.120	27	37.0
3	0.090	36	27.8
4	0.060	60	16.7
5	0.030	111	9.0

^{*} The reciprocal of the time in seconds can be used as a measure of the rate of reaction.

[Source: Adapted from http://www.flinnsci.com/]



											2,5								10.00
							12.21		\$ 5		, L , L ,					S 212		2 20, 20	
		1																	
(d) Draw the best fit axes provided.	line o	of -	ag	ains	st c	onc	ent	rati	on	of s	sodi	un	n thi	osul	fate	on	the		
•					_							_							
	45-	1																	
	45-							F											
	40-																		
τ.	35-							l											
S . 0	30-																		
Rate of reaction $\left(\frac{1}{t}\right)/10^{-3}s^{-1}$																			
	25-																		
actic	20-				d			h	E										
of re	15-							P											
Rate	10-																		
- (1					
	5-			H															
	_			0.03	2	-	00		0	00		0	10	0	15)			
	0		4	U.U.			.06			09 nole		0.1	12	U	.15				



	 In a different experiment, this reaction was found to be first order with respect to hydrochloric acid. Deduce the overall rate expression for the reaction. 	[1]
(f)	A student decided to carry out another experiment using 0.075 mol dm ⁻³ solution of sodium thiosulfate under the same conditions. Determine the time taken for the mark to be no longer visible.	[2]
 (g)	 An additional experiment was carried out at a higher temperature, T₂. (i) On the same axes, sketch Maxwell–Boltzmann energy distribution curves at the two temperatures T₁ and T₂, where T₂ > T₁. 	[2]
	SMASHING 1 1 1	

	(ii)	_	хрісіі	,				ses the r					Į.
		• • •/•											
10.0						* * * * * *				•••••			
(h)				reason may be			of rates	of react	ions obta	ained at	higher		ĺ
						* * * * * * *							
								on A/Hig		tial) fact	or, A.		[1]
											or, A.		[1]
T1 87	he act	e wh	on en	ergy, E_a	for the	e reaction	on of 1-nential)	odoetha	exponen	sodium	hydroxid ol ⁻¹ dm ³ s	-1	[1]
TI 87	he acti	ivati	on en	ergy, E_a	for the	e reaction	on of 1-nential)	odoetha	exponen	sodium	hydroxid	-1	[2]
TI 87	he acti	ivati	on en	ergy, E ₃ , the frequence	for the	e reaction	on of 1-nential)	odoetha	exponen	sodium	hydroxid ol ⁻¹ dm ³ s	-1	
TI 87	he acti	ivati	on en	ergy, E ₃ , the frequence	for the	e reaction	on of 1-nential)	odoetha	exponen	sodium	hydroxid ol ⁻¹ dm ³ s	-1	
TI 87	he acti	ivati	on en	ergy, E ₃ , the frequence	for the	e reaction	on of 1-nential)	odoetha	exponen	sodium	hydroxid ol ⁻¹ dm ³ s	-1	
TI 87	he acti	ivati	on en	ergy, E ₃ , the frequence	for the	e reaction	on of 1-nential)	odoetha	exponen	sodium	hydroxid ol ⁻¹ dm ³ s	-1	
TI 87	he acti	ivati	on en	ergy, E ₃ , the frequence	for the	e reaction	on of 1-nential)	odoetha	exponen	sodium	hydroxid ol ⁻¹ dm ³ s	-1	
TI 87	he acti	ivati	on en	ergy, E ₃ , the frequence	for the	e reaction	on of 1-nential)	odoetha	exponen	sodium	hydroxid ol ⁻¹ dm ³ s	-1	



Topic Chem 6 Q# 109/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/

The reaction between hydrogen and nitrogen monoxide is thought to proceed by the mechanism shown below.

$2NO(g) \rightleftharpoons N_2O_2(g)$	fast equilibrium
$N_2O_2(g) + H_2(g) \rightarrow N_2O(g) + H_2O(g)$	slow reaction
$N_2O(g) + H_2(g) \rightarrow N_2(g) + H_2O(g)$	fast reaction

		1.20 (9) 1.2(9) 1.20(9)	
(a)	(i)	State the equation for the overall reaction.	[1]
((e)(e) e			****
			* * * * *
	(ii)	Deduce the rate expression consistent with this mechanism.	[1]
	(iii)	Explain how you would attempt to confirm this rate expression, giving the	results

you would expect.	[3]
***************************************	* * * * * * * *



i	(iv)	State, giving your reason, whether confirmation of the rate expression would prove that the mechanism given is correct.	[1]
	(v)	Suggest how the rate of this reaction could be measured experimentally.	[1]
b)	(iii)	Sketch and label a second Maxwell–Boltzmann energy distribution curve representing the same system but at a higher temperature, T _{higher} .	[1]
	Probability of this energy	Energy Energy	
	(iv)	Explain why an increase in temperature increases the rate of this reaction.	[2]

Topic Chem 6 Q# 110/ IB CHEM/2015/w/TZ0/Paper 2 Section B/Higher Level/Q7/ www.SmashingScience.org :o)

(c) The oxidation of nitrogen monoxide takes place as follows:

$$2NO(g) + O_2(g) \rightarrow 2NO_2(g)$$

The following experimental data was obtained at 101.3 kPa and 298 K.

Experiment	Initial [NO] / mol dm ⁻³	Initial [O ₂] / mol dm ⁻³	Initial rate / mol dm ⁻³ s ⁻¹
1	2.30 × 10 ⁻²	1.15×10^{-2}	1.05 × 10 ⁻³
2	2.30 × 10 ⁻²	2.30 × 10 ⁻²	2.09 × 10 ⁻³
3	4.60 × 10 ⁻²	4.60 × 10 ⁻²	1.68 × 10 ⁻²

[1]
[2]
*



							44 5 5 5 5 5 5
* * *							
1.1.1	**********		*******	na nama sa s			halada na
* * *							
* * *		********					
Chem 6 Q#	111/ IB Chem/	/2015/w/TZ0	/Paper 2 Sec	tion A/Highe	er Level/Q1/	www.Smashi	ngScience.c
The studer	nt found by fu	irther experi	mentation th	hat oxidation	n of ascorbi	c acid follow	s
first-order	kinetics. The	graph of In	k against 1	is shown be	elow.		
	the activation		1			units	[3
200000000					,		
			$\frac{1}{\tau}$	/ K-1			
		0.0004	0.0032	0.0033	0.0034	0.0035	0.0036
	0.0030	0.0031					
7	0.0030	0.0031	0.0032		111111111	0.0033	0.0036
	0.0030	0.0031	0.0032			0.0033	0.0036
-10.4	0.0030	0.0031	0.0032			0.0055	0.0036
-10.4	0.0030	0.0031	0.0032			0.0030	0.0036
-10.5	0.0030	0.0031	0.0032			0.0030	0.0036
-10.5	0.0030	0.0031	0.0032			0.0050	0.0036
-10.5	0.0030	0.0031	0.0032			0.0050	0.0036
-10.5	0.0030	0.0031	0.0032			0.0050	0.0036
-10.5	0.0030	0.0031	0.0032			0.0030	0.0036
-10.5 -10.6 -10.7 -10.8	0.0030	0.0031	0.0032			0.0050	0.0036
-10.5 -10.6 -10.7	0.0030	0.0031	0.0032			0.0033	0.0036
-10.5 -10.6 -10.7 -10.8 -10.9	0.0030	0.0031	0.0032			0.0033	0.0036
-10.5 -10.6 -10.7 -10.8	0.0030	0.0031	0.0032			0.0055	0.0036
-10.5 -10.6 -10.7 -10.8 -10.9	0.0030	0.0031	0.0032				0.0036
-10.5 -10.6 -10.7 -10.8 -10.9	0.0030						0.0036
-10.5 -10.6 -10.7 -10.8 -10.9	0.0030	0.0031					0.0036

2.

(a)	Define the term rate of reaction.	[1]
(b)	Explain why increasing the particle size of a solid reactant decreases the rate of reaction.	[2]
(c)	Nitrogen(II) oxide reacts with hydrogen according to the equation below.	
	$2NO(g) + 2H_2(g) \rightarrow N_2(g) + 2H_2O(g)$ A suggested mechanism for this reaction is:	
	Step 1: $NO + H_2 \rightleftharpoons X$ fast	
	Step 2: $X + NO \rightarrow Y + H_2O$ slow	
	Step 3: $Y + H_2 \rightarrow N_2 + H_2O$ fast	
	(i) Identify the rate-determining step.	[1]

	 (ii) A student hypothesized that the order of reaction with respect to H₂ is 2. Evaluate this hypothesis. 	[2]



Topic Chem 6 Q# 113/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q7/ www.SmashingScience.org :o)

(c) (i) Determine the orders of reaction of the reactants and the overall rate expression for the reaction between 2-bromobutane and aqueous sodium hydroxide using the data in the table.

Experiment	[NaOH] / mol dm ⁻³	[C ₄ H ₉ Br] / mol dm ⁻³	Rate / mol dm ⁻³ s ⁻¹
1	1.00	1.00	1.66 × 10 ⁻³
2	0.50	1.00	8.31 × 10 ⁻⁴
3	0.25	0.25	1.02 × 10 ⁻⁴
4	1.00	0.50	8.29 × 10 ⁻⁴

(ii) Determine the rate constant, *k*, with its units, using the data from experiment 3. [2]

(iii) Identify the molecularity of the rate-determining step in this reaction. [1]



[2]

Topic Chem 6 Q# 114/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/ The rate of reaction is an important factor in industrial processes such as the Contact process to make sulfur trioxide, SO₃(g). Define the term rate of reaction. (a) [1] Describe the collision theory. [3] (b) Topic Chem 7 Q# 115/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q5/ www.SmashingScience.org:o) This reaction is used in the manufacture of sulfuric acid. $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$ $K_c = 280 \text{ at } 1000 \text{ K}$ (e) 0.200 mol sulfur dioxide, 0.300 mol oxygen and 0.500 mol sulfur trioxide were mixed in a 1.00 dm3 flask at 1000 K. Predict the direction of the reaction showing your working. [3]

Topic Chem 7 Q# 116/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o)

Urea, (H₂N)₂CO, is excreted by mammals and can be used as a fertilizer.

(b) The structural formula of urea is shown.

(d) Urea can also be made by the direct combination of ammonia and carbon dioxide gases.

	$2NH_3(g) + CO_2(g) \rightleftharpoons (H_2N)_2CO(g) + H_2O(g) \qquad \Delta H < 0$	
(i)	State the equilibrium constant expression, K_c .	[1]
(ii)	Predict, with a reason, the effect on the equilibrium constant, K_c , when the temperature is increased.	[1]
(iii)	Determine an approximate order of magnitude for K_c , using sections 1 and 2 of the data booklet. Assume ΔG^e for the forward reaction is approximately +50 kJ at 298 K.	[2]
	OLIVER HILLERY	



Topic Chem 7 Q# 117/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q4/ www.SmashingScience.org:o)

(d) The equilibrium for a mixture of NO₂ and N₂O₄ gases is represented as:

$$2NO_2(g) \rightleftharpoons N_2O_4(g)$$

At 100°C, the equilibrium constant, K_c, is 0.21.

(i) At a given time, the concentration of NO₂(g) and N₂O₄(g) were 0.52 and 0.10 mol dm⁻³ respectively.
 Deduce, showing your reasoning, if the forward or the reverse reaction is favoured at this time.

[2]

٠.	-	•				:	•		-	 •	•	1		•	ं	•		8.	•	•	 •			•	•	•	-	•		•			Ť	•	ं							
		*	•	100			٠		*	 *	•					+0			•	•		*	•	*				*	 *		٠			• •	*	1 5		*	* :		•	
		-	-			* *	*	000	-	 **				***	-	***	•	 	100	*	 -	-	 -	W-010			-				***	 		1	-			*		4		
		*	 -	-	 -			 		 *			-	* *										*			-	- 1			**	 4	2		*	 	-		¥ 54			


(ii) Comment on the value of ΔG when the reaction quotient equals the equilibrium constant, Q = K.





Topic Chem 7 Q# 118/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q3/ www.SmashingScience.org:o)

PCl₅(g) and Cl₂(g) were placed in a sealed flask and allowed to reach equilibrium at 200°C.
 The enthalpy change, ΔH, for the decomposition of PCl₅(g) is positive.



[Source: Source: http://education.alberta.ca/media]

(a)	(i)	Deduce the equilibrium constant expression, K_c , for the decomposition of $\mathrm{PCl}_5(g)$.	[1]
5.55.			
	(ii)	Deduce, giving a reason, the factor responsible for establishing the new equilibrium after 14 minutes.	[2]
***		**************************************	



Topic Chem 7 Q# 119/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/

- Ethane-1,2-diol, HOCH₂CH₂OH, has a wide variety of uses including the removal of ice from aircraft and heat transfer in a solar cell.
 - (a) Ethane-1,2-diol can be formed according to the following reaction.

$$2CO(g) + 3H_2(g) \rightleftharpoons HOCH_2CH_2OH(g)$$

(i)	Deduce the equilibrium constant expression, \mathcal{K}_{c} , for this reaction.	[1]
		* */*

(ii) State how increasing the pressure of the reaction mixture at constant temperature will affect the position of equilibrium and the value of K_e.

[2]

Position of eq	uilibriu <mark>m</mark> :				
K _c :					
		****	 	 	*

Topic Chem 7 Q# 120/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/

2. One of the main constituents of acid deposition is sulfuric acid, H₂SO₄. This acid is formed from the sulfur dioxide pollutant, SO₂.

A mechanism proposed for its formation is:

$$\text{HO} \bullet (g) + \text{SO}_2(g) \rightarrow \text{HOSO}_2(g)$$

 $\text{HOSO}_2(g) + \text{O}_2(g) \rightarrow \text{HOO} \bullet (g) + \text{SO}_3(g)$
 $\text{SO}_3(g) + \text{H}_2\text{O}(1) \rightarrow \text{H}_2\text{SO}_4(aq)$



(b) Consider the following equilibrium between the two oxides of sulfur, sulfur dioxide and sulfur trioxide:

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$
 $\Delta H = -198 \text{ kJ}$

Predict, with a reason, in which direction the position of equilibrium will shift for each of the changes listed below.

[3]

Change	Shift	Reason
Increase in temperature		****************
Increase in pressure		
Addition of a catalyst to the mixture		

Topic Chem 7 Q# 121/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/

 Phosgene, COCl₂, is usually produced by the reaction between carbon monoxide and chlorine according to the equation:

$$CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$$

(a) (i) Deduce the equilibrium constant expression, K., for thi	reaction

[1]

	The second of th	

(ii) At exactly 600 °C the value of the equilibrium constant is 0.200. Calculate the standard Gibbs free energy change, ΔG^e, for the reaction, in kJ, using sections 1 and 2 of the data booklet. State your answer to three significant figures.

[3]





Topic Chem 7 Q# 122/ IB CHEM/2015/w/TZ0/Paper 2 Section B/Higher Level/

(a) The following reaction is used in industry to obtain hydrogen from natural gas by partial oxidation with steam.

$$CH_4(g) + H_2O(g) \rightleftharpoons 3H_2(g) + CO(g)$$
 $\Delta H^{\Theta} = +206 \text{ kJ}$

 Describe the effect, if any, of each of the following changes on the equilibrium amount of hydrogen, giving a reason in each case.

[4]

	Increasing the pressure, at constant temperature:	
	Increasing the temperature, at constant pressure:	

i)	Discuss the effects of adding a solid catalyst to the mixture of methane and steam, at constant pressure and temperature.	

(iii) Deduce the equilibrium constant expression, K_c , for the reaction.



(iv) Identify which of the changes in part (a) (i) will affect the value of K_c and whether the value will increase or decrease.

[1]

[1]

(b)	Th	e equilibrium constant, K_c , for the reaction	
		$CO(g) + H_2O(g) \rightleftharpoons H_2(g) + CO_2(g)$	
	wa	s found to be 10.0 at 420°C.	
	Ca	00 mol of CO(g) and 1.00 mol of H ₂ O(g) are mixed in a 1.00 dm ³ container at 420 °C. Iculate the equilibrium concentration of each component in the mixture, showing ur working.	[3]
	5.5		
	202		
	• • •	***************************************	
	2%		
	:		
	* *		
Topi	c Che r	n 7 Q# 123/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/	
7.		en nitrogen gas and hydrogen gas are allowed to react in a closed container the follow illibrium is established.	ing
		$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ $\Delta H = -92.6 \text{ kJ}$	
The a	above	equation is needed in question (d)(i)	
(d)	(i)	Deduce the equilibrium constant expression, K_c , for the reaction on page 10.	[1]
	<u> </u>		

		CMA CHIMA I I I	
	(ii)	When 1.00 mol of nitrogen and 3.00 mol of hydrogen were allowed to reach equilibrium in a 1.00 dm ³ container at a temperature of 500 °C and a pressure of 1000 atm, the equilibrium mixture contained 1.46 mol of ammonia.	
		Calculate the value of K _c at 500 °C.	[2]

Topic Chem 7 Q# 124/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/

 When nitrogen gas and hydrogen gas are allowed to react in a closed container the following equilibrium is established.

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$
 $\Delta H = -92.6 \text{ kJ}$

Outline two characteristics of a reversible reaction in a state of dynamic equilibrium.

Predict, with a reason, how each of the following changes affects the position of equilibrium.
The volume of the container is increased.

Ammonia is removed from the equilibrium mixture.

ii)) Define the term activation energy, E _a .	
		, //



		onia is manufactured by the Haber process in which iron is used as a catalyst. ain the effect of a catalyst on the rate of reaction.	[2]

(c)		eal conditions used in the Haber process are 500°C and 200 atm, resulting in eximately 15% yield of ammonia.	
	(i)	Explain why a temperature lower than 500 °C is not used.	[2]
	(ii)	Outline why a pressure higher than 200 atm is not often used.	[1]
	(ii)	Outline why a pressure higher than 200 atm is not often used.	[1]
pic ([1]
	Chem 7		[1]
	Chem 7 The rat	Q# 125/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/ te of reaction is an important factor in industrial processes such as the Contact	[1]
	Chem 7 The rat	Q# 125/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/ te of reaction is an important factor in industrial processes such as the Contact s to make sulfur trioxide, SO ₃ (g).	[1]
	Chem 7 The rat	Q# 125/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/ te of reaction is an important factor in industrial processes such as the Contact s to make sulfur trioxide, $SO_3(g)$. The Contact process involves this homogeneous equilibrium: $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g) \qquad \Delta H = -198 \text{kJ}$	
	The rat process (c) T	Q# 125/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/ te of reaction is an important factor in industrial processes such as the Contact s to make sulfur trioxide, $SO_3(g)$. The Contact process involves this homogeneous equilibrium: $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g) \qquad \Delta H = -198 \text{kJ}$ i) State and explain how increasing the pressure of the reaction mixture affects the	[1] e
	The rat process (c) T	Q# 125/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/ te of reaction is an important factor in industrial processes such as the Contact s to make sulfur trioxide, $SO_3(g)$. The Contact process involves this homogeneous equilibrium: $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g) \qquad \Delta H = -198 \text{kJ}$ i) State and explain how increasing the pressure of the reaction mixture affects the	
	The rat process (c) T	Q# 125/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/ te of reaction is an important factor in industrial processes such as the Contact s to make sulfur trioxide, $SO_3(g)$. The Contact process involves this homogeneous equilibrium: $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g) \qquad \Delta H = -198 \text{kJ}$ i) State and explain how increasing the pressure of the reaction mixture affects the	

	temperature and pressure.	[4]
(iii)	State the effect of increasing temperature on the value of $K_{\rm e}$ for this reaction.	[1]


Outl	ine the economic importance of using a catalyst in the Contact process.	[2]
		[2]
	ine the economic importance of using a catalyst in the Contact process.	[2]
		[2]
		[2]

Cher		
Cher	n 8 Q# 126/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org	:o)
Cher	m 8 Q# 126/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org ber forms two chlorides, copper(I) chloride and copper(II) chloride. (iv) Examine the relationship between the Brønsted–Lowry and Lewis definit	:o)
Cher	m 8 Q# 126/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org ber forms two chlorides, copper(I) chloride and copper(II) chloride. (iv) Examine the relationship between the Brønsted–Lowry and Lewis definit	:o)
Cher	m 8 Q# 126/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org ber forms two chlorides, copper(I) chloride and copper(II) chloride. (iv) Examine the relationship between the Brønsted–Lowry and Lewis definit	:o)
Cher	m 8 Q# 126/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org ber forms two chlorides, copper(I) chloride and copper(II) chloride. (iv) Examine the relationship between the Brønsted–Lowry and Lewis definit	:o)
Cher	m 8 Q# 126/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org ber forms two chlorides, copper(I) chloride and copper(II) chloride. (iv) Examine the relationship between the Brønsted–Lowry and Lewis definit	:o)



Topic Chem 8 Q# 127/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q6./ www.SmashingScience.org:o)

- Another common acid found in food is ethanoic acid.
 - (a) A sample of ethanoic acid was titrated with sodium hydroxide solution, and the following pH curve obtained.



Annotate the graph to show the buffer region and the volume of sodium hydroxide at the equivalence point.

[2]

 (b) (i) Identify the most suitable indicator for the titration using section 22 of the data booklet.

[1]

Trica.	TOUR / /	1 7	
	the second second second second		

(ii) Describe, using a suitable equation, how the buffer solution formed during the titration resists pH changes when a small amount of acid is added. [2]



Topic Chem 8 Q# 128/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q5./ www.SmashingScience.org:o)

A molecule of citric acid, C₈H₈O₇, is shown.

The equation for the first dissociation of citric acid in water is

$$C_6H_8O_7(aq) + H_2O(l) \rightleftharpoons C_6H_7O_7^-(aq) + H_3O^+(aq)$$

(a) (i)	Identify a conjugate acid-base pair in the equation.		[1]
(ii)	The value of K_a at 298 K for the first dissociation is 5.0	1 × 10 ⁻⁴ .	<u> </u>
	State, giving a reason, the strength of citric acid.		[1]
(iii)	The dissociation of citric acid is an endothermic proce hydrogen ion concentration, $[H^{+}]$, and on K_{a} , of increase		[2]
(iii)			[2]
(iii)	hydrogen ion concentration, [H ⁺], and on K _a , of increase	sing the temperature.	[2]



	The state of the s		The state of the s		
		names neares na <mark>s</mark> an			
	/ IB Chem/2018/w/Tz id, CH ₃ CH ₂ CH ₂ CO0 State the equation	OH, is a weak acid	d and ethylamin	e, CH ₃ CH ₂ NH ₂	
oic a	id, CH ₃ CH ₂ CH ₂ CO	OH, is a weak acid	d and ethylamin	e, CH ₃ CH ₂ NH ₂	
(i) noic a	id, CH ₃ CH ₂ CH ₂ CO	OH, is a weak acid	d and ethylamin	e, CH ₃ CH ₂ NH ₂	
(i) noic a	id, CH ₃ CH ₂ CH ₂ CO0 State the equation id:	OH, is a weak acid	d and ethylamin	e, CH ₃ CH ₂ NH ₂	
(i) noic a	id, CH ₃ CH ₂ CH ₂ CO0 State the equation id:	OH, is a weak acid	d and ethylamin	e, CH ₃ CH ₂ NH ₂	
(i) noic a	id, CH ₃ CH ₂ CH ₂ CO0 State the equation id:	OH, is a weak acid	d and ethylamin	e, CH ₃ CH ₂ NH ₂	
(i) noic a	id, CH ₃ CH ₂ CH ₂ CO0 State the equation id:	OH, is a weak acid	d and ethylamin	e, CH ₃ CH ₂ NH ₂	
			ydrochloric acid of equal concentration, sta	ydrochloric acid of equal concentration, stating the expecte	Outline two laboratory methods of distinguishing between solutions of citrydrochloric acid of equal concentration, stating the expected observation




(b)	(i)	A 0.250 moldm ⁻³ aqueous solution of butanoic acid has a concentration of hydrogen ions, [H ⁺], of 0.00192 moldm ⁻³ . Calculate the concentration of hydroxide ions, [OH ⁻], in the solution at 298 K.	[1]
	(ii)	Determine the pH of a 0.250 mol dm ⁻³ aqueous solution of ethylamine at 298 K, using section 21 of the data booklet.	[3]
23 1 24 2 24 2 24 2 24 2			



(c) Sketch the pH curve for the titration of 25.0 cm³ of ethylamine aqueous solution with 50.0 cm³ of butanoic acid aqueous solution of equal concentration. No calculations are required.

[3]



Topic Chem 8 Q# 130/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q5/ www.SmashingScience.org:o)

 Limescale, CaCO₃ (s), can be removed from water kettles by using vinegar, a dilute solution of ethanoic acid, CH₃COOH (aq).

(a)	Predict, giving a reason, a difference between the reactions of the same concentrations
	of hydrochloric acid and ethanoic acid with samples of calcium carbonate.

	 	 	*										 						 •	 								
		 	•) . ()	 -				*>	*		 (oe)(e			 	× (• •					60)e				*		
 	 	 	2			•						 2								 				 2.0		4		
	 	 	-	 •	•		*	i.					7:1	•	 			or o	 	•		•			•			

[2]

	[1]

(c) Write an equation to show ammonia, NH ₃ , acting as a Brønsted–Lowry base and a different equation to show it acting as a Lewis base.	
Brønsted-Lowry base:	
Lewis base:	
K_a (2,2-dimethylpropanoic acid) = 9.333×10^{-6}	
(e) Explain, using appropriate equations, how a suitably concentrated solution formed by the partial neutralization of 2,2-dimethylpropanoic acid with sodium hydroxide acts as buffer solution.	
the partial neutralization of 2,2-dimethylpropanoic acid with sodium hydroxide acts as	
the partial neutralization of 2,2-dimethylpropanoic acid with sodium hydroxide acts as	
the partial neutralization of 2,2-dimethylpropanoic acid with sodium hydroxide acts as	
the partial neutralization of 2,2-dimethylpropanoic acid with sodium hydroxide acts as	

Topic Chem 8 Q# 131/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q2/ www.SmashingScience.org :o)

(f) Calcium carbide reacts with water to form ethyne and calcium hydroxide.

$$CaC_{2}(s) + H_{2}O(l) \rightarrow C_{2}H_{2}(g) + Ca(OH)_{2}(aq)$$

			$CaC_2(s) + H_2C(t) \rightarrow C$	$G_2 \cap G_2(g) + Ga(O \cap G_2(aq))$	
		Esti	mate the pH of the resultant solution.		[1]
opic 8.			32/ IB Chem/2017/s/TZ 2/Paper 2/Higher ids and bases ionize in water.	Level/Q8/ www.SmashingScience.org :o)	
	(a)	Sodi	um hypochlorite ionizes in water.		
			$OCl^-(aq) + H_2O(l) \rightleftharpoons O$	H ⁻ (aq) + HOCl(aq)	
		(i)	Identify the amphiprotic species.		[1]
					CF7 404 YF9
		(ii)	Identify one conjugate acid-base p	air in the reaction.	[1]
			Acid	Base	

	(b)			own monoprotic acid, HA, was titrated required to reach the equivalence po	
		(i)	Calculate the amount, in mol, of Na	aOH (aq) used.	[1]
		(ii)	Calculate the molar mass of the ac	id.	[1]
	0.000.00				
	1212/4/				

(iii) Calculate [H ⁺] in the NaOH solution. [1]	
(iv) The following curve was obtained using a pH probe.	
14	
12	
10	
6 -	
2	
0 5 10 15 20 25 30 35 Volume of NaOH added / cm ³	
State, giving a reason, the strength of the acid.	[1]
(v) State a technique other than a pH titration that can be used to detect the equivalence point.	[1]
(vi) Deduce the pK_a for this acid.	[1]

	to two decimal places.	
w.	Two hydrides of nitrogen are ammonia and hydrazine, N ₂ H ₄ . One derivative of ammonia is methanamine whose molecular structure is shown below. H H H H Ammonia socoto reversible with treater	
	(c) Ammonia reacts reversibly with water.	
	$NH_3(g) + H_2O(l) \rightleftharpoons NH_4^+(aq) + OH^-(aq)$	
	Explain the effect of adding H ⁺ (aq) ions on the position of the equilibrium.	2
		_

 (d) (i) Hydrazine reacts with water in a similar way to ammonia. (The association of a molecule of hydrazine with a second H⁺ is so small it can be neglected.)

$$N_2H_4(aq)+H_2O(l) \rightleftharpoons N_2H_5^+(aq)+OH^-(aq)$$

$$pK_b$$
 (hydrazine) = 5.77

Calculate the pH of a 0.0100 mol dm⁻³ solution of hydrazine.

are ere ere er		
ಾರ್ಯಗಳಿಂದ ಮುನುವರು		
(ii)	Suggest a suitable indicator for the titration of hydrazine solut sulfuric acid using section 22 of the data booklet.	tion with dilute

(e) Outline, using an ionic equation, what is observed when magnesium powder is added to a solution of ammonium chloride. [2]



[3]

Topic Chem 8 Q# 134/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/

by titration with a standard solution of sodium hydroxide, NaOH(aq). Distinguish between a weak acid and a strong acid. [1] Weak acid: Strong acid: (b) Suggest why it is more convenient to express acidity using the pH scale instead of using the concentration of hydrogen ions. [1] 5.00 g of an impure sample of hydrated ethanedioic acid, (COOH)2.2H2O, was dissolved in water to make 1.00 dm³ of solution. 25.0 cm³ samples of this solution were titrated against a 0.100 mol dm⁻³ solution of sodium hydroxide using a suitable indicator. $(COOH)_2(aq) + 2NaOH(aq) \rightarrow (COONa)_2(aq) + 2H_2O(l)$ The mean value of the titre was 14.0 cm³. Suggest a suitable indicator for this titration. Use section 22 of the data booklet. (i) [1]

The concentration of a solution of a weak acid, such as ethanedioic acid, can be determined



Topic Chem 8 Q# 135/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/


- 7. This question is about the weak acid methanoic acid, HCOOH.
 - (a) Calculate the pH of $0.0100\,\mathrm{mol\,dm^{-3}}$ methanoic acid stating any assumption you make. $K_a = 1.6 \times 10^{-4}$.

[3]

Assumption:		

(b) (i) Sketch a graph of pH against volume of a strong base added to a weak acid showing how you would determine pK_a for the weak acid.

[2]





(ii)	Explain, using an equation, why the pH increases very little in the buffer region when a small amount of alkali is added.	[2

A mechan	ism proposed for its formation is:	
	$HO_{\bullet}(g) + SO_{2}(g) \rightarrow HOSO_{2}(g)$	
	$HOSO_2(g) + O_2(g) \rightarrow HOO \cdot (g) + SO_3(g)$	
	$SO_3(g) + H_2O(1) \rightarrow H_2SO_4(aq)$	
(a) State	e what the symbol (•) represents in the species shown in this mechanism.	[1]

		, 7
rormul	the an equation for the reaction of introgen dioxide with water.	
	One of the from the s A mechan (a) State	when a small amount of alkali is added. c Chem 8 Q# 136/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/ One of the main constituents of acid deposition is sulfuric acid, H ₂ SO ₄ . This acid is formed from the sulfur dioxide pollutant, SO ₂ . A mechanism proposed for its formation is: HO•(g)+SO ₂ (g) → HOSO ₂ (g) HOSO ₂ (g) → HOO•(g)+SO ₃ (g) SO ₃ (g)+H ₂ O(l) → H ₂ SO ₄ (aq) (a) State what the symbol (•) represents in the species shown in this mechanism.

Topic Chem 8 Q# 137/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/

A 0.12 mol dm⁻³ sodium methanoate solution dissociates completely into its ions.

(a)	Formulate the equation, including state symbols, for the equilibrium reaction of the hydrolysis of the methanoate anion.	[.

ons 1	and 21 of the data booklet may be used for parts (b) to (e).	
(b)	Calculate the value of K_a , the acid dissociation constant at 298K, for an aqueous solution of methanoic acid.	I
(c)	Calculate the value of K_b , the base dissociation constant, for the conjugate base.	1
<u> </u>	THE ATO	



	Determine the concentration, in mol dm ⁻³ , of hydroxide ion, [OH ⁻ (aq)], in the original 0.12 mol dm ⁻³ sodium methanoate solution, mentioning one assumption made.	[3]
	Calculate the pH of the 0.12 mol dm ⁻³ sodium methanoate solution.	[2]
_		


C		
	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution	
C	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution	
C	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	
C	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	
	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	
	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	
	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	
	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	
	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	
	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	
	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	
	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	
	them 8 Q# 138/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience (d) Phenylamine can act as a weak base. Calculate the pH of a 0.0100 mol dm ⁻³ solution of phenylamine at 298 K using section 21 of the data booklet.	

Topic Chem 8 Q# 139/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q1(a) Not with 1(a)(

Phosphine (IUPAC name phosphane) is a hydride of phosphorus, with the formula PH₃.

It's Lewis structure is this:





1		
(ii	 Deduce, giving your reason, whether phosphine would act as a Lewis acid, a Lewis base, or neither. 	
(V) Ammonia acts as a weak Brønsted–Lowry base when dissolved in water.	
	$NH_3(aq) + H_2O(l) \rightleftharpoons NH_4^+(aq) + OH^-(aq)$	
	Outline what is meant by the terms "weak" and "Brønsted-Lowry base".	[
Weak:		
Brønste	d–Lowry base:	
Brønste	I–Lowry base:	
(b) Pr	osphine is usually prepared by heating white phosphorus, one of the allotropes phosphorus, with concentrated aqueous sodium hydroxide. The equation for the action is:	
(b) Pr	osphine is usually prepared by heating white phosphorus, one of the allotropes phosphorus, with concentrated aqueous sodium hydroxide. The equation for the	
(b) Pr	osphine is usually prepared by heating white phosphorus, one of the allotropes phosphorus, with concentrated aqueous sodium hydroxide. The equation for the action is: $P_4(s) + 3OH^-(aq) + 3H_2O(l) \rightarrow PH_3(g) + 3H_2PO_2^-(aq)$	
(b) Proof of re	osphine is usually prepared by heating white phosphorus, one of the allotropes phosphorus, with concentrated aqueous sodium hydroxide. The equation for the action is: $P_4(s) + 3OH^-(aq) + 3H_2O(l) \rightarrow PH_3(g) + 3H_2PO_2^-(aq)$ The ion $H_2PO_2^-$ is amphiprotic. Outline what is meant by amphiprotic, giving the	
(b) Proof of re	osphine is usually prepared by heating white phosphorus, one of the allotropes phosphorus, with concentrated aqueous sodium hydroxide. The equation for the action is: $P_4(s) + 3OH^-(aq) + 3H_2O(l) \rightarrow PH_3(g) + 3H_2PO_2^-(aq)$ The ion $H_2PO_2^-$ is amphiprotic. Outline what is meant by amphiprotic, giving the	
(b) Proof of re	osphine is usually prepared by heating white phosphorus, one of the allotropes phosphorus, with concentrated aqueous sodium hydroxide. The equation for the action is: $P_4(s) + 3OH^-(aq) + 3H_2O(l) \rightarrow PH_3(g) + 3H_2PO_2^-(aq)$ The ion $H_2PO_2^-$ is amphiprotic. Outline what is meant by amphiprotic, giving the	



(iv)	State the equation for the reaction of this oxide of phosphorus with water.	[1
(v)	Suggest why oxides of phosphorus are not major contributors to acid deposition.	[1
(vi)	The levels of sulfur dioxide, a major contributor to acid deposition, can be minimized by either pre-combustion and post-combustion methods. Outline one technique of each method.	[2
Pre-combu	ustion:	
Post-comb	oustion:	
Chem 8 Q#	140/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/	
Tin(II) chlo	oride is a white solid that is commonly used as a reducing agent.	
(iii)	Tin(II) chloride reacts with water to precipitate the insoluble basic chloride, Sn(OH)Cl.	
	$SnCl_2(aq) + H_2O(l) \rightleftharpoons Sn(OH)Cl(s) + H^+(aq) + Cl^-(aq)$	
	Suggest why tin(II) chloride is usually dissolved in dilute hydrochloric acid.	[1



Topic Chem 8 Q# 141/ IB CHEM/2015/w/TZ0/Paper 2 Section B/Higher Level/Q7/ www.SmashingScience.org :o)

(d) Consider the following spontaneous reactions.

Fe(s) + Ni²⁺(aq)
$$\rightarrow$$
 Fe²⁺(aq) + Ni(s)
Zn(s) + Fe²⁺(aq) \rightarrow Zn²⁺(aq) + Fe(s)

$$Ni(s) + Pb^{2+}(aq) \rightarrow Ni^{2+}(aq) + Pb(s)$$


(i)	Deduce the order of increasing reactivity of the metals based on the reactions above.	[2]
(ii)	Identify the strongest oxidizing agent in the reactions above.	[1]

De	educe the half-equations for the formation of the major product at the positive ectrode (anode) when the following aqueous solutions are electrolysed.	[2]
ele	educe the half-equations for the formation of the major product at the positive ectrode (anode) when the following aqueous solutions are electrolysed.	[2]
ele	ectrode (anode) when the following aqueous solutions are electrolysed.	[2]
ele	ectrode (anode) when the following aqueous solutions are electrolysed.	[2]
Dil	ute sodium chloride:	[2]
Dil	ute sodium chloride:	[2]



Topic Chem 8 Q# 142/ IB CHEM/2015/w/TZ0/Paper 2 Section B/Higher Level/

(a) 20.0 cm³ aqueous solutions of two bases, each with a concentration of 0.100 mol dm⁻³ were separately titrated with 0.100 mol dm⁻³ hydrochloric acid, HCl (aq), and the following graph was obtained.





(i)	Deduce the pH at the equivalence points for base I and base II.	[2]
(ii)	Suggest why the titration curve for base I is different from base II.	[1]
(iii)	State the formulas of two possible bases which could be used as base I.	[1]
(iv)	Calculate, using data from the graph, the dissociation constant, $K_{\rm b}$, of base II, showing your working.	[3]
(v)	Suggest an indicator that can be used for both titrations.	[1]



1		

(ii)	Calculate the pH of a solution prepared by mixing $40.0\mathrm{cm^3}$ of $0.200\mathrm{moldm^{-3}}$ NH ₃ (aq) and $40.0\mathrm{cm^3}$ of $0.100\mathrm{moldm^{-3}}$ HCl (aq), showing your working. (pK _b NH ₃ = 4.75 at $298\mathrm{K}$)	

	Reaction B $HCO_3^-(aq) + H_2O(l) \rightleftharpoons CO_3^{2-}(aq) + H_3O^+(aq)$	
(i)	Reaction B $HCO_3^-(aq) + H_2O(l) \rightleftharpoons CO_3^{2^-}(aq) + H_3O^+(aq)$ Explain whether $HCO_3^-(aq)$ behaves as an acid or a base in each of the reactions A and B.	
(i)	Explain whether HCO ₃ ⁻ (aq) behaves as an acid or a base in each of the	
(i)	Explain whether $HCO_3^-(aq)$ behaves as an acid or a base in each of the reactions A and B .	
(i)	Explain whether $HCO_3^-(aq)$ behaves as an acid or a base in each of the reactions A and B .	
(i)	Explain whether $HCO_3^-(aq)$ behaves as an acid or a base in each of the reactions A and B .	
(i)	Explain whether HCO ₃ ⁻ (aq) behaves as an acid or a base in each of the reactions A and B. Reaction A:	
(i)	Explain whether HCO ₃ ⁻ (aq) behaves as an acid or a base in each of the reactions A and B. Reaction A:	
	Explain whether HCO ₃ ⁻ (aq) behaves as an acid or a base in each of the reactions A and B. Reaction A: Reaction B:	
(i) (ii)	Explain whether HCO ₃ ⁻ (aq) behaves as an acid or a base in each of the reactions A and B. Reaction A:	
	Explain whether HCO ₃ ⁻ (aq) behaves as an acid or a base in each of the reactions A and B. Reaction A: Reaction B:	
	Explain whether HCO ₃ ⁻ (aq) behaves as an acid or a base in each of the reactions A and B. Reaction A: Reaction B: Deduce two conjugate acid-base pairs from reactions A and B.	

(i)	Distinguish between strong and weak acids.]
(ii)	A 1.00 g sample of solid magnesium carbonate, MgCO ₃ , is added to separate solutions of HNO ₃ and HNO ₂ of the same concentration and temperature. State one similarity and one difference in the observations made in these reactions.	[2
	Similarity:	
	Difference:	
(iii)	A solution of HNO ₃ has a pH of 1, while a solution of HNO ₂ has a pH of 5. Determine the ratio of the hydrogen ion concentration in HNO ₃ :HNO ₂ .	[1
Chem	8 Q# 143/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience	ce.
(b)	Cu ²⁺ (aq) reacts with ammonia to form the complex ion [Cu(NH ₃) ₄] ²⁺ . Explain this reaction in terms of acid-base theory, and outline the bonding in the complex formed between Cu ²⁺ and NH ₃ .	
		5 5
		117.00


(d) Nitric acid, HNO₃, and nitrous acid, HNO₂, are described as strong and weak acids respectively.



(i)	Define the term base according to the Le	wis theory.	[
(ii)	Define the term weak base according to t	he Brønsted–Lowry theory.	[*
	********************		* * *
(iii)	Deduce the formulas of conjugate acid-b	ase pairs in the reaction below.	
	$CH_3NH_2(aq) + H_2O(l) \rightleftharpoons$	$^{+}$ CH ₃ NH ₃ ⁺ (aq) + OH ⁻ (aq)	[
	Acid	ā	
-	ACId	Conjugate base	
Det	termine the pH of a 0.100 mol dm ⁻³ solution		
Det			
Det	termine the pH of a 0.100 mol dm ⁻³ solution les 2 and 15 of the data booklet.	of ammonia, NH ₃ (aq), using	
Det	termine the pH of a 0.100 mol dm ⁻³ solution	of ammonia, NH ₃ (aq), using	
Det	termine the pH of a 0.100 mol dm ⁻³ solution les 2 and 15 of the data booklet.	of ammonia, NH ₃ (aq), using	
Det	termine the pH of a 0.100 mol dm ⁻³ solution les 2 and 15 of the data booklet.	of ammonia, NH ₃ (aq), using	
Det tabl	termine the pH of a 0.100 mol dm ⁻³ solution les 2 and 15 of the data booklet.	of ammonia, NH ₃ (aq), using	
Det tabl	termine the pH of a 0.100 mol dm ⁻³ solution les 2 and 15 of the data booklet.	of ammonia, NH ₃ (aq), using	
Dettabl	termine the pH of a 0.100 mol dm ⁻³ solution les 2 and 15 of the data booklet.	of ammonia, NH ₃ (aq), using	
Det tabl	termine the pH of a 0.100 mol dm ⁻³ solution les 2 and 15 of the data booklet.	of ammonia, NH ₃ (aq), using	
Det tabl	termine the pH of a 0.100 mol dm ⁻³ solution les 2 and 15 of the data booklet.	of ammonia, NH ₃ (aq), using	
Dettabl	termine the pH of a 0.100 mol dm ⁻³ solution les 2 and 15 of the data booklet.	of ammonia, NH ₃ (aq), using	
Det tabl	termine the pH of a 0.100 mol dm ⁻³ solution les 2 and 15 of the data booklet.	of ammonia, NH ₃ (aq), using	

(g) (i) Sketch the pH titration curve obtained when 50.0 cm³ of 0.100 mol dm⁻³ NH₃ (aq) is added to 25.0 cm³ of 0.100 mol dm⁻³ HCl (aq).

[3]



(ii) Identify an indicator from table 16 of the data booklet that could be used for this titration.

[1]

Topic Chem 8 Q# 145/ IB Chem/2015/s/TZ2/Paper 2 Section A/Higher Level/

A buffer solution with a pH of 3.87 contains 7.41g dm⁻³ of propanoic acid, CH₃CH₂COOH, together with an unknown quantity of sodium propanoate, CH₃CH₂COONa.

(a) Define the term *buffer solution*.

[2]

(b) Explain, using appropriate equations, how this solution acts as a buffer solution.

[2]

(c) Calculate the concentration, in moldm⁻³, of sodium propanoate in this buffer solution. The pK₃ of propanoic acid is 4.87 at 298 K.

[4]




Topic Chem 8 Q# 146/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/ Need table 21 and 22/ www.SmashingScience.org:o)

8.	Acide can	he described	as strong or weak.
ο.	Acius can	ne described	as subliq of wear.

(i)	Outline the difference in dissociation between strong and weak acids of the same concentration.	
(ii)	Describe three tests that can be carried out in the laboratory, and the expected results, to distinguish between 0.10 mol dm ⁻³ HCl (aq) and 0.10 mol dm ⁻³ CH ₃ COOH (aq).	
	1 1600	_
Cal	culate the pH, using table 15 of the data booklet, of a solution of ethanoic acid de by dissolving 1.40 g of the acid in distilled water to make a 500 cm ³ solution.	
	1965 * 327 A 1	_

(c) (i) Determine the pH at the equivalence point of the titration and the pK_a of an unknown acid using the acid-base titration curve below.



** ***	***	 		

(ii) Identify, using table 16 of the data booklet, a suitable indicator to show the end-point of this titration. [1]



	Describe how an indicator, that is a weak acid, works. Use Le Chatelier's principle in your answer.	[2
(i)	State the formula of the conjugate base of chloroethanoic acid, CH ₂ ClCOOH.	[1
(ii)	Identify, with a reason, whether chloroethanoic acid is weaker or stronger than ethanoic acid using table 15 of the data booklet.	[1
(iii)	Determine the pH of the solution resulting when 100 cm ³ of 0.50 mol dm ⁻³ CH ₂ ClCOOH is mixed with 200 cm ³ of 0.10 mol dm ⁻³ NaOH.	[4]
(iii)		[4
(iii)		[4
(iii)		[4
(iii)	CH ₂ ClCOOH is mixed with 200 cm ³ of 0.10 mol dm ⁻³ NaOH.	[4
(iii)	CH ₂ ClCOOH is mixed with 200 cm ³ of 0.10 mol dm ⁻³ NaOH.	[4
(III)	CH ₂ ClCOOH is mixed with 200 cm ³ of 0.10 mol dm ⁻³ NaOH.	[4
(iii)	CH ₂ ClCOOH is mixed with 200 cm ³ of 0.10 mol dm ⁻³ NaOH.	[4
(iii)	CH ₂ ClCOOH is mixed with 200 cm ³ of 0.10 mol dm ⁻³ NaOH.	[4



Topic Chem 8 Q# 147/ IB Chem/2015/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.SmashingScience.org:o)

(c) Predict and explain the pH of the following aqueous solutions, using equations to support your answer. [4]


	nium ch		40-1	4/-					
					 	 	 	 ****	* 5
Sodiun	n metha	noate,	HCOOI	Na (aq)					
				* * * * *	 	 	 	 * * *	

Topic Chem 9 Q# 148/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q7./ www.SmashingScience.org:0)

- Copper forms two chlorides, copper(I) chloride and copper(II) chloride.
 - (ii) Copper(II) chloride is used as a catalyst in the hydrogen chloride.

$$4HCl(g) + O_2(g) \rightarrow 2Cl_2(g) + 2H_2O(g)$$

(c) Two electrolysis cells were assembled using graphite electrodes and connected in series as shown.



Patrick Brannac

 State how current is conducted through the wires and through the electrolyte. 	[2]
Wires:	
Electrolyte:	
(ii) Write the half-equation for the formation of gas bubbles at electrode 1.	[1]
(iii) Bubbles of gas were also observed at another electrode. Identify the electrode and the gas.	[1
Electrode number (on diagram):	
Name of gas:	
(iv) Deduce the half-equation for the formation of the gas identified in (c)(iii).	[1
	2.2.2
	*25.5
SMASHINGILL	



chlo	ride and copper.	
	$2Cu^+(aq) \rightarrow Cu(s) + Cu^{2+}(aq)$	
(i)	Calculate the cell potential at 298 K for the disproportionation reaction, in V, using section 24 of the data booklet.	[1
(ii)	Comment on the spontaneity of the disproportionation reaction at 298 K.	[1
(iii)	Calculate the standard Gibbs free energy change, ΔG° , to two significant figures, for the disproportionation at 298 K. Use your answer from (e)(i) and sections 1 and 2 of the data booklet.	[1

(iv)	Suggest, giving a reason, whether the entropy of the system increases or decreases during the disproportionation.	[1

Copper(I) chloride undergoes a disproportionation reaction, producing copper(II)

(v)				a reason, on reaction			tandar	d e	nth	alpy	/ ch	ange	e, ΔF	f ^e , for	the	[
	****					*****	*** *** ***		4 40			× 600		********		
								272 2				0.270				
								1-7 T	51 50	ten t		ক হচত	ten t	ritiron ti		
(vi)				reason, le solutio		t of incre	easing (ten	pe	ratı	ıre o	on th	ne sta	ability o	of	[
									5 5							
				*****				e//e =				* * *				
hem 9 O#	149/ IB	Chemi	/2019/v	/TZ0/Papi	er 2/Highe	er Level/C	13 / www	w Sr	nas	hins	 Scie	nce	org 'o	1		
The biocl series of Na ₂ S ₂ O ₃ (nemica reactio	l oxyg	gen de	mand of a	water s	ample o	an be	dete	ern	nine	d by	y the	follo	wing	ın,	
The bioch series of	nemica reactio (aq).	al oxyg ons. Ti	gen der he fina	mand of a	water s tration o	ample of the sar	an be o	dete	ern	nine diun	d by	y the	follo	wing	n,	
The bioch series of	nemica reactio (aq).	nl oxyg ons. Ti 2Mn ²	gen der he fina (aq) +	mand of a	water s tration o	ample of the same $(q) \rightarrow 21$	an be omple with	dete ith : s) +	ern soc	nine diun	ed by n thi (l)	y the	follo	wing	n,	
The bioch series of	nemica reactio (aq).	al oxygons. The 2Mn ²	gen der he fina (aq) +) + 2I	mand of a step is to $O_2(aq) +$	a water s tration o 4OH⁻(a †(aq) →	ample of the said q $\rightarrow 21$ Mn^{2+} (a	an be on the mapping of the mapping	dete ith: s)+ (aq)	2H	nine diun	ed by n thi (l)	y the	follo	wing	n,	
The bioch series of	nemica reactio (aq). M t analy ely (da y 5). Ti	2Mn ² 2Mn ² sed tv	gen der he fina (aq) +) + 2Γ 2S ₂ O ₃ vo 300 and the owing	O ₂ (aq) + (aq) + 4H (aq) + I 0 cm ³ sale other afresults we	a water stration of 40H ⁺ (aq) → mples of ter leaving	ample of f the san aq) → 2l Mn ²⁺ (a 2I ⁻ (aq) water tang it sea	an be on the matter $MnO_2(s)$ $q) + I_2(s)$ $+ S_4O_6(s)$ aken from the matter s	determination (aq) + (aq) companda da	2F + eq)	nine diun 1 ₂ O 2H ₂	(l) O(l) the osul	follo	owing solution ne	n,	
The biocl series of Na ₂ S ₂ O ₃ (A student immediat days (day	nemica reactio (aq). M t analy ely (da y 5). Ti	2Mn ² 2Mn ² sed tv	yo 300 and the owing of the second se	O ₂ (aq) + (aq) + 4H (aq) + 1 O cm ³ sale other afresults well)	a water stration of 40H ⁺ (aq) → anples of ter leaving tere obtains	ample of the san and the san	an be on the matter $MnO_2(s)$	determination (aq) (aq) a dation	2F + eq)	nine diun 1 ₂ O 2H ₂	(l) O(l) the osul	follo	owing solution ne	ın,	
The biocl series of Na ₂ S ₂ O ₃ (A student immediat days (day	nemica reactio (aq). M t analy ely (da y 5). Ti	2Mn ² 2Mn ² sed tv	yen der he fina (aq) +) + 2I 2S ₂ O ₃ yo 300 and the owing of	onand of a step is to the country of a step is to the country of a step is to the country of a step in the country of a s	a water stration of 40H ⁺ (aq) → anples of ter leaving tere obtains	ample of the san $q) \rightarrow 2l$ $Mn^{2+}(aq)$ $Vater tang it seaned for q Var d d d d d$	an be on the matter $MnO_2(s)$	determination (aq) (aq) a dation	2F + eq)	nine diun 1 ₂ O 2H ₂	(l) O(l) the osul	follo	owing solution ne	n,	
The biocl series of Na ₂ S ₂ O ₃ (A student immediat days (day	nemica reactio (aq). M t analy ely (da y 5). Ti	2Mn ² 2Mn ² sed tv	gen der he fina (aq) +) + 2Γ 2S ₂ O ₃ vo 300 and the owing (ac	O ₂ (aq) + (aq) + 4H (aq) + 1 O cm ³ sale other afresults well)	a water stration of 40H ⁺ (aq) → anples of ter leaving tere obtains	ample of the san and the san	an be on the matter $MnO_2(s)$	determination (aq) (aq) a dation	2F + eq)	nine diun 1 ₂ O 2H ₂	(l) O(l) the osul	follo	owing solution ne	n,	



[1]

(ii)	Calculate the number of moles of oxygen in the day 0 sample.	[2]
	ga en swa ni nga isinga ising basi na nakini gab na nga na nga naka indika inaka isinga naanka	

(iii)	The day 5 sample contained 5.03×10^{-5} moles of oxygen.	
	Determine the 5-day biochemical oxygen demand of the pond, in mg dm ⁻³ ("parts per million", ppm).	[2]

	50/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q6/ www.SmashingScience.org:o)	-
	acid, CH ₃ CH ₂ CH ₂ COOH, is a weak acid and ethylamine, CH ₃ CH ₂ NH ₂ , is a weak base	
(iii)	Deduce the average oxidation state of carbon in butanoic acid.	[1]

	51/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q3/ www.SmashingScience.org :o)	
(d) (i)	an form the bromate(V) ion, BrO ₃ ⁻ . Bromate(V) ions act as oxidizing agents in acidic conditions to form bromide ions.	
(4)	med in demandable dead for 1	101
	Deduce the half-equation for this reduction reaction.	[2]




		(ii)	Bromate(V) ions oxidize iron(II) ions, Fe ²⁺ , to iron(III) ions, Fe ³⁺ .	
			Deduce the equation for this redox reaction.	ľ
		(iii)	Calculate the standard Gibbs free energy change, ΔG^{Θ} , in J, of the redox reaction in (ii), using sections 1 and 24 of the data booklet.	
			E^{\oplus} (BrO ₃ ⁻ /Br ⁻) = +1.44 V	[2
	* * * *		*****	

opic 1.	3.26	g of i	52/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o) ron powder are added to 80.0 cm ³ of 0.200 mol dm ⁻³ copper(II) sulfate solution. The reaction occurs:	
			$Fe(s) + CuSO_4(aq) \rightarrow FeSO_4(aq) + Cu(s)$	
		State	ident electrolyzed aqueous iron(II) sulfate, FeSO ₄ (aq), using platinum electrodes. e half-equations for the reactions at the electrodes, using section 24 of the booklet.	[2
	Anod	e (po	sitive electrode):	
	Catho	ode (ı	negative electrode):	
			*** *** *** *** *** *** *** *** *** **	



Topic Chem 9 Q# 153/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q6/ www.SmashingScience.org:o)

The diagram shows an incomplete voltaic cell with a light bulb in the circuit.





	(e)	Determine the loss in mass of one electrode if the mass of the other electrode increases by 0.10 g.	[2]
	* * * /		
	#09E0#20		
Topic 4.		Q# 154/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q4/ www.SmashingScience.org :o) Ids can be formed in many ways.	
	(a)	The landing module for the Apollo mission used rocket fuel made from a mixture of hydrazine, N_2H_4 , and dinitrogen tetraoxide, N_2O_4 .	
		$2N_2H_4(l) + N_2O_4(l) \rightarrow 3N_2(g) + 4H_2O(g)$	
		(iii) Determine the oxidation state of nitrogen in the two reactants.	[1
	N ₂ H	l ₄ :	
	N ₂ C	Date of the state	
750	7.15.7		
		(iv) Deduce, giving a reason, which species is the reducing agent.	[1
L Topic	Chem 9	Q# 155/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q2/ www.SmashingScience.org :o)	
2.	(a) K ₂ (An acidic sample of a waste solution containing $\mathrm{Sn^{2+}}(\mathrm{aq})$ reacted completely with $\mathrm{Cr_2O_7}$ solution to form $\mathrm{Sn^{4+}}(\mathrm{aq})$.	
		(i) State the oxidation half-equation.	[1



 (b) (i) Corrosion of iron is similar to the processes that occur in a voltaic cell. The initial steps involve the following half-equations: Fe^{2*} (aq) + 2e⁻ ⇒ Fe(s) 1/2 O₂(g) + H₂O(l) + 2e⁻ ⇒ 2OH⁻ (aq) Calculate E^o, in V, for the spontaneous reaction using section 24 of the data booklet. (ii) Calculate the Gibbs free energy, ΔG^o, in kJ, which is released by the corrosion of 1 mole of iron. Use section 1 of the data booklet. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. (d) Finem9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.5mashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air. (a) Formulate an equation for the oxidation of nickel(II) sulfide to nickel(II) oxide. 		Deduce the overall redox equation for the reaction between acidic Sn ²⁺ (aq) and Cr ₂ O ₇ ²⁻ (aq), using section 24 of the data booklet. [1]
steps involve the following half-equations: Fe²+ (aq) + 2e⁻ ⇌ Fe (s) 1/2 O₂(g) + H₂O (l) + 2e⁻ ⇌ 2OH⁻ (aq) Calculate E⁰, in V, for the spontaneous reaction using section 24 of the data booklet. (ii) Calculate the Gibbs free energy, ΔG⁰, in kJ, which is released by the corrosion of 1 mole of iron. Use section 1 of the data booklet. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes.		
steps involve the following half-equations: Fe²+ (aq) + 2e⁻ ⇌ Fe (s) 1/2 O₂(g) + H₂O (l) + 2e⁻ ⇌ 2OH⁻ (aq) Calculate E⁰, in V, for the spontaneous reaction using section 24 of the data booklet. (ii) Calculate the Gibbs free energy, ΔG⁰, in kJ, which is released by the corrosion of 1 mole of iron. Use section 1 of the data booklet. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes.		
steps involve the following half-equations: Fe²+ (aq) + 2e⁻ ⇌ Fe (s) 1/2 O₂(g) + H₂O (l) + 2e⁻ ⇌ 2OH⁻ (aq) Calculate E⁰, in V, for the spontaneous reaction using section 24 of the data booklet. (ii) Calculate the Gibbs free energy, ΔG⁰, in kJ, which is released by the corrosion of 1 mole of iron. Use section 1 of the data booklet. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes.		
 \frac{1}{2} O_2(g) + H_2O(l) + 2e^- \infty 2OH^-(aq) Calculate E°, in V, for the spontaneous reaction using section 24 of the data booklet. (ii) Calculate the Gibbs free energy, ΔG°, in kJ, which is released by the corrosion of 1 mole of iron. Use section 1 of the data booklet. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. (chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.5mashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air. 	(b) (i)	
 \frac{1}{2} O_2(g) + H_2O(l) + 2e^- \infty 2OH^-(aq) Calculate E°, in V, for the spontaneous reaction using section 24 of the data booklet. (ii) Calculate the Gibbs free energy, ΔG°, in kJ, which is released by the corrosion of 1 mole of iron. Use section 1 of the data booklet. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. (chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.5mashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air. 		$Fe^{2+}(aq) + 2e^{-} \rightleftharpoons Fe(s)$
(ii) Calculate the Gibbs free energy, ΔG°, in kJ, which is released by the corrosion of 1 mole of iron. Use section 1 of the data booklet. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.		$\frac{1}{2}O_2(g) + H_2O(l) + 2e^- \rightleftharpoons 2OH^-(aq)$
1 mole of iron. Use section 1 of the data booklet. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.		
1 mole of iron. Use section 1 of the data booklet. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.	*****	
1 mole of iron. Use section 1 of the data booklet. (c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.		
(c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.	(ii	
(c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.	222222	
(c) Zinc is used to galvanize iron pipes, forming a protective coating. Outline how this process prevents corrosion of the iron pipes. Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.	*****	
process prevents corrosion of the iron pipes. Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org:o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.	*****	
Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org :o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.		
Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org :o) Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.		
Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.		***************************************
Millerite, a nickel sulfide mineral, is an important source of nickel. The first step in extracting nickel is to roast the ore in air.	 :hem 9 O#	156/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q4/ www.SmashingScience.org :o)
(a) Formulate an equation for the oxidation of nickel(II) sulfide to nickel(II) oxide.		a nickel sulfide mineral, is an important source of nickel. The first step in extracting
	nickel is t	mulate an equation for the oxidation of nickel(II) sulfide to nickel(II) oxide.
	nickel is t	rmulate an equation for the oxidation of nickel(II) sulfide to nickel(II) oxide.
	nickel is t	mulate an equation for the oxidation of nickel(II) sulfide to nickel(II) oxide.



(b) The nickel obtained from another ore, nickeliferous limonite, is contaminated with iron. Both nickel and iron react with carbon monoxide gas to form gaseous complexes, tetracarbonylnickel, Ni(CO)₄(g), and pentacarbonyliron, Fe(CO)₅(g). Suggest why the nickel can be separated from the iron successfully using carbon monoxide.

[1]

	• •	• •		• •		• •			*5		•	•	-	٠	*		*	٠	*	•	• •	•	*	•	•	•		**	•		• •	•	*	• •	*	÷	*	*	*	*	*	•		*:	• •		•
67		• •	• •		(*)		•		*	tot		•		ঃ		. :	č		:	•	• •		•		٠	•	•	*	•	*5	•				٠		*	•	*()	*	•	2)2	•	٠	•	. 1	•
					0.00			-00-0		***) e 1					. ,	 *		80				*							*		-				+			**	 	en c						

Topic Chem 9 Q# 157/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q3/ www.SmashingScience.org:o)

- 3. Vanadium has a number of different oxidation states.
 - (a) Determine the oxidation state of vanadium in each of the following species.

[2]

V ₂ O ₅ :	
VO ²⁺ :	

(b) Electrode potentials for the reactions of vanadium and other species are shown below.

$V^{3+}(aq) + e^- \rightleftharpoons V^{2+}(aq)$	-0.26 V
$VO^{2+}(aq) + 2H^{+}(aq) + e^{-} \rightleftharpoons V^{3+}(aq) + H_2O(l)$	+0.34V
$VO_2^+(aq) + 2H^+(aq) + e^- \rightleftharpoons VO^{2+}(aq) + H_2O(l)$	+1.0V

$Zn^{2+}(aq) + 2e^- \rightleftharpoons Zn(s)$	-0.76V
$Pb^{2+}(aq) + 2e^- \rightleftharpoons Pb(s)$	-0.13V
$SO_4^{2-}(aq) + 4H^+(aq) + 2e^- \rightleftharpoons H_2SO_3(aq) + H_2O(l)$	+0.17V
$Ag^{+}(aq) + e^{-} \rightleftharpoons Ag(s)$	+0.80V
$\frac{1}{2}Cl_2(g) + e^- \rightleftharpoons Cl^-(aq)$	+1.36V



3.	Vanadium h	as a	number	of differer	t oxidation	states.

1-1	Datamaina tha	id-ti	t-tf	alicema im	anala aftha	fellousia a anasia a
(a)	Determine the	oxidation s	iare or vana	acinim in i	each of the	following species.

[2]

V ₂ O ₅ :				
(*)*(*)*(*)*(*)	 	 *** *** *** ***	 	
\/O ² +-				
VO-:				

(b) Electrode potentials for the reactions of vanadium and other species are shown below.

$V^{3+}(aq) + e^- \rightleftharpoons V^{2+}(aq)$	-0.26V
$VO^{2+}(aq) + 2H^{+}(aq) + e^{-} \rightleftharpoons V^{3+}(aq) + H_2O(l)$	+0.34V
$VO_2^+(aq) + 2H^+(aq) + e^- \rightleftharpoons VO^{2+}(aq) + H_2O(l)$	+1.0V

$Zn^{2+}(aq) + 2e^- \rightleftharpoons Zn(s)$	-0.76V
$Pb^{2+}(aq) + 2e^{-} \rightleftharpoons Pb(s)$	-0.13V
$SO_4^{2-}(aq) + 4H^+(aq) + 2e^- \rightleftharpoons H_2SO_3(aq) + H_2O(l)$	+0.17V
$Ag^+(aq) + e^- \rightleftharpoons Ag(s)$	+0.80V
$\frac{1}{2}Cl_2(g) + e^- \rightleftharpoons Cl^-(aq)$	+1.36V

(i)	Identify, from the table, a non-vanadium species that can reduce VO2+ (aq) to)
	V ³⁺ (aq) but no further.	

[1]

(ii)	Identify, from the table, a non-vanadium species that could convert VO ₂ ⁺ (aq) to
	V^{2+} (aq).

[1]



 (c) (i) Formulate an equation for the reaction between VO²⁺(aq) and V²⁺(aq) in acidic solution to form V³⁺(aq). 	
 (ii) Comment on the spontaneity of this reaction by calculating a value for ΔG^Θ using the data given in (b) and in section 1 of the data booklet. 	
Chem 9 Q# 158/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/Q4/ www.SmashingScience.org:o)	
 (i) Magnesium chloride can be electrolysed. (i) Deduce the half-equations for the reactions at each electrode when molten magnesium chloride is electrolysed, showing the state symbols of the products. The melting points of magnesium and magnesium chloride are 922 K and 987 K respectively. 	
Anode (positive electrode):	
Cathode (negative electrode):	
(ii) Identify the type of reaction occurring at the cathode (negative electrode). [1]	N. C.



(iii) State the products when a very dilute aqueous solution of magnesium chlorid is electrolysed.	[2]
Anode (positive electrode):	
Cathode (negative electrode):	
hem 9 Q# 159/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/Q4/ www.SmashingScience (j) Standard electrode potentials are measured relative to the standard hydrogen electrode. Describe a standard hydrogen electrode.	ce.org :o) [2

***************************************	*/*/*(*)
 (k) A magnesium half-cell, Mg (s)/Mg²⁺ (aq), can be connected to a copper half-cell, Cu (s)/Cu²⁺ (aq). 	
 Formulate an equation for the spontaneous reaction that occurs when the circles completed. 	cuit [1
(ii) Determine the standard cell potential, in V, for the cell. Refer to section 24 of data booklet.	the [1
(iii) Predict, giving a reason, the change in cell potential when the concentration of copper ions increases.	of [2

Topic Chem 9 Q# 160/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/

- Ethane-1,2-diol, HOCH₂CH₂OH, has a wide variety of uses including the removal of ice from aircraft and heat transfer in a solar cell.
 - (c) Determine the average oxidation state of carbon in ethene and in ethane-1,2-diol.

Eth	en	e:																				
						 	 	 	 	 * *	 	 	 	 		 	cul,	a,	 	 	 	
Eth	an	e-	1,2-	dic	ol:																	
	2.0					 	 	 	 	 	 		 	 	21,	 	2 .			 	 	

Topic Chem 9 Q# 161/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/

- 1. Two IB students carried out a project on the chemistry of bleach.
 - (a) The bleach contained a solution of sodium hypochlorite, NaClO(aq). The students determined experimentally the concentration of hypochlorite ions, ClO, in the bleach:

Experimental procedure:

- The bleach solution was first diluted by adding 25.00 cm³ of the bleach to a 250 cm³ volumetric flask. The solution was filled to the graduation mark with deionized water.
- 25.00 cm³ of this solution was then reacted with excess iodide in acid.

$$ClO^{-}(aq) + 2l^{-}(aq) + 2H^{+}(aq) \rightarrow Cl^{-}(aq) + I_{2}(aq) + H_{2}O(1)$$

 The iodine formed was titrated with 0.100 mol dm⁻³ sodium thiosulfate solution, Na₂S₂O₃(aq), using starch indicator.

$$I_2(aq) + 2S_2O_3^{2-}(aq) \rightarrow 2\Gamma(aq) + S_4O_6^{2-}(aq)$$

The following data were recorded for the titration:

SIMA	First titre	Second titre	Third titre
Final burette reading of $0.100 \text{mol dm}^{-3} \text{Na}_2 \text{S}_2 \text{O}_3(\text{aq})$ (in cm ³ ± 0.05)	23.95	46.00	22.15
Initial burette reading of $0.100 \text{mol dm}^{-3} \text{Na}_2 \text{S}_2 \text{O}_3(\text{aq})$ (in cm ³ ± 0.05)	0.00	23.95	0.00




(i)	Deduce the oxidation states of chlorine and iodine in the following species.	[
	NaClO:	
	\mathbf{I}_{2} :	
(ii)	Deduce, with a reason, the oxidizing agent in the reaction of hypochlorite ions with iodide ions in part (a).	
(iii)	From a health and safety perspective, suggest why it is not a good idea to use hydrochloric acid when acidifying the bleach.	[1
	SMASHINGILL	



(iv) The thiosulfate ion, S₂O₃², is an interesting example of oxidation states. The sulfur atoms can be considered to have an oxidation state of +6 on one atom and -2 on the other atom. Discuss this statement in terms of your understanding of oxidation state.

[2]



Lewis (electron dot) structure of thiosulfate

*	+	*	+	*		0			*	+	*	•			*	*	+		6	•			•						•			•			-					*			•) •		9			+	-						
•		-			•		-				1	ne:	200	10	(A)	÷	*	Ž:	10	100	te	•	5	100	Đợi.	511	1		o to	e i	20	-	2115	100			-	o.T.	•	Ť.	Ť:		10.5		100	Vē	68	•	૽	0	d	*	•		1
-																																																							
*	•		•							•		-			*	-		***		P 4			60	0.00		.)(-				***	•				000			-			400						///					-			
-	to	 e to			 ***				-			Per c					2	700	ra e	10.0					7) (T		***	513								20		5.17				127	-		ı					
											4				-			i							-			2/4		10					112				3					-					4	2					
*			*					*	*	-	*	5			(%)	*	*5	•0	*	*35		*	40.7		53			60		*	*	*10				*	*		*	*:	•	50	63			3	V.S	*	*	15	*	-	*	*	•

(c) The various changes that have been made to the definitions of oxidation and reduction show how scientists often broaden similarities to general principles.

Combustion is also a redox type of reaction.

With reference to the combustion reaction of methane, explore two different definitions of oxidation, choosing one which is valid and one which may be considered not valid.

[2]

$$CH_4(g) + 2O_2(g) \rightarrow CO_2(g) + 2H_2O(1)$$

 	 	e e e:e:	 		
 	 		 	+ + + +	
 * * * *	 		 		

Topic Chem 9 Q# 162/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/

Tin(II) chloride is a white solid that is commonly used as a reducing agent.

(b) Tin can also exist in the +4 oxidation state.

$$Sn^{4+}(aq) + 2e^{-} \rightleftharpoons Sn^{2+}(aq)$$
 $E^{\Theta} = +0.15V$

Vanadium can be reduced from an oxidation state of +4 to +3 according to the equation:

$$VO^{2+}(aq) + 2H^{+}(aq) + e^{-} \rightleftharpoons V^{3+}(aq) + H_{2}O(l)$$
 $E^{\Theta} = +0.34 V$

(i) Calculate the cell potential, E^{Θ} , and the standard free energy, ΔG^{Θ} , change for the reaction between the VO²⁺ and Sn²⁺ ions, using sections 1 and 2 of the data

ron

	DOOKIEL.	[4]
	E [⊕] :	
	ΔG^{Θ} :	
	(ii) Deduce, giving your reason, whether a reaction between Sn ²⁺ (aq) and VO ²⁺ (aq) would be spontaneous.	[1]
1.	c Chem 9 Q# 163/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/ Phosphine (IUPAC name phosphane) is a hydride of phosphorus, with the formula PH ₃ .	
b)	(iii) State the oxidation state of phosphorus in P, and H,PO,-	[2]

P ₄ :					
H ₂ PO ₂ -:					



		(iv)	Oxidation is now defined in terms of change of oxidation number. Explore how earlier definitions of oxidation and reduction may have led to conflicting answers for the conversion of P ₄ to H ₂ PO ₂ ⁻ and the way in which the use of oxidation numbers has resolved this.	[3]
	4.30			
 opio	. Chen	n 9 O#	164/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/	Į.
١.			used the technique of titration to determine the concentration of ascorbic acid	
	(C ₆ H	H ₈ O ₆) ir	a sample of orange juice. Excess potassium iodide, KI (aq), was added to	
	in th	e pres	range juice. The resulting solution was titrated with potassium iodate, KIO ₃ (aq), ence of starch as an indicator. The end-point of the titration was shown by a colour.	
	Step	1	$IO_3^-(aq) + 5I^-(aq) + 6H^+(aq) \rightarrow 3I_2(aq) + 3H_2O(l)$	
	lodii	ne is o	nly slightly soluble in water; but in the presence of excess iodide ions, I ⁻ (aq),	
			e soluble tri-iodide ion, I ₃ (aq).	
	Step	2	$I_2(aq) + I^-(aq) \rightleftharpoons I_3^-(aq)$	
	oich	, 2	$I_2(aq) + I(aq) \leftarrow I_3(aq)$	
	Asc	orbic a	cid reacts with tri-iodide ions as follows.	
	Step	3	$C_6H_8O_6(aq) + I_3^-(aq) \rightarrow C_6H_8O_6(aq) + 2H^+(aq) + 3I^-(aq)$	
	(a)	(i)	Deduce the changes in oxidation number of iodine in step 1.	[2]
	. /			
			IO ₃ to I ₂ :	
			GRAD CHILDREN	
			Γ to I_2 :	
		(ii)	Identify the oxidizing and reducing agents in step 1.	[1]
			Oxidizing agent:	
			Reducing agent:	
				-3

	presence of an acid to form $Cr^{3+}(aq)$ and $IO_3^-(aq)$ ions. Deduce the half-equation for the reaction of I^- to IO_3^- and the overall equation for this reaction.	[2]
	Half-equation:	

	Overall equation:	

(ii)	Explain in terms of oxidation numbers whether iodine is oxidized or reduced in part (d) (i).	[1]
A vo		e.org :
A vo	9 Q# 166/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/Q8/ www.SmashingScience obtaic cell is constructed as follows. One half-cell contains a chromium electrode nersed in a solution containing Cr ³⁺ (aq) ions. The other half-cell contains a copper strode immersed in a solution containing Cu ²⁺ (aq) ions. The two electrodes are	e.org:
A vo	9 Q# 166/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/Q8/ www.SmashingScience obtains cell is constructed as follows. One half-cell contains a chromium electrode nersed in a solution containing Cr³+(aq) ions. The other half-cell contains a copper strode immersed in a solution containing Cu²+(aq) ions. The two electrodes are nected to a voltmeter and the two solutions by a salt bridge.	e.org:
A vo	9 Q# 166/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/Q8/ www.SmashingScience obtaic cell is constructed as follows. One half-cell contains a chromium electrode persed in a solution containing Cr³+(aq) ions. The other half-cell contains a copper strode immersed in a solution containing Cu²+(aq) ions. The two electrodes are nected to a voltmeter and the two solutions by a salt bridge. Voltmeter Voltmeter	e.org:

(ii)	Calculate the cell potential, in V, under standard conditions, for this voltaic cell, using table 14 of the data booklet and $E^{\Theta}_{cr^{3}}$ %cr = -0.74 V.	[1
(iii)	Predict the balanced equation for the spontaneous reaction which will produce a current in this voltaic cell.	[1]
(iv)	Identify the negative and the positive electrodes in this cell.	[1]
(v)	Predict the direction of movement of electrons in the external circuit.	[1]
(vi)	State the directions in which the negative ions (anions) and the positive ions (cations) flow in the salt bridge.	[1]

Topic Chem 9 Q# 167/ IB Chem/2015/s/TZ2/Paper 2 Section A/Higher Level/

- Electrolysis is an important industrial process used to obtain very reactive elements from their common ores.
 - (a) Molten magnesium chloride can be electrolysed using inert graphite electrodes at 800 °C.

Deduce the half-equations, including state symbols, for the reactions occurring at each electrode. (The melting points of ${\rm MgCl_2}$ and ${\rm Mg}$ are 714°C and 649°C respectively.)

[3]

	Positive electrode (anode):	
	Negative electrode (cathode):	
(b)	Aluminium can also be obtained by electrolysis. Suggest one reason why aluminium is often used instead of iron by engineers.	[1]

Topic Chem 9 Q# 168/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q7/ www.SmashingScience.org:o) The structure of ethanol is CH3CH2OH and for ethanal it is CH3CHO

		Ethanal:	
	(iii)	Deduce the half-equation for the oxidation of ethanol to ethanal.	[
	(iv)	Deduce the overall redox equation for the reaction of ethanol to ethanal with acidified potassium dichromate(VI).	[2
hem (i)	Bro	169/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q6/ www.SmashingScience mine can be produced by the electrolysis of molten sodium bromide. Juce the half-equation for the reaction at each electrode.	.org :o)
	Bro	169/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q6/ www.SmashingScience mine can be produced by the electrolysis of molten sodium bromide. duce the half-equation for the reaction at each electrode.	
	Pos	169/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q6/ www.SmashingScience mine can be produced by the electrolysis of molten sodium bromide. duce the half-equation for the reaction at each electrode.	
	Pos	169/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q6/ www.SmashingScience mine can be produced by the electrolysis of molten sodium bromide. duce the half-equation for the reaction at each electrode.	
	Pos Neg	169/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q6/ www.SmashingScience mine can be produced by the electrolysis of molten sodium bromide. duce the half-equation for the reaction at each electrode.	
(i)	Pos Neg Precon	169/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q6/ www.SmashingScience mine can be produced by the electrolysis of molten sodium bromide. duce the half-equation for the reaction at each electrode. Sitive electrode (anode): gative electrode (cathode):	[2]
(i)	Pos Neg Precon	169/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q6/ www.SmashingScience mine can be produced by the electrolysis of molten sodium bromide. duce the half-equation for the reaction at each electrode. Sitive electrode (anode): gative electrode (cathode): dict the products formed at the electrodes during the electrolysis of icentrated aqueous sodium bromide.	[2]
(i)	Pos Neg Pre con	169/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q6/ www.SmashingScience mine can be produced by the electrolysis of molten sodium bromide. duce the half-equation for the reaction at each electrode. Sitive electrode (anode): dict the products formed at the electrodes during the electrolysis of identrated aqueous sodium bromide.	[2]

(ii) Determine the oxidation number of carbon in ethanol and ethanal.

[2]

		$Br_2(aq) + 2NaI(aq) \rightarrow I_2(aq) + 2NaBr(aq)$	
	Iden	ntify the oxidizing agent in this reaction.	[1]
	• • •		• • •
)	(i)	Define the term standard electrode potential, E^{\ominus} .	[1]
	(ii)	Draw a labelled diagram for the voltaic cell in which the following reaction occurs $Mg(s) + Cu^{2+}(aq) \rightarrow Mg^{2+}(aq) + Cu(s)$	urs.
	(ii)		urs.

(iii) A student measures a voltage of 2.65 V in the voltaic cell formed between magnesium and copper half-cells using a digital voltmeter.

State the random uncertainty of this value, in V, and the number of significant figures in the answer.

[2]

Rar	ndo	m ı	ınc	erta	ain	ty:																		
						• •	 					 *			•	 100				٠				
Sig	nifi	can	t fic	jure	es:																			

	(iv)	Outli	ne how the student can reduce the random error in her results. [1]	
opio 3.	Prop	ene is	Q# 171/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q4. NOT with Q3b(i) an important starting material for many products. The following shows some ds which can be made from propene, C ₃ H ₆ .	
			Propene $(C_3H_6) \rightarrow C_3H_7Cl \rightarrow C_3H_8O \rightarrow C_3H_6O$	
	(a)	Con	sider the conversion of propene to C ₃ H ₇ Cl.	
		(i)	State the type of reaction.	[1]
	41.47¥			
		(ii)	State the IUPAC name of the major product.	[1]
	\$ (\$1.\$)			
		(iii)	Outline why it is the major product.	[1]
	5.5.5			
		(iv)	Write an equation for the reaction of the major product with aqueous sodium hydroxide to produce a C ₃ H ₈ O compound, showing structural formulas.	[1]

(b) An experiment was carried out to determine the order of reaction between one of the isomers of C₃H₇Cl and aqueous sodium hydroxide. The following results were obtained.

Experiment	[C ₃ H ₇ Cl] / mol dm ⁻³	[OHT] / mol dm ⁻³	Initial rate / mol dm ⁻³ s ⁻¹
1	0.05	0.10	3.1 × 10 ⁻⁴
2	0.10	0.20	1.3 × 10 ⁻³
3	0.15	0.10	9.2 × 10 ⁻⁴

The rate equation for this reaction is: Rate = $k [C_3H_7Cl][OH^-]$



		(ii)	Deduce the type of mechanism for the reaction of this isomer of C ₃ H ₇ Cl with aqueous sodium hydroxide.	[1]
		(iii)	Sketch the mechanism using curly arrows to represent the movement of electrons.	[4]
	(d)	(i)	State the reagents for the conversion of the compound C_3H_8O formed in (a)(iv) into C_3H_8O .	[1]
	* ***	* ***		
u		(ii)	Explain why the compound C_3H_8O , produced in (a)(iv), has a higher boiling point than compound C_3H_8O , produced in d(i).	[2]
	18 828	****		
	2.52			
ı	(e)		pene is often polymerized. Draw a section of the resulting polymer, showing two eating units.	[1]



9.	Halo	genoalkanes undergo nucleophilic substitution reactions with sodium hydroxide.	
	(a)	State a reason why most halogenoalkanes are more reactive than alkanes.	[1]
	(b)	Classify 1-bromopropane as a primary, secondary or tertiary halogenoalkane, giving a reason.	[1]
	1111		
	(c)	Explain the mechanism of the reaction between 1-bromopropane with aqueous sodium hydroxide using curly arrows to represent the movement of electron pairs.	[4]
	(d)	State, giving your reason, whether the hydroxide ion acts as a Lewis acid, a Lewis base, or neither in the nucleophilic substitution.	[1]

	(e)	Suggest two advantages of understanding organic reaction mechanisms.	[2]

	* * * *		

Topic Chem 10 Q# 172/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q9/ www.SmashingScience.org :o)

Topic Chem 10 Q# 173/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q8/ www.SmashingScience.org :o)

Carbon dioxide contributes significantly to global warming. It can be used as a raw material with methyloxirane to form polymers.

$$n CO_2 + n HC O CH_2 \longrightarrow \begin{bmatrix} O & O & CH CH_2 \\ CH_3 & CH_3 & CH_3 \end{bmatrix}$$

methyloxirane

(;	a)	Sug	gest w	hy the three-membered ring in methyloxirane is unstable.	[1]
(1	b)	(i)	Drav	v two structural isomers of methyloxirane.	[2]
		(ii)	State	e, giving a reason, whether methyloxirane can form cis-trans isomers.	[1]



Topic Chem 10 Q# 174/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q7/ www.SmashingScience.org :o)

7. Consider the following Hess's law cycle:

$$CH_2 = CH_2(g) + H_2(g) \xrightarrow{Step 1} CH_3CH_3(g)$$

$$+ \frac{7}{2} O_2(g) \\ Step 2 \\ + \frac{7}{2} O_2(g) \\ Step 3$$

$$2CO_2(g) + 3H_2O(l)$$

	(a)	Identify the type of reaction in step 1.	[1
Copic		10 Q# 175/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q6/ www.SmashingScience.org :o) anoic acid, CH ₃ CH ₂ CH ₂ COOH, is a weak acid and ethylamine, CH ₃ CH ₂ NH ₂ , is a weak base.	
	(d)	Explain why butanoic acid is a liquid at room temperature while ethylamine is a gas at room temperature.	[2]
	•••		
Copic		10 Q# 176/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q6/ www.SmashingScience.org :o) anoic acid, CH ₃ CH ₂ CH ₂ COOH, is a weak acid and ethylamine, CH ₃ CH ₂ NH ₂ , is a weak base.	
	(e)	(i) State a suitable reagent for the reduction of butanoic acid.	[1]

33		(ii) Deduce the product of the complete reduction reaction in (e)(i).	[1]



Topic Chem 10 Q# 177/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q7/ www.SmashingScience.org:o)
7. This question is about the reactions of halogenoalkanes.

Compare and contrast the mechanisms by which 1-chlorobutane, $CH_3CH_2CH_2CH_2CI$, and 2-chloro-2-methylpropane, $(CH_3)_3CCI$, react with aqueous sodium hydroxide,

giving two similarities and one difference.	[3]
Two similarities:	
One difference:	
(b) Outline why the rate of reaction of the similar bromo-compounds is faster.	[1]
(c) (i) State the organic product of the reaction between 1-chlorobutane, CH ₃ CH ₂ CH ₂ CI, and aqueous sodium hydroxide.	[1]
(ii) Suggest how this product could be synthesized in one step from butanoic acid.	[1]



	(III)	reacts with butanoic acid. [1]	

opio	Chem 10 Q	# 178/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q3/ www.SmashingScience.org :o)	
3.	This que	stion is about ethene, C ₂ H ₄ , and ethyne, C ₂ H ₂ .	
	(a) (i)	Ethyne, like ethene, undergoes hydrogenation to form ethane. State the conditions required.	[2]

	(ii)	Outline the formation of polyethene from ethene by drawing three repeating units of the polymer.	[1]
		ryne reacts with chlorine in a similar way to ethene. Formulate equations for the owing reactions.	[2]
	One mole	e of ethyne reacts with one mole of chlorine:	
	One mole	e of benzene reacts with one mole of chlorine:	



Topic Chem 10 Q# 179/ IB Chem/2017/s/TZ 2/Paper 2/Higher Level/Q6/ www.SmashingScience.org :o)

- 6. The photochemical chlorination of methane can occur at low temperature.
 - (a) Using relevant equations, show the initiation and the propagation steps for this reaction.

[3]

Initiation:			
Propagation:			


(b) The overall equation for monochlorination of methane is:

$$CH_4(g) + Cl_2(g) \rightarrow CH_3Cl(g) + HCl(g)$$

Calculate the standard enthalpy change for the reaction, ΔH^{\oplus} , using section 12 of the data booklet.

[2]

(c) Polyvinyl chloride (PVC) is a polymer with the following structure.



State the structural formula for the monomer of PVC.

[1]





Topic ((v) Identify one organic functional group that can react with acidified K ₂ Cr ₂ O ₇ (aq).	[1
Горіс (7 .	Chem 10 Q# 181/ IB Chem/2017/s/TZ 1/Paper 2/Higher Level/Q7/ www.SmashingScience.org :o) Benzene is an aromatic hydrocarbon.	
	(a) Discuss the physical evidence for the structure of benzene.	[2]
<u> </u>	(b) State the typical reactions that benzene and cyclohexene undergo with bromine.	[1]
	Benzene:	
	Cyclohexene:	
	(c) State the reagents used to convert benzene to nitrobenzene and the formula of the electrophile formed.	[2]
	(d) Explain the mechanism for the nitration of benzene, using curly arrows to show the movement of electron pairs.	[4]
		Ĭ

(b) When one enantiomer undergoes substitution by alkaline hydrolysis approximately 75% of the product molecules show inversion of configuration. Comment on the		(e)	State the reagents used in the two-stage conversion of nitrobenzene to aniline.	[2]
One structural isomer of C ₄ H ₅ Br is a chiral molecule. (a) Draw the three-dimensional shape of each enantiomer of this isomer showing their spatial relationship to each other. (b) When one enantiomer undergoes substitution by alkaline hydrolysis approximately 75 % of the product molecules show inversion of configuration. Comment on the mechanisms that occur. (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater				
One structural isomer of C ₄ H ₅ Br is a chiral molecule. (a) Draw the three-dimensional shape of each enantiomer of this isomer showing their spatial relationship to each other. (b) When one enantiomer undergoes substitution by alkaline hydrolysis approximately 75 % of the product molecules show inversion of configuration. Comment on the mechanisms that occur. (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater				
One structural isomer of C ₄ H ₅ Br is a chiral molecule. (a) Draw the three-dimensional shape of each enantiomer of this isomer showing their spatial relationship to each other. (b) When one enantiomer undergoes substitution by alkaline hydrolysis approximately 75 % of the product molecules show inversion of configuration. Comment on the mechanisms that occur. (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater		9.07		
(a) Draw the three-dimensional shape of each enantiomer of this isomer showing their spatial relationship to each other. (b) When one enantiomer undergoes substitution by alkaline hydrolysis approximately 75 % of the product molecules show inversion of configuration. Comment on the mechanisms that occur. (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater	ic	Chem	10 Q# 182/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/	
(b) When one enantiomer undergoes substitution by alkaline hydrolysis approximately 75% of the product molecules show inversion of configuration. Comment on the mechanisms that occur.		One	structural isomer of C ₄ H ₉ Br is a chiral molecule.	
75 % of the product molecules show inversion of configuration. Comment on the mechanisms that occur. [2] (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater		(a)		[2
75 % of the product molecules show inversion of configuration. Comment on the mechanisms that occur. [2] (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater				
75 % of the product molecules show inversion of configuration. Comment on the mechanisms that occur. [2] (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater				
75 % of the product molecules show inversion of configuration. Comment on the mechanisms that occur. [2] (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater				
75% of the product molecules show inversion of configuration. Comment on the mechanisms that occur. [2] (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater				
75% of the product molecules show inversion of configuration. Comment on the mechanisms that occur. [2] (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater				
75 % of the product molecules show inversion of configuration. Comment on the mechanisms that occur. [2] (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater				
75 % of the product molecules show inversion of configuration. Comment on the mechanisms that occur. [2] (c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater			1-2-2-110	
(c) Suggest why the rate of alkaline hydrolysis of an enantiomer of iodopropane is greater		(b)		
				[2
		15500		

		(c)	Suggest why the rate of alkaline hydrolysis of an enantiomer of indopropage is greater	
		(0)	than that of an enantiomer of bromopropane.	[:
			• • • • • • • • • • • • • • • • • • • •	
		22.2		

a) Draw the	full structural formulas of propane and propene.	[1]
Propane:		
Propene:		
	pane and propene react with bromine. te an equation and the condition required for the reaction of 1 mol of propane	
(i) Sta	te an equation and the condition required for the reaction of 1 mol of propane 1 mol of bromine.	[2]
(i) Sta	te an equation and the condition required for the reaction of 1 mol of propane	[2]
(i) Sta	te an equation and the condition required for the reaction of 1 mol of propane	[2]
(i) Sta	te an equation and the condition required for the reaction of 1 mol of propane	[2]
(i) Sta	te an equation and the condition required for the reaction of 1 mol of propane	[2]
(i) Sta	te an equation and the condition required for the reaction of 1 mol of propane in 1 mol of bromine.	
(i) Sta	te an equation and the condition required for the reaction of 1 mol of propane in 1 mol of bromine.	
(ii) Sta	te an equation and the condition required for the reaction of 1 mol of propane in 1 mol of bromine.	

	d) Construct the mechanism of the formation of 2-bromopropane from hydrogen b and propene using curly arrows to denote the movement of electrons.	
_		
c (hem 10 Q# 184/IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/Q7/ www.Smashi	ingScience.or
	an alcohol with the following formula:	ingScience.or
		ingScience.or
	an alcohol with the following formula: H H H	ingScience.or
	an alcohol with the following formula:	ingScience.or
	An alcohol with the following formula: H H H C——————————————————————————————	ingScience.or
	an alcohol with the following formula: H H H C——————————————————————————————	ingScience.or
	An alcohol with the following formula: H H H C——————————————————————————————	ingScience.or
is	an alcohol with the following formula: H H H C——————————————————————————————	ingScience.or
is an	an alcohol with the following formula: H H H C C C H H O H Somer of X but has a different functional group Both X and Y are soluble in water. Deduce whether or not both X and Y show	ingScience.or
is an	An alcohol with the following formula: H H H C C C H H O H Somer of X but has a different functional group Both X and Y are soluble in water. Deduce whether or not both X and Y show hydrogen bonding with water molecules, representing any hydrogen bonding	
(is	An alcohol with the following formula: H H H C C C H H O H Somer of X but has a different functional group Both X and Y are soluble in water. Deduce whether or not both X and Y show hydrogen bonding with water molecules, representing any hydrogen bonding	ingScience.or
is an	An alcohol with the following formula: H H H C C C H H O H Somer of X but has a different functional group Both X and Y are soluble in water. Deduce whether or not both X and Y show hydrogen bonding with water molecules, representing any hydrogen bonding	
is	An alcohol with the following formula: H H H C C C H H O H Somer of X but has a different functional group Both X and Y are soluble in water. Deduce whether or not both X and Y show hydrogen bonding with water molecules, representing any hydrogen bonding	
is	an alcohol with the following formula: H H H C C C H H O H Somer of X but has a different functional group Both X and Y are soluble in water. Deduce whether or not both X and Y show hydrogen bonding with water molecules, representing any hydrogen bonding present by means of a diagram.	



(iv)	Compare and contrast the bonding types formed by nitrogen in [PtCl ₂ (NH ₃) ₂].	[2]
	Similarity:	
	Difference:	

(v)	Deduce all of the intermolecular forces present for the ammonia molecule.	[2]


	1 person 1	\square

Topic Chem 10 Q# 185/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/Q7/ www.SmashingScience.org:o)

Topic Chem 10 Q# 186/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/

- The biopharmaceutical industry is now a global contributor to the world economy.
 - (a) Atorvastatin, a drug used to lower cholesterol, recently gained attention from the global media.

Atorvastatin has the structure shown below.



Identify the four functional groups, I, II, III and IV.

[2]

- (b) Bute is a chemical with the following molecular formula: C19H20N2O2
 - (iii) Deduce the degree of unsaturation (index of hydrogen deficiency IHD) of bute. [1]



(i)	Diaw dom isomer	ers of the complex.	[1]
(ii)	Explain the polar your answer.	rity of each isomer, using a diagram for each isomer to suppor	t [2]
			a a
			6

12	**********		
		m/2016/s/TZoSP/Paper 2 Section A/Higher Level/	
1-100	doethane reacts wit	th sodium hydroxide.	
	doethane reacts wit Explain the mech		
1-100	doethane reacts wit Explain the mech	th sodium hydroxide. thanism of this reaction, using curly arrows to represent the move and showing any stereochemical features of the reaction mechanical	
1-100	doethane reacts wit Explain the mech	th sodium hydroxide. hanism of this reaction, using curly arrows to represent the move	
1-100	doethane reacts wit Explain the mech	th sodium hydroxide. thanism of this reaction, using curly arrows to represent the move and showing any stereochemical features of the reaction mechanical	
1-100	doethane reacts wit Explain the mech	th sodium hydroxide. thanism of this reaction, using curly arrows to represent the move and showing any stereochemical features of the reaction mechanical	
1-100	doethane reacts wit Explain the mech	th sodium hydroxide. thanism of this reaction, using curly arrows to represent the move and showing any stereochemical features of the reaction mechanical	
1-100	doethane reacts wit Explain the mech	th sodium hydroxide. thanism of this reaction, using curly arrows to represent the move and showing any stereochemical features of the reaction mechanical	
1-100	doethane reacts wit Explain the mech	th sodium hydroxide. thanism of this reaction, using curly arrows to represent the move and showing any stereochemical features of the reaction mechanical	
1-100	doethane reacts wit Explain the mech	th sodium hydroxide. thanism of this reaction, using curly arrows to represent the move and showing any stereochemical features of the reaction mechanism.	


(d) The two isomers of $[Pt(NH_3)_2Cl_2]$ are crystalline. One of the isomers is widely used as a

(b)	State the rate expression for this reaction and identify the molecularity of the rate- determining step (RDS).	1
	Rate expression:	
	Molecularity of RDS:	
	ggest why polar, aprotic solvents are more suitable for S_N2 reactions whereas polar, this solvents favour S_N1 reactions. [2]	
S _N 2		
900		

6 1	153-30-19	
S_N^1	11/	
-		
	164 - 159 A T	
	WALES / L	
Б	to a side of the second of the	
	duce, with a reason, if water or DMF (N,N-Dimethylformamide, HCON(CH ₃) ₂) is a ter solvent for this reaction. [1]	
000	[2]	
• • •		
3.53		

Topic Chem 10 Q# 188/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/

Compound A and compound B are hydrocarbons.



(a) (i) State the term that is used to describe molecules that are related to each other in the same way as compound A and compound B.

[1]

 Suggest a chemical test to distinguish between compound A and compound B, giving the observation you would expect for each.

[2]

Test:	
Observation with A:	

Observation with B:	

Topic Chem 10 Q# 189/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/


 Phosgene, COCl₂, is usually produced by the reaction between carbon monoxide and chlorine according to the equation:

$$CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$$



(b) One important industrial use of phosgene is the production of polyurethanes. Phosgene is reacted with diamine X, derived from phenylamine.

Classify diamine X as a primary secondary or tertiary amine



(ii)	Phenylamine, C ₆ H ₅ NH ₂ , is produced by the reduction of nitrobenzene, C ₆ H ₅ NO ₂ . Suggest how this conversion can be carried out.



(iii) Nitrobenzene can be obtained by nitrating benzene using a mixture of concentrated nitric and sulfuric acids. Formulate the equation for the equilibrium established when these two acids are mixed.

[1]




(iv) Deduce the mechanism for the nitration of benzene, using curly arrows to indicate the movement of electron pairs.

[4]

[1]



(c) The other monomer used in the production of polyurethane is compound Z shown below.



(i)	State the name, applying IUPAC rules, of compound Z and the class of
	compounds to which it belongs.

[2]

Name:	
Class:	

Topic Chem 10 Q# 190/ IB CHEM/2015/w/TZ0/Paper 2 Section B/Higher Level/

9. (a) A 0.842 g sample of a liquid halogenoalkane, RBr(l), was heated under reflux with 1.35 × 10⁻² mol of aqueous sodium hydroxide, NaOH (aq). After cooling the mixture, the excess NaOH was titrated with hydrochloric acid, HCl (aq), and required 7.36 × 10⁻³ mol of the acid.

(i)	State the equation for the substitution reaction of the halogenoalkane with sodium
	hydroxide.

	•	7
1	8	

VICE VI	

(ii) Calculate the amount, in mol, of sodium hydroxide that reacted with the halogenoalkane. [1]





(iii)	Calculate the molar mass of the halogenoalkane.
	,
(iv)	Given that each molecule of the halogenoalkane contains one bromine atom, determine its molecular formula.

base tertia (If yo	ou have not been able to determine the molecular formula in part (a) (iv), use
C.H.	Br to deduce the four structural isomers.)
051	The state of the s
051	
05.1	
reacti	on between a primary halogenoalkane drawn in part (a) (v) and potassium llows a S _N 2 mechanism.
reacti	on between a primary halogenoalkane drawn in part (a) (v) and potassium
reacti	on between a primary halogenoalkane drawn in part (a) (v) and potassium flows a S _N 2 mechanism.

(b)

(ii)	Explain the mechanism of the reaction using curly arrows to represent the movement of electron pairs.	
		_
(iii)	The organic product obtained in part (b) (ii) can be reduced to form an amine. State an equation for this reaction and a suitable catalyst.	
is mole	cule is needed in the question that follows it:	
	CH ₃	
C	=c	
/	CU	
	CH ₃	
(ii)	Under certain conditions, the major product obtained in the elimination reaction	
	can undergo polymerization. Identify the type of polymerization and draw a section of the polymer consisting of two repeating units.	
	Type of polymerization:	
		*
	Section of polymer:	

Ethane can react with chlorine. Explain the free-radical mechanism of this including any necessary reaction conditions.	[4

***************************************	* * * * * * * * *

Chem 10 Q# 191/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/	
Some reactions of but-2-ene are given below.	
Same reasons of Date 2 and Same Same	
Poly(but-2-ene)	
A	
Br.(I) concentrated H.SO	
Compound A \longrightarrow Br ₂ (l) H ₃ C—CH=CH—CH ₃ $\xrightarrow{\text{concentrated H}_2SO}$ H ₂ O(l)	→ C4HOH
in the absence But-2-ene H ₂ O(l)	C ₄ H ₉ OH Compound C
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(l)	→ C4HOH
compound A H_3C —CH $=$ CH $=$ CH $_3$ in the absence But-2-ene	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(l)	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(l) HBr	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(l) HBr H ₃ C—CHBr—CH ₂ —CH ₃	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(l) H ₂ O(l)	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(I) H ₂ O(I) HBr H ₃ C—CHBr—CH ₂ —CH ₃ Compound B	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(I) H ₃ C—CHBr—CH ₂ —CH ₃	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(I) H ₂ O(I) HBr CH ₂ CHBr—CH ₂ —CH ₃ Compound B	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(I) H ₂ O(I) HBr CH ₂ CHBr—CH ₂ —CH ₃ Compound B	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(I) H ₂ O(I) HBr H ₃ C—CHBr—CH ₂ —CH ₃ Compound B	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(I) H ₂ O(I) HBr H ₃ C—CHBr—CH ₂ —CH ₃ Compound B	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(I) H ₂ O(I) HBr H ₃ C—CHBr—CH ₂ —CH ₃ Compound B	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(I) H ₂ O(I) HBr H ₃ C—CHBr—CH ₂ —CH ₃ Compound B	→ C4HOH
in the absence of UV H ₃ C—CH=CH—CH ₃ H ₂ O(I) H ₂ O(I) HBr H ₃ C—CHBr—CH ₂ —CH ₃ Compound B	→ C4HOH

	(iii) Describe the colour change observed when excess but-2-ene reacts with bromin to form compound A.	15
		*
(i)	Outline two reasons why the polymerization of alkenes is of economic importance.	
(ii)	Identify the structure of the repeating unit of poly(but-2-ene).	
(i)	Compound C , C ₄ H ₉ OH, can also be formed by reacting compound B , CH ₃ CHBrCH ₂ CH ₃ , with aqueous potassium hydroxide. This reaction proceeds by both S _N 1 and S _N 2 mechanisms. Explain the S _N 2 mechanism, using curly arrows to represent the movement of electron pairs.	
	SMASHINGILL	

[1]

(ii)	Explain why the hydroxide ion is a better nucleophile than water.	[2
(i)	Compound B , CH ₃ CHBrCH ₂ CH ₃ , also reacts with potassium cyanide. Apply IUPAC rules to name the organic product formed.	[1]
(ii)	The organic product of the reaction in part (d) (i) can be reduced to: $ \begin{array}{cccccccccccccccccccccccccccccccccc$	[1]
(iii)	Deduce the full structural formula of the organic product formed when the compound in part (d) (ii) reacts with ethanoic acid in the presence of an acid catalyst.	[1]
	SMASHINGI	

(e)		npound C , C ₄ H ₉ OH, can be oxidized by acidified potassium dichromate(VI) to form pound F .	
	(i)	State the name of the functional group present in compound F .	[1]
	(ii)	Deduce the structural formula of an alcohol which is a structural isomer of compound C and cannot be oxidized by acidified potassium dichromate(VI).	[1]
(f)	Exp	ain why but-2-ene is more volatile than compound C.	[2]
	2/1/2 2/1/2		
(g)	Ded	uce the equation for the complete combustion of compound C.	[1]
	4.4.4		
		A A A A A A A A A A A A A A A A A A A	

OI St	ereoisomensm.	
(i)	Define the term stereoisomers.	[1
(ii)	State the conditions needed for a compound to show geometrical isomerism.	[2
(iii)	Draw the structures of the two geometrical isomers of but-2-ene, clearly identifying each as <i>cis</i> or <i>trans</i> .	[2

But-2-ene can exist as two geometrical isomers. Geometrical isomerism is a form

(a)	10 Q# 192/ IB Chem/2015/s/TZ2/Paper 2 Section A/Higher Level/ State two features of a homologous series.
(b)	Ethane, a member of the homologous series of alkanes, can react with bromine. Explain the free-radical mechanism of this reaction, including any necessary reaction conditions.

	110 Q# 193/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/ anol has many industrial uses. State an equation for the formation of ethanol from ethene and the necessary reaction conditions.
C. H	Equation:
	Equation:
	Equation: Conditions:



(a)		e the equation for the acid-catalysed reaction of ethanol with propanoic acid and the name of the organic product.	[2]
	Equ	ation:	
	Nan	ne of the organic product:	
	× * *	***************************************	
(e)	(i)	A polyester can be formed when ethane-1,2-diol reacts with benzene-1,4-dicarboxylic acid.	
		Deduce the structure of the repeating unit and state the other product formed.	[2]
		Repeating unit:	
		Other product:	
	(ii)	State the type of polymerization that occurs.	[1]
		SMASHING!!!	

Patrick Brannac

Topic Chem 10 Q# 194/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/

Bromomethane was used as a pesticide until it was found to be ozone-depleting.

(a)	State	e the equation for the reaction between methane and bromine to form bromomethane.	[1]
	* * *		
(b)	(i)	Explain, using equations, the complete free-radical mechanism for the reaction of methane with bromine, including necessary reaction conditions.	[4
	(ii)	Bromomethane reacts with aqueous sodium hydroxide. State the organic product of this reaction.	[1
(c)	Expl	lain why the rate of the reaction between iodomethane, CH ₃ I, and NaOH(aq) is er than the rate of the reaction between CH ₃ Br and NaOH(aq).	[2]
	* * *		



Topic Chem 10 Q# 195/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/ Ethanol is a primary alcohol that can be oxidized by acidified potassium dichromate(VI). Distinguish between the reaction conditions needed to produce ethanal and ethanoic acid. [2] Ethanal: Ethanoic acid: Ethanol can be made by reacting aqueous sodium hydroxide with bromoethane. Explain the mechanism for this reaction, using curly arrows to represent the movement [4] of electron pairs. 2-bromobutane exists as optical isomers. (d) (i) State the essential feature of optical isomers. [1] [2] (11) Outline how a polarimeter can distinguish between these isomers.

Topic Chem 11 Q# 196/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q4.NOT with Q3c


 Propene is an important starting material for many products. The following shows some compounds which can be made from propene, C₃H₆.

Propene
$$(C_3H_8) \rightarrow C_3H_7Cl \rightarrow C_3H_8O \rightarrow C_3H_8O$$

(b) An experiment was carried out to determine the order of reaction between one of the isomers of C₃H₇Cl and aqueous sodium hydroxide. The following results were obtained.

Experiment	[C ₃ H ₇ CI] / mol dm ⁻³	[OHT] / mol dm ⁻³	Initial rate / mol dm ⁻³ s ⁻¹
1	0.05	0.10	3.1 × 10 ⁻⁴
2	0.10	0.20	1.3×10^{-3}
3	0.15	0.10	9.2 × 10 ⁻⁴

C₃H₆O mentioned in this question has a molecular formula of CH₃COCH₃



[1]



Topic Chem 11 Q# 197/ IB Chem/2019/w/TZ0/Paper 2/Higher Level/Q3./ www.SmashingScience.org:0)

 The biochemical oxygen demand of a water sample can be determined by the following series of reactions. The final step is titration of the sample with sodium thiosulfate solution, Na₂S₂O₃(aq).

$$2Mn^{2+}(aq) + O_2(aq) + 4OH^{-}(aq) \rightarrow 2MnO_2(s) + 2H_2O(l)$$

 $MnO_2(s) + 2\Gamma(aq) + 4H^{+}(aq) \rightarrow Mn^{2+}(aq) + I_2(aq) + 2H_2O(l)$
 $2S_2O_3^{2-}(aq) + I_2(aq) \rightarrow 2\Gamma(aq) + S_4O_8^{2-}(aq)$

A student analysed two $300.0\,\mathrm{cm^3}$ samples of water taken from the school pond: one immediately (day 0), and the other after leaving it sealed in a dark cupboard for five days (day 5). The following results were obtained for the titration of the samples with $0.0100\,\mathrm{mol\,dm^{-3}\,Na_2S_2O_3(aq)}$.

Sample	Titre / $\mathrm{cm^3}\pm0.1\mathrm{cm^3}$
Day 0	25.8
Day 5	20.1




(ii) Suggest a modification to the procedure that would make the results more reliable.	[1]
opic Chem 11 Q# 198/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q8/ www.SmashingScience.org :o) (c) Predict the chemical shift and splitting pattern of the signal produced by the hydrogen atoms labelled X in the ¹ H NMR spectrum of the polymer. Use section 27 of the data booklet. CH ₃ CH ₃ CH ₃ n	[2]
Chemical shift:	
Splitting pattern:	


An organic compound containing carbon, hydrogen and oxygen has 62.02% carbon and 10.43% hydrogen by mass.

lt's empirical formula is C₃H₆O










[Source: NIST Mass Spec Data Center, S.E. Stein, director, "Mass Spectra" in NIST Chemistry WebBook, NIST Standard Reference Database Number 69, Eds. P.J. Linstrom and W.G. Mallard, National Institute of Standards and Technology, Gaithersburg MD, 20899]

The mass spectrum of the compound is shown. Deduce the relative molecular mass of the compound.





[Source: NIST Mass Spec Data Center, S.E. Stein, director, "Mass Spectra" in NIST Chemistry WebBook, NIST Standard Reference Database Number 69, Eds. P.J. Linstrom and W.G. Mallard, National Institute of Standards and Technology, Gaithersburg MD, 20899]

The compound could not be oxidized using acidified potassium dichromate(VI).

Deduce the structural formula of the compound.

[1]

Topic Chem 11 Q# 200/ IB Chem/2018/w/TZ 0/Paper 2/Higher Level/Q1/ www.SmashingScience.org :o)

 3.26 g of iron powder are added to 80.0 cm³ of 0.200 mol dm⁻³ copper(II) sulfate solution. The following reaction occurs:

$$Fe(s) + CuSO_4(aq) \rightarrow FeSO_4(aq) + Cu(s)$$

- (b) (i) The reaction was carried out in a calorimeter. The maximum temperature rise of the solution was 7.5 °C.
 - (iii) The only significant uncertainty is in the temperature measurement.

Determine the absolute uncertainty in the calculated value of ΔH if the uncertainty in the temperature rise was ± 0.2 °C. [2]

					27.			-				2 2								201		-										-			-				2.6			4
8.5	ţ	•	 •		7 .3	200				10.5		7	-5						 	*			 e to		đạ.		le.t.	•		•	•	•	30	n t	্ৰ	•	:::			•		ē
	100		 •		-		-				-		 -				100		 	 *			 ٠					+		٠		20			-	•			-		-	
2.3	1913		 - 1		20	1		4	2:4	50-2	2%		1	26	1 1	-	150	2		 21	- 1	-		٠.	 157	- 12			1 1	(a)	20	20			ŭ.	10	202	- 2	2		 121	2

Topic Chem 11 Q# 201/ IB Chem/2018/s/TZ 2/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o)

 (g)	Outline	why repea	ting quantitative	measuremen	its is impor	tant.		
				-	JUL OF		-	

Topic Chem 11 Q# 202/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q2/ www.SmashingScience.org: o)

(b)	Describe how the relative atomic mass of a sample of calcium could be determined
	from its mass spectrum.

+			*	-			-	•	•		•	-	•	*	+	5			*	*	*	*	-	#il			iż	-	0.0	16	*	*	-	H	*	•	*	*		73.		9	-	- 0	*		- 0			+	- 19	•	02					•) •	*	5 3	•	*	-	*	7	+
														\$ 9			04	1	2		2	40		-		+	34							¥		¥	2	-		114			- 12			4.5	- 7		. 3						. ;									*		
•			1	-	•	•	•					•	-	-					-	-	1	-		*	*	*	*	-			•			*	*		*	-	-												-									1			*	*	•	•
	2.59	÷		-	-	-	-	-		-	•	-		+13					-		-	-	-	-						- 13					+			- *		10	. ,			 - 17			-	-00		+==	. ,			 .,.		-	*:0*	-0.5	*11	٠.		-	-	*		-


Topic Chem 11 Q# 203/ IB Chem/2018/s/TZ 1/Paper 2/Higher Level/Q1/ www.SmashingScience.org:o)

- Urea, (H₂N)₂CO, is excreted by mammals and can be used as a fertilizer.
 - (b) The structural formula of urea is shown.



[2]

(j) The mass spectrum of urea is shown below.



[Source: NIST Mass Spec Data Center, S.E. Stein, director, "Mass Spectra" in NIST Chemistry WebBook, NIST Standard Reference Database Number 69, Eds. P.J. Linstrom and W.G. Mallard, National Institute of Standards and Technology, Gaithersburg MD, 20899, doi:10.18434/T4D303, (retrieved May 31, 2018).]

Identify the species responsible for the peaks at m/z = 60 and 44.


				-/
	¥			

[2]



60:

(k) The IR spectrum of urea is shown below.



[Source: SDBS, National Institute of Advanced Industrial Science and Technology]

Identify the bonds causing the absorptions at 3450 cm⁻¹ and 1700 cm⁻¹ using section 26 of the data booklet.

	[2

3450 cm ⁻¹ :			

1700 cm ⁻¹ :			
	(0	 	



(1)	(i)	Predict the number of signals in the 'H NMR spectrum of urea.	3
	(ii)	Predict the splitting pattern of the ¹ H NMR spectrum of urea.	
			-
	(iii)	Outline why TMS (tetramethylsilane) may be added to the sample to carry out ¹ H NMR spectroscopy and why it is particularly suited to this role.	
Se e			i.
	u orași a		îs
d) 1 (1)			8

Topic Chem 11 Q# 204/ IB Chem/2016/w/TZ0/Paper 2 Section A/Higher Level/


- Ethane-1,2-diol, HOCH₂CH₂OH, has a wide variety of uses including the removal of ice from aircraft and heat transfer in a solar cell.
 - (f) Predict the ¹H NMR data for ethanedioic acid and ethane-1,2-diol by completing the table.
 [2]

	Number of signals	Splitting pattern
Ethanedioic acid:		
Ethane-1,2-diol:		Not required

Topic Chem 11 Q# 205/ IB Chem/2016/s/TZoSP/Paper 2 Section A/Higher Level/Q7/ www.SmashingScience.org: 0) (b) Bute is a chemical with the following molecular formula: C19H20N2O2



(iv) The infrared (IR) spectrum of bute is shown below.



[Source: SDBS web: www.sdbs.riodb.aist.go.jp (National Institute of Advanced Industrial Science and Technology, 2014)]

Using information from section 26 of the data booklet, identify the bonds corresponding to A and B. [1]

A :	
B:	

(v) Based on analysis of the IR spectrum, predict, with an explanation, one bond containing oxygen and one bond containing nitrogen that could not be present in the structure.

Bond containing oxygen not present in structure:


Bond containing nitrogen not present in structure:

Explanation:



[2]

(c) An alcohol, X, of molecular formula C₃H₈O, used as a disinfectant in hospitals, has the following ¹HNMR spectrum:



[Source: SDBS web: www.sdbs.riodb.aist.go.jp (National Institute of Advanced Industrial Science and Technology, 2014)]


The three peaks in the ¹HNMR spectrum of X have chemical shift values centred at $\delta = 4.0, 2.3$ and 1.2 ppm.

- (i) From the integration trace, estimate the ratio of hydrogen atoms in different chemical environments. [1]
- (ii) Deduce the full structural formula of X. [1]

(iii)	Y is an isomer of X containing a different functional group. State the condensed structural formula of Y.	[1]
(iv)	Compare and contrast the expected mass spectra of X and Y using section 28 of the data booklet.	[2]
	One similarity:	
	One difference:	
Горіс	Chem 11 Q# 206/ IB Chem/2016/s/TZ1/Paper 2 Section A/Higher Level/Q2/ www.Smashir (c) The other monomer used in the production of polyurethane is compound Z sbelow.	
	(ii) Deduce the number of signals you would expect to find in the ¹ H NMR of compound Z, giving your reasons.	spectrum [1]


The mass spectrum and infrared (IR) spectrum of compound **Z** are shown below:

Mass spectrum





Compound A and compound B are hydrocarbons.



(b)	Outline how you could use the IR spectra of compounds A and B and section 26 of the
	data booklet to identify them.

(c) Two signals occur in the ¹H NMR spectrum of compound A. Deduce their expected chemical shift and their splitting pattern, using section 27 of the data booklet. [2]

Signal	1	2
Chemical shift / ppm	.,	
Splitting pattern	***********	



[1]

Topic Chem 11 Q# 208/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/

- A student used the technique of titration to determine the concentration of ascorbic acid
 (C₆H₈O₆) in a sample of orange juice. Excess potassium iodide, KI(aq), was added to
 acidified orange juice. The resulting solution was titrated with potassium iodate, KIO₃(aq),
 in the presence of starch as an indicator. The end-point of the titration was shown by a
 blue-black colour.
- (c) The concentration of KIO₃ used in the titration was 2.00 × 10⁻³ mol dm⁻³. The titration produced the following results.

	Titration 1	Titration 2	Titration 3
Final volume of KIO ₃ (± 0.05 cm ³)	7.10	14.40	21.60
Initial volume of KIO ₃ (± 0.05 cm ³)	0.00	7.10	14.40
Volume added of KIO ₃ (± 0.10 cm ³)	7.10	7.30	7.20
Mean volume added of KIO ₃ (± 0.10 cm ³)		7.20	

(i)	Calculate the percentage uncertainty associated with the mean volume of KIO ₃ (aq).	[1
	A ST A M	
(ii)	The colour of orange juice interfered with the blue-black colour at the equivalence point. State the name of this type of error and suggest how this can be minimized.	[2
(ii)	point. State the name of this type of error and suggest how this can be minimized.	[2
(ii)	point. State the name of this type of error and suggest how this can be	[2
(ii)	point. State the name of this type of error and suggest how this can be minimized.	[2



estion 6 continued) In the operation of a mass spectrometer, the first stage is vaporization and the last is (e) detection. State the names of the other three stages and outline what happens in each one. [3] Topic Chem X 93 Q# 210/ IB CHEM/2015/s/TZ2/Paper 2 Section B/Higher Level/ Consider the structure and bonding in MgCl, and PCl₃. (i) State and explain the electrical conductivities of these two chloride compounds in their liquid state. [3] (ii) Suggest, giving your reasons, the approximate pH values of the solutions formed by adding each chloride compound separately to distilled water. [4] MgCl,: PCI,:

Topic Chem X 92 Q# 209/ IB CHEM/2015/w/TZ0/Paper 2 Section B/Higher Level/Q6/ www.SmashingScience.org:o)



Patrick Brannac

Topic Chem 99x99 Q# 211/ IB Chem/2015/w/TZ0/Paper 2 Section A/Higher Level/

(a) The monomers hexanedioic acid and 1,6-diaminohexane react together to form a

	synthetic polymer.	
	Deduce the structural formula of each monomer.	[2
(b)	State the type of polymerization reaction that occurs between these two monomers and identify the structural feature needed in the monomers.	[2
	Type:	
	Structural feature:	
(c)	Draw the structure of the linkage formed in this polymer, and identify the other product of this polymerization reaction.	[2]
	VAR 1/2	



Topic Chem x 910 Q# 212/ IB CHEM/2015/w/TZ0/Paper 2 Section B/Higher Level/Q9/ www.SmashingScience.org

(c) The reaction between the primary halogenoalkane, obtained in part (a) (v), and hot, concentrated alcoholic NaOH is an example of an elimination reaction.

(i) Explain the mechanism of the elimination reaction using curly arrows to represent the movement of electron pairs.

[4]

Topic Chem x 910 Q# 213/ IB CHEM/2015/s/TZ1/Paper 2 Section B/Higher Level/Q7/ www.SmashingScience.org

(e) Describe, using an equation, the elimination of HBr from 2-bromobutane, stating the reagent used.

[2]



Mark Scheme

Topic Chem 1 Q# 1/ IB Chem/2019/w/TZ0/P2/Higher Level/Q8./

			1	1	T.
7.	a	i	$C(NH_2)_3NO_3(s) \rightarrow 2N_2(g) + 3H_2O(g) + C(s)$		1
7.	a	ii	moles of gas = «5 × 10.0 g / 122.11g mol ⁻¹ =» 0.409 «mol» ✓		1
7.	а	iii	$\alpha p = \frac{0.409 \text{mol} \times 8.31 \text{J K}^{-1} \text{mol}^{-1} \times \left(127 + 273\right) \text{K}}{10.0 \text{dm}^3}$ » = 136 αkPa» ✓		1
7.	а	iv	Any two of: nitrogen non-polar/London/dispersion forces AND water polar/H-bonding ✓ water has «much» stronger intermolecular forces ✓ water molecules attract/condense/occupy smaller volume «and therefore deviate from ideal behaviour» ✓		2 max
7.	b		2Na (s) + 2H ₂ O (l) → 2NaOH (aq) + H ₂ (g) ✓ hydrogen explosive OR highly exothermic reaction OR sodium reacts violently with water OR forms strong alkali ✓	Accept the equation of combustion of hydrogen. Do not accept just "sodium is reactive/dangerous".	2

Topic Chem 1 Q# 2/ IB Chem/2019/w/TZ0/P2/Higher Level/Q7./

		The state of the s		48
6.	b	mass of H ₂ O = «18.360 g − 17.917 g =» 0.443 «g» <i>AND</i> mass of CuCl ₂ = «17.917 g − 16.221 g =» 1.696 «g» ✓	Award [3] for correct final answer.	
		moles of $H_2O = \frac{0.443 \text{ g}}{18.02 \text{ g mol}^{-1}} = 0.0246 \text{ (mol)}$		
		OR		3
		moles of CuCl₂ = « 1.696 g = » 0.0126 «mol» ✓	1	
		«water: copper(II) chloride = 1.95: 1»		
		«x =» 2 ✓	Accept «x =» 1.95.	

Topic Chem 1 Q#3/ IB Chem/2018/w/TZ 0/P2/Higher Level/

Topic Chem 1 Q#4/ IB Chem/2018/w/TZ 0/P2/Higher Level/

2.	a	«in 100 g sample» 62.02 g AND 10.43 g 1.01g mol ⁻¹	
		OR	
		«in 100 g sample» 5.164 mol C <i>AND</i> 10.33 mol H ✓	
		27.55 %	3
		OR	
		1.722 mol O √	
		«empirical formula» C₃H ₆ O √	



Topic Chem 1 Q# 5/ IB Chem/2018/w/TZ 0/P2/Higher Level/

1.	a	i	$n_{CuSO4} = 0.0800 \text{ dm}^3 \times 0.200 \text{ mol dm}^{-3} = 0.0160 \text{ mol } AND$ $n_{Fe} = \frac{3.26 \text{ g}}{55.85 \text{ g mol}^{-1}} = 0.0584 \text{ mol } \checkmark$ $CuSO_4 \text{ is the limiting reactant } \checkmark$	Do not award M2 if mole calculation is not shown.	2
1.	a	H	ALTERNATIVE 1: «0.0160 mol × 63.55 g mol ⁻¹ = » 1.02 «g» ✓ «0.872g/1.02g × 100 = » 85.5 «%» ✓ ALTERNATIVE 2: «0.872g/63.55 gmol ⁻¹ = » 0.0137 «mol» ✓ «0.0137 mol/0.0160	Accept answers in the range 85–86 %. Award [2] for correct final answer.	2

Topic Chem 1 Q# 6/ IB Chem/2018/s/TZ 2/P2/Higher Level/

			1	1
1.	a	$n(H_2SO_4) \approx 0.0500 \text{ dm}^3 \times 0.100 \text{ mol dm}^{-3} = 0.00500/5.00 \times 10^{-3} \approx \text{mol} \Rightarrow \checkmark$		1
1.	b	$H_2SO_4(aq) + Mg(OH)_2(s) \rightarrow MgSO_4(aq) + 2H_2O(l) \checkmark$	Accept an ionic equation.	1
1.	С			1
1.	d	n(H₂SO ₄) reacted «= 0.00500 − 0.001178» = 0.00382/3.82 × 10 ⁻³ «mol» ✓		1
1.	e	$n(Mg(OH)_2) \ll = n(H_2SO_4) = w = 0.00382/3.82 \times 10^{-3} \text{ wmolw } \checkmark$ $m(Mg(OH)_2) \ll = 0.00382 \text{ mol} \times 58.33 \text{ g mol}^{-1} w = 0.223 \text{ wgw } \checkmark$	Award [2] for correct final answer.	2
1.	f	% Mg(OH) ₂ «= $\frac{0.223 \text{ g}}{1.24 \text{ g}}$ × 100» = 18.0 «%» √	Answer must show three significant figures.	1

Topic Chem 1 Q#7/ IB Chem/2018/s/TZ 1/P2/Higher Level/

1.	а	ı	molar mass of urea $\ll 4 \times 1.01 + 2 \times 14.01 + 12.01 + 16.00$ $\approx 60.07 \text{ (g mol}^{-1}) \checkmark$ $\ll \% \text{ nitrogen} = \frac{2 \times 14.01}{60.07} \times 100 = \% 46.65 \text{ (%%)} \checkmark$	Award [2] for correct final answer. Award [1 max] for final answer not to two decimal places.	2
1.	a	ii	«cost» increases AND lower N % «means higher cost of transportation per unit of nitrogen» OR «cost» increases AND inefficient/too much/about half mass not nitrogen ✓	Accept other reasonable explanations. Do not accept answers referring to safety/explosions.	1
1.	С		$n(\text{KNCO}) \approx 0.0500 \text{ dm}^3 \times 0.100 \text{ mol dm}^{-3} = 5.00 \times 10^{-3} \text{ emol} \text{ s} \checkmark$ $\text{emass of urea} = 5.00 \times 10^{-3} \text{ mol} \times 60.07 \text{ g mol}^{-1} = 0.300 \text{ egs} \checkmark$	Award [2] for correct final answer.	2
1.	d	iii	In $K = \frac{-\Delta G^{\circ}}{RT} = \frac{-50 \times 10^{3} \text{ J}}{8.31 \text{ J K}^{-1} \text{ mol}^{-1} \times 298 \text{ K}} = -20 \text{ J}$	Accept range of 20-20.2 for M1. Award [2] for correct final answer.	2



1.	f		$2(H_2N)_2CO\left(s\right) + 3O_2(g) \rightarrow 4H_2O\left(l\right) + 2CO_2\left(g\right) + 2N_2\left(g\right)$ correct coefficients on LHS \checkmark correct coefficients on RHS \checkmark	Accept $(H_2N)_2CO(s) + \frac{3}{2}O_2(g) \rightarrow$ $2H_2O(l) + CO_2(g) + N_2(g)$. Accept any correct ratio.	2
in a	g		$ \text{«V} = \frac{0.600 \text{g}}{60.07 \text{g mol}^{-1}} \times 22700 \text{cm}^3 \text{mol}^{-1} = \text{» } 227 \text{«cm}^3 \text{»} \checkmark $		1
pi	ic Ch	em 1	. Q#8/ IB Chem/2017/s/TZ 2/P2/Higher Level/		
	а	iii	« [K ₂ Cr ₂ O ₇] = $\frac{13.239 \text{ g}}{294.20 \text{ g mol}^{-1} \times 0.100 \text{ dm}^{-3}}$ =» 0.450 «mol dm ⁻³ » ✓		1
	a	iv	$n(Sn^{2+}) = \text{$0.450 mol dm}^{-3} \times 0.01324 dm^{3} \times \frac{3 mol}{1 mol} = \text{$0.0179 $mol $} \checkmark$ $\text{$(Sn^{2+})$} = \frac{0.0179 mol}{0.0100 dm^{3}} = \text{$1.79 $mol dm}^{-3} \text{$$$} \checkmark$	Award [2] for correct final answer.	2
			. Q# 9/ IB Chem/2017/s/TZ 2/P2/Higher Level/		1
pi	ic Ch	em 1			1

Topic Chem 1 Q# 10/ IB Chem/2017/s/TZ 2/P2/Higher Level/	Topic Chem		IB Chem/2017/s/TZ 2/P2/Higher Level
--	------------	--	-------------------------------------

oxide not decomposed completely v

heat sample to constant mass «for three or more trials» ▼

≈ 1 / ratio of Ag to O approximately 1 : 1, so»

0.03036

0.03031 AgO ✓

temperature too low

heating time too short

1

ii

a

1	b	A _r closer to 107/less than 108 «so more ¹⁰⁷ Ag» OR	Accept calculation that gives greater than 50% 107Ag.	1
	8 98	A _r less than the average of (107 + 109) «so more ¹⁰⁷ Ag» ✓	9458	228



2

Accept "not heated strongly enough".

If M1 as per markscheme, M2 can only be scored for constant mass technique.

Accept "soot deposition" (M1) and any

suitable way to reduce it (M2).

dessicator" (M2).

Accept "absorbs moisture «from atmosphere» (M1) and "cool in

Award [1 max] for reference to "impurity" AND design improvement.

1 c	i	Flask	Colour of solution	Product formula	Do not accept name for the products. Accept "Na $^+$ + OH $^-$ " for NaOH.	
		Na ₂ O	blue AND	NaOH ✓	Ignore coefficients in front of formula.	3
		P4O 10	yellow ✓	H₃PO ₄ ✓		
1 c	ii	«molten» P ₄ O ₁₀		d particles AND conducts el ns/charged particles AND d electricity ✓	mobile charges being present	2
d		energy levels or	crete/specific/certain/diffe onverge/get closer togeth onverge with distance fro		Accept appropriate diagram for either M1, M2 or both. Do not give marks for answers that refe to the lines in the spectrum.	2
pic Ch	em 1	Q# 11/ IB Che	em/2017/s/TZ 1/P2/	/Higher Level/		,
6. с	i	12.01	$021 \text{ AND H} = \frac{4.08}{1.01} = 4.0$	04 AND CI = $\frac{71.65}{35.45}$ = 2.02	Accept 24.27 : 4.08 : 71.65 12.01 : 35.45	2
20.000	em 1	total mass of oxy	em/2017/s/TZ 1/P2/ rgen «= 8.0 × 10 ⁻³ g dm ⁻¹	/Higher Level/ 3 × 1000 dm3» = 8.0 «g»	Do not accept C ₂ H ₄ Cl ₂ . Award [2] for correct final answer. Award [3] for correct final answer.	
Commence of the Control of the Contr		Q# 12/ IB Che total mass of oxy $n(O_2) = \frac{8.0}{32.00 \text{ g}}$ OR $n(N_2H_4) = n(O_2)$	em/2017/s/TZ 1/P2/ gen «= 8.0 × 10 ⁻³ g dm ⁻¹ gmol ⁻¹ = » 0.25 «mol»	³ × 1000 dm ³ » – 8.0 *g*	Do not accept C ₂ H ₄ Cl ₂ . Award [2] for correct final answer.	3
Comment of the State of Comment		Q# 12/ IB Che total mass of oxy $n(O_2) = \frac{8.0}{32.00 \text{ g}}$ OR $n(N_2H_4) = n(O_2) = \frac{8.0}{32.00 \text{ g}}$ «mass of hydraz	em/2017/s/TZ 1/P2/ rgen «= 8.0 × 10 ⁻³ g dm ⁻¹ gmol ⁻¹ = » 0.25 «mol» ine = 0.25 mol × 32.06 g 32.00 gmol ⁻¹ = » 0.25	3 × 1000 dm³» = 8.0 «g» mol⁻¹ =» 8.0 «g» ✓ 5 «mol»	Do not accept C ₂ H ₄ Cl ₂ . Award [2] for correct final answer.	
. h	i	Q# 12/ IB Che total mass of oxy $n(O_2) = \frac{8.0}{32.00 \text{ g}}$ OR $n(N_2H_4) = n(O_2) = \text{mass of hydraz}$ $\text{mn}(N_2H_4) = n(O_2) = \text{mn}(N_2H_4)$ $\text{modern}(N_2H_4) = n(O_2) = \text{mn}(N_2H_4)$	em/2017/s/TZ 1/P2/ gen «= 8.0 × 10 ⁻³ g dm ⁻¹ gmor ⁻¹ = » 0.25 «mol» gmor ⁻¹ = 0.25 mol × 32.06 g	3 × 1000 dm ³ » = 8.0 «g» mol ⁻¹ =» 8.0 «g» ✓ 5 «mol» m ³ mol ⁻¹ » = 6.2 «dm ³ » ✓	Do not accept C ₂ H ₄ Cl ₂ . Award [2] for correct final answer. Award [3] for correct final answer.	3
5. h	i	Q# 12/ IB Che total mass of oxy $n(O_2) = \frac{8.0}{32.00 \text{ g}}$ OR $n(N_2H_4) = n(O_2) = \frac{8.0}{32.00 \text{ g}}$ "mass of hydraz $\text{"n(N_2H_4)} = n(O_2) = \frac{8.0}{32.00 \text{ g}}$ "volume of nitro $Q# 13/ IB Che$	em/2017/s/TZ 1/P2/gen «= 8.0×10^{-3} g dm ⁻¹ /gmol ⁻¹ = $^{\circ}$ 0.25 «mol» gine = 0.25 mol × 32.06 g $^{\circ}$ = $\frac{8.0g}{32.00g \text{mol}^{-1}}$ = $^{\circ}$ 0.25 gen = 0.25 mol × 24.8 dr em/2017/s/TZ 1/P2/giv > TiO ₂ (s) + 4HCl (according)	3 × 1000 dm ³ » = 8.0 «g» mol ⁻¹ =» 8.0 «g» ✓ 5 «mol» m ³ mol ⁻¹ » = 6.2 «dm ³ » ✓ /Higher Level/	Do not accept C ₂ H ₄ Cl ₂ . Award [2] for correct final answer. Award [3] for correct final answer.	3

2.	a Weak acid: partially dissociated/ionized «in solution/water» AND Strong acid: «assumed to be almost» completely/100% dissociated/ ionized «in solution/water» ✓	Accept answers relating to pH, conductivity, reactivity if solutions of equal concentrations stated.	1
----	---	--	---



Question		estion Answers		Notes	Total
2.	b		«log scale» reduces a wide range of numbers to a small range OR simple/easy to use OR converts exponential expressions into a linear scale/simple numbers ✓	Do not accept "easy for calculations".	1
2.	С	i	phenolphthalein OR phenol red ✓		1
),	С	ii			1
2.	С	iii	« 1/2 × 1.40 × 10 ⁻³ =» 7.00 × 10 ⁻⁴ «mol» ✔		1

C	uest	ion	Answers	Notes	Total
2.	c	iv	Answers ALTERNATIVE 1: «mass of pure hydrated ethanedioic acid in each titration = $7.00 \times 10^{-4} \text{mol} \times 126.08 \text{g mol}^{-1} = \text{w} 0.0883 / 8.83 \times 10^{-2} \text{«g»} \checkmark$ mass of sample in each titration = $\frac{25}{1000} \times 5.00 \text{g} = \text{w} 0.125 \text{«g»} \checkmark$ «% purity = $\frac{0.0883 \text{g}}{0.125 \text{g}} \times 100 = \text{w} 70.6 \text{«%»} \checkmark$ ALTERNATIVE 2: «mol of pure hydrated ethanedioic acid in 1 dm³ solution = $7.00 \times 10^{-4} \times \frac{1000}{25} = \text{w} 2.80 \times 10^{-2} \text{«mol»} \checkmark$ «mass of pure hydrated ethanedioic acid in sample = $2.80 \times 10^{-2} \text{mol} \times 126.08 \text{g mol}^{-1} = \text{w} 3.53 \text{«g»} \checkmark$ «% purity = $\frac{3.53 \text{g}}{5.00 \text{g}} \times 100 = \text{w} 70.6 \text{«%»} \checkmark$	Notes Award suitable part marks for alternative methods. Award [3] for correct final answer. Award [2 max] for 50.4% if anhydrous ethanedioic acid assumed.	Total
			5.00 g ALTERNATIVE 3: mol of hydrated ethanedioic acid (assuming sample to be pure) = $\frac{5.00 \mathrm{g}}{126.08 \mathrm{gmol}^{-1}} = 0.03966 \mathrm{«mol} \mathrm{»} \checkmark$ actual amount of hydrated ethanedioic acid = $ (7.00 \times 10^{-4} \times \frac{1000}{25}) = 2.80 \times 10^{-2} \mathrm{mol} \mathrm{»} \checkmark$ $ (\% \text{ purity}) = \frac{2.80 \times 10^{-2}}{0.03966} \times 100 = 70.6 \text{«%»} \checkmark$		

Topic Chem 1 Q# 15/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

b i	$n_{\rm c}: \left\langle \frac{73.99}{12.01} \right\rangle = 6.161 ({ m mol}) \text{ and } n_{\rm H}: \left\langle \frac{6.55}{1.01} \right\rangle = 6.49 ({ m mol}) \text{ and } n_{\rm N}: \left\langle \frac{9.09}{14.01} \right\rangle = 0.649 ({ m mol}) \text{ and } n_{\rm C}: \left\langle \frac{10.37}{16.00} \right\rangle = 0.6481 ({ m mol}) \checkmark$	Award [2 max] for correct final answer without working	3
	$n_{\rm c}:n_{\rm H}:n_{\rm o}:n_{\rm o}=9.5:10:1:1~\checkmark$	1/2 1/9	
	Empirical formula: C ₁₉ H ₂₀ N ₃ O ₂ ✓	1 11 10	

Topic Chem 1 Q# 16/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

(Question		Answers	Notes	Total
1.	a	i	<(22.05+22.15)(0.5) →22.10 (cm³) ✓		1
	а	ii	$\left\langle \frac{22.10 \times 0.100}{1000} \right\rangle = 2.21 \times 10^{-3} / 0.00221 < \text{mol} > \checkmark$		1
	а	iii	$\left\langle \frac{0.5 \times 2.21 \times 10^{-3} \times 1000}{25.00} \right. = \left. \right\rangle 4.42 \times 10^{-2} / 0.0442 \langle \text{mol dm}^{-3} \rangle \checkmark$		1
	а	iv	(4.42×10 ⁻² ×10 => 4.42×10 ⁻¹ /0.442 (mol dm ⁻³) ✓		1

Topic Chem 1 Q# 17/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

1.



1.	С	i	€\(\left(\frac{2.478}{4×30.97}\right) > - 0.02000 €mols \(\not\)		1
	С	ii	n(NaOH) = €0.1000 × 5.00 => 0.500 €mols AND P ₄ /phosphorus is limiting reagent ✓	Accept $n(H_2O) = \frac{100}{18} = 5.50$ AND P_4 is limiting reagent.	1
	С	iii	amount in excess ← 0.500 – (3 × 0.02000)» = 0.440 ←mol» ✓		1
	С	iv		Accept methods employing pV = nRT, with p as either 100 (454 cm²) or 101.3kPa (448 cm²). Do not accept answers in dm².	1

(Question		Answers	Notes	Total
1.	d		$n(P) \leftarrow \frac{43.6}{30.97} \Rightarrow -1.41 \text{ emols} \checkmark$ $n(O) \leftarrow \frac{100 - 43.6}{16.00} \Rightarrow -3.53 \text{ emols} \checkmark$ $\leftarrow \frac{n(O)}{n(P)} = \frac{3.53}{1.41} - 2.50 \text{ so empirical formula is} P_2O_s \checkmark$	Accept other methods where the working is shown.	3
1.	d	iii	€ 285 141.9 - 2.00 , so molecular formula - 2 × P ₂ O ₈ -> P ₄ O ₁₀ ✓		1

Topic Chem 1 Q# 18/ IB Chem/2015/w/TZ0/P2 Section A/Higher Level/

1. (4) (1) 10, 101, 1710 100	1.	(a)	(i)	IO_3 to I_2 : V/+5 to 0
------------------------------	----	-----	-----	-----------------------------

$$I^-$$
 to $I_2:-1/-1$ to 0;

[2]

Accept change in oxidation number -5 and +1. Penalize incorrect notation such as 5+ or 5 once only.

(ii) Oxidizing agent: IO_/iodate and Reducing agent. I_/iodide; [1]

(b) $0.250 \times 2.00 \times 10^{-3} / 5.00 \times 10^{-4}$ (mol of KIO₃); $(5.00 \times 10^{-4} \times 214.00 =) 0.107(g);$ Award [2] for the correct final answer.

[2]

(C) 1.4 (%); Accept 1 (%). [1]

systematic;

dilute the orange juice;

[2]

Accept other valid suggestions, eg. compare with a standard (showing colour at equivalence) / look at mixture through a yellow filter / add more starch (for a sharper colour change) / filter orange juice (through charcoal). Do not accept repeat titrations or alternative indicator.

1.44×10⁻⁵ (mol); (iii)

[1]

IO₃⁻: 3C₆H₈O₆ / 1:3 mole ratio; (d)

$$(1.44 \times 10^{-5} \text{ mol} \times 3 =) 4.32 \times 10^{-5} \text{ (mol)};$$

[2]

Award [2] for the correct final answer.

Award [1 max] for "4.80 x 10-6 (mol)" obtained from reversed ratio, 3:1.

(e)
$$(4.32 \times 10^{-5} \times 176.14 =) 7.61 \times 10^{-3} (g)$$
;
Accept $M_r = 176$ and mass = $7.60 \times 10^{-3} (g)$.

[1]



(f) gradient =
$$-\frac{E_a}{R}/-1233$$
 (K);

Accept value from -1200 to -1260 (K).

$$E_a = (1233 \times 8.31 =) 1.02 \times 10^4 \text{ Jmol}^{-1} / 10.2 \text{kJ mol}^{-1}$$

correct E_a value;

correct units;

Allow value in range 9.97 - 10.5 kJ mol-1.

Answer must be given to three significant figures.

Award [3] for correct final answer and units.

Accept J or kJ instead of J mol-1 or kJ mol-1.

Topic Chem 1 Q# 19/ IB Chem/2015/s/TZ1/P2 Section A/Higher Level/

1. (a)
$$\left(\frac{(12.70 + 12.50)}{2}\right) = 12.60 \text{ (cm}^3)$$
;

 $(0.01260 \times 0.100 =) 1.26 \times 10^{-3} \text{ (mol)};$

[2]

[3]

Award [2] for correct final answer.

(b) (i)
$$\left(\frac{1.26 \times 10^{-3}}{2}\right) = 6.30 \times 10^{-4} \text{(mol)};$$

[1]

(ii)
$$(6.30 \times 10^{-4} \times 10 =) 6.30 \times 10^{-3} \text{ (mol)};$$

[1]

(iii)
$$\left(\frac{0.795}{6.30\times10^{-3}}\right) = 126 (gmol^{-1});$$


[1]

(iv)
$$M_r(C_2H_2O_4) = 90.04$$
 and $M_r(H_2O) = 18.02$;
 $x = 2$

[2]

Accept integer values for M_r's of 90 and 18 and any reasonable calculation. Award [1 max] if no working shown.

Topic Chem 2 Q# 20/ IB Chem/2018/w/TZ 0/P2/Higher Level/





Topic Chem 2 Q# 21/ IB Chem/2017/s/TZ 2/P2/Higher Level/

4	C	Any two of:		
		IE4: electron in lower/inner shell/energy level		
		OR		
		IE₄: more stable/full electron shell ✓		
		IE ₄ : electron closer to nucleus	Accept "increase in effective nuclear	2 max
		OR	charge" for M2.	
		IE₄: electron more tightly held by nucleus ✓		
		IE₄: less shielding by complete inner shells ✓		

Topic Chem 2 Q# 22/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

Question		ion	Answers	Notes	Total
4.	a		26 12 Mg ✓		1
4.	b		$\alpha A_r = \frac{24 \times 78.60 + 25 \times 10.11 + 26 \times 11.29}{100}$ $\alpha = 24.3269 = \frac{24.33}{2}$	Award [2] for correct final answer. Do not accept data booklet value (24.31).	2
4.	С		contamination with sodium/other «compounds» ✓		1
4.	d	i	energy levels are closer together at high energy / high frequency / short wavelength ✓		1
4.	d	ii	ionisation energy ✓		1

Topic Chem 2 Q# 23/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

d	i	[Ne]3s³3p⁴ ✓	Electrons must be given as superscript.	1
d	ii	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	100	1

Topic Chem 2 Q# 24/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

e	ionic radius of nitrogen is 146pm/146×10 ⁻¹² m which is greater than	Values must be given to score mark.	
	atomic radius which is 71pm/71×10 ⁻¹² m due to increased repulsion	117/	1
	between electrons ✓		

Topic Chem 2 Q# 25/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

4.	e	ì	First 4s AND Second: 4s AND Third: 3d AND Fourth: 3d ✓	Do not apply ECF from (c).	1
4.	e	ii	(in the same sub-shell and a) decrease in electron-electron repulsion OR (in the same sub-shell and) as more electrons removed, the pull of of the nucleus/positive ions holds the remaining electrons more tightly ✓	Do not accept "greater nuclear charge/ effective nuclear charge".	1
4.	e	iii	electron 5 is lost from the 3d orbital OR electron 5 is lost from the valence shell ✓ electron 6 is lost from a 3p orbital OR electron 6 is lost from a ∢complete inner shell ✓ 3p orbital/complete inner shell experiences a much larger effective nuclear charge OR 3p orbital/complete inner shell is less well shielded OR 3p orbital/complete inner shell is nearer the nucleus ✓	Award [1 max] (for M1/M2) (ECF) if candidate recognises electrons 5 and 6 are from different levels.	3
4.	e	iv	28 ✓		1

Topic Chem 2 Q# 26/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

79.91 = 79x + 81(1-x);

Award M1 for any suitable calculation.

(abundance 79Br =) 54.5 %;

Award [2] for correct final answer.

Topic Chem 3 Q# 27/ IB Chem/2019/w/TZ0/P2/Higher Level/Q7./

6.	а	i	[Ar] 3d ¹⁰	
			OR	1
			1s² 2s² 2p6 3s² 3p6 3d10 ✓	

[2]

6.	f	i	«ligands cause» d-orbitals «to» split ✓ light absorbed as electrons transit to higher energy level «in d–d transitions» OR light absorbed as electrons promoted ✓ energy gap corresponds to «orange» light in visible region of spectrum ✓ colour observed is complementary ✓		3 max
6.	f	ii	full «3»d sub-level/orbitals OR no d–d transition possible «and therefore no colour» ✓		1
opi	Che	em 3	Q# 28/ IB Chem/2018/w/TZ 0/P2/Higher Level/		- k
4.	a		nuclear charge/number of protons/Z _{et} increases «causing a stronger pull on the outer electrons» ✓	Accept "atomic number" for "number of protons".	2

4.	a	nuclear charge/number of protons/Z _{et} increases «causing a stronger pull on the outer electrons» ✓ same number of shells/«outer» energy level/shielding ✓	Accept "atomic number" for "number of protons".	2
4.	b	isoelectronic/same electronic configuration/«both» have 2.8 ✓ more protons in Na ⁺ ✓		2

Topic Chem 3 Q# 29/ IB Chem/2018/w/TZ 0/P2/Higher Level/

3.	a	i	1s ² 2s ² 2p ⁶ 3s ² 3p ⁶ 4s ² 3d ¹⁰ 4p ⁵ OR [Ar] 4s ² 3d ¹⁰ 4p ⁵ ✓	Accept 3d before 4s.	1
3.	a	ii	1 1 445» 1 445»	Accept double-headed arrows.	1
3.	b	1	Structure I – follows octet rule: [:○──Br→○:] / [:○──Br──○:] / [:○──Br──○:] / Structure II – does not follow octet rule: [:○──Br──○] / [:○──Br──○] / (:○○──Br──○) / (:○○──Br──○) / (:○○──Br──○) / (:○○──Br──○) / (:○○──Br──○) / (:○○───────────────────────────────────	Accept dots, crosses or lines to represent electron pairs.	2
3.	b	II	«structure I» formal charge on Br = +2 OR «structure II» formal charge on Br = 0/+1 structure II is preferred AND it produces formal charge closer to 0 ✓	Ignore any reference to formal charge on oxygen.	2



		1			
3.	C		Geometry:		
			trigonal/pyramidal ✔		
			Reason:		
			three bonds AND one lone pair		
			OR		3
			four electron domains ✓	Accept "charge centres" for "electron domains".	
			O-Br-O angle:	Accept answers in the range 104-109°.	
			107° ✓	and the first section of the section	
3.	d	i	$BrO_3^-(aq) + 6e^- + 6H^+(aq) \rightarrow Br^-(aq) + 3H_2O(l)$	Accept reversible arrows.	
			correct reactants and products ✓		2
			balanced equation ✓		
3.	d	ii	$BrO_3^-(aq) + 6Fe^{2^+}(aq) + 6H^+(aq) \rightarrow Br^-(aq) + 3H_2O(l) + 6Fe^{3^+}(aq)$		1
3.	d	iii	E ⁰ reaction = «+1.44 ∨ - 0.77 ∨ =» 0.67 «∨» ✓		
			ΔG^{Θ} = «-nFE $^{\Theta}$ _{reaction} = -6 × 96500 C mol ⁻¹ × 0.67 V =» -3.9 × 10 ⁵ «J» \checkmark		2
3.	e		both are paramagnetic ✓	Accept orbital diagrams for both ions showing	Attion
			«both» contain unpaired electrons ✓	unpaired electrons.	2

Topic Chem 3 Q# 30/ IB Chem/2018/s/TZ 1/P2/Higher Level/

2. с	«promoted» electrons fall back to lower energy level ✓ energy difference between levels is different ✓	Accept "Na and Ca have different nuclear charge" for M2.	2
2. e	(i)		2

Topic Chem 3 Q# 31/ IB Chem/2018/s/TZ 1/P2/Higher Level/

1	Sec			7 -
1.	h	lone/non-bonding electron pairs «on nitrogen/oxygen/ligand» given to/shared with	D 40	
		metal ion ✓		2
1		co-ordinate/dative/covalent bonds ✓		
1		co-ordinate/dative/covalent bonds v		

Topic Chem 3 Q# 32/ IB Chem/2017/s/TZ 2/P2/Higher Level/

	100					1	1
1	С	i	Flask containing	Colour of solution	Product formula	Do not accept name for the products. Accept "Na $^+$ + OH $^-$ " for NaOH.	
			Na ₂ O	blue	NaOH ✓	Ignore coefficients in front of formula.	3
			P4O ₁₀	AND yellow ✓	H₃PO4 ✓		
-	+	+	.^-	<u> </u>			4



Topic Chem 3 O# 33/ IB Chem/2017/s/TZ 1/P2/Higher Level/

2.	d	I.E.	A log graph is acceptable. Accept log plot on given axes (without amendment of y-axis). Award mark if gradient of 5 to 6 is greater than "best fit line" of 1 to 5.	1
2.		regular increase for first five AND sharp increase to the 6th ✓	Do not occupify for a fee	1
2.	e	pair of electrons provided by the ligand ✓	Do not accept "dative" or "co- ordinate bonding" alone.	1
2.	f	partially filled d-orbitals <pre></pre>		4 max

	c Che	≥m 3	Q# 34/ IB Chem/2017/s/TZ 1/P2/Higher Level/ (46 × 7.98) + (47 × 7.32) + (48 × 73.99) + (49 × 5.46) + (50 × 5.25) 100 = 47.93 ✓	Answer must have two decimal places with a value from 47.90 to 48.00. Award [2] for correct final answer. Award [0] for 47.87 (data booklet value).	2
2.	С		Protons: 22 AND Neutrons: 26 AND Electrons: 22 ✓		1
2.	d	i	1s ² 2s ² 2p ⁶ 3s ² 3p ⁶ 3d ² ✓		1

Topic Chem 3 Q# 35/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

4.	е	$MgO(s) + H_2O(l) \rightarrow Mg(OH)_2(s)$ OR $MgO(s) + H_2O(l) \rightarrow Mg^{2+}(aq) + 2OH^{-}(aq) \checkmark$	Accept ⇒.	1
4.	f	from basic to acidic ✓ through amphoteric ✓	Accept "alkali/alkaline" for "basic". Accept "oxides of Na and Mg: basic AND oxide of Al: amphoteric" for M1. Accept "oxides of non-metals/Si to Ct acidic" for M2. Do not accept just "become more acidic".	2
4.	g	Mg₃N₂ ✓	Accept MgO ₂ , Mg (OH) ₂ , Mg (NO ₃) ₂ , MgCO ₃	1

Topic Chem 3 Q# 36/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

2.	f	coordinate/dative/covalent bond from O to «transition» metal «ion» OR acts as a Lewis base/nucleophile ✓ can occupy two positions OR provide two electron pairs from different «O» atoms OR form two «coordinate/dative/covalent» bonds «with the metal ion» OR	2
		chelate «metal/ion» ✓	



Topic Chem 3 Q# 37/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

4.	c		1s²2s²2p°3s²3p°3d³4s² OR 1s²2s²2p°3s²3p°4s²3d³ incomplete d ∢sub-» level/orbital/shell ∢in its compounds/ions» ✓		2
4.	d	i	give/donate a lone/non-bonding electron pair ✔	Accept "through the formation of a dative/ coordinate bond". Accept "by acting as Lewis bases". Do not accept "act as ligands".	1

Question		Answers	Notes	Total
4. d	ii	⊄more chlorido ligandss smaller energy gap between split d-orbitals OR Cl⁻ is lower than H₂O in spectrochemical series OR Cl⁻ is a weaker ligand/has lower charge density ✓ the absorption will move to longer wavelengths OR the absorption wavelength will increase ✓	Do not accept answers in terms of change of frequency.	2

Topic Chem 3 Q# 38/ IB CHEM/2015/w/TZ0/P2 Section B/Higher Level/

(e) (i) Na and Mg: basic

Al: amphoteric

Do not accept amphiprotic.

Si to Cl: acidic

Ar: no oxide ;;

Award [2] for three or four correct, award [1] for two correct.

Award [1] for stating oxides become more acidic towards the right/chlorine or more basic towards the left/sodium.

Do not penalize incorrect formulas of oxides.

(ii) $Na_2O(s) + H_2O(l) \rightarrow 2NaOH(aq) / Na_2O(s) + 2HCl(aq) \rightarrow 2NaCl(aq) + H_2O(l);$

Accept a correct equation with any acid or acidic oxide.

 $SO_3(l) + H_2O(l) \rightarrow H_2SO_4(aq) / SO_3(l) + 2NaOH(aq) \rightarrow Na_2SO_4(aq) + H_2O(l);$ [2]

Accept a correct equation with any metal hydroxide, metal oxide, metal carbonate or metal hydrogen carbonate.

Do not accept equation with SO2.

Ignore state symbols.

Accept ionic equations for M1 and/or M2.

Topic Chem 3 Q# 39/ IB Chem/2015/w/TZ0/P2 Section A/Higher Level/

- (a) (i) ability of an atom to attract (a pair of) electrons in a covalent bond/molecule
 / ability of an atom to attract a shared pair of electrons;
 Do not accept element/nucleus instead of atom.
 - (ii) do not form bonds/compounds / do not share electrons / have (full/stable)
 octet / have full/stable outer shell;
 Accept (chemically) inert / do not react / stable electron arrangements/
 configurations.

 [1]
 - (c) Na⁺ has one less energy level/shell compared to Na atom / Na⁺ has 2 energy levels/shells compared to 3 in Na atom;
 (compared to Na atom) effective nuclear charge greater in Na⁺ (so valence electrons attracted more strongly to nucleus) / (compared to Na atom) more protons than electrons in Na⁺ (so valence electrons attracted more strongly to nucleus);

 [1 max]



[1]

[2]

Topic Chem 3 Q# 40/ IB Chem/2015/w/TZ0/P2 Section A/Higher Level/

(a) (Cu) 1s²2s²2p⁶3s²3p⁶4s¹3d¹⁰ / 1s²2s²2p⁶3s²3p⁶3d¹⁰4s¹;
 Do not accept [Ar] 4s¹3d¹⁰.

[2]

Do not accept [Ar] 3d9.

(c) d orbitals are split (into two sets of different energies); (frequency/wavelength of visible) light absorbed by electrons moving between d levels/orbitals:

colour due to remaining frequencies/complementary colour transmitted;

Cu²⁺ has unpaired electrons/partially filled d sub-level / Zn²⁺ has filled d sub-level;

Accept d-d transitions possible between orbitals for M2.

[4]

Topic Chem 3 Q# 41/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(b) (i) Na, Mg (oxides): basic

Al (oxide): amphoteric
Do not accept amphiprotic.

Si to Cl (oxides): acidic

[2]

Award [2] for all three listed sets correct.

Award [1] for one or two listed sets correct.

Award [1] for stating oxides become more acidic towards right/Cl or more basic towards left/Na.

Do not penalize if reference is to Ar instead of Cl.

Do not penalize for incorrect formulas of oxides.

(ii) Na₂O(s)+H₂O(l) \rightarrow 2NaOH(aq);

$$P_4O_{10}(s) + 6H_2O(l) \rightarrow 4H_3PO_4(aq)$$
;

[2]

Ignore state symbols.

Accept $P_2O_5(s) + 3H_2O(l) \rightarrow 2H_3PO_4(ag)$.

Do not award marks if incorrect formulas of the oxides are used.

Topic Chem 3 Q# 42/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(iv)

[2]

Accept any other octahedral arrangement of ligands.

Ignore missing square brackets and charge.



Topic Chem 3 Q# 43/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/ 8. (a) (i) [1] 45 Accept full-arrows. Accept boxes in reverse order or at different energy levels. Do not award the mark if sub-levels are incorrectly labelled or if no boxes are drawn. (electrostatic) attraction between (lattice of) cations/positive/Cr3+ ions and delocalized electrons; [1] (delocalized electrons allows) the layers/rows of cations/positive/Cr3+ ions (iii) to slide past each other (without disrupting the metallic bonding); [1] Accept atoms instead of ions. III / +3: (C) (i) [1] Do not eccept incorrect notation such as 3+/3. ligand has lone/non-bonding electron pair / (II) dative (covalent)/coordinate/coordination bond forms; ligand is Lewis base / ion is Lewis acid; [2] (iii) partially filled/incomplete d sub levels/orbitals; d orbitals split into two levels; energy difference is in visible part of spectrum / electrons absorb visible light/one colour/frequency/wavelength; electron transitions occur from lower to higher energy level (within d sub-level); complementary colour/colour not absorbed is seen; [4 max] Do not accept complementary colour "emitted". Topic Chem 3 Q# 44/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/ (e) Cl has 7 valence electrons and is in group 7; Accept "group 17" as suggested by IUPAC. Ct has 3 occupied (electron) shells/energy levels and so is in period 3; [2] Topic Chem 3 Q# 45/ IB Chem/2015/s/TZ1/P2 Section A/Higher Level/ 2. $Na_{2}O(s) + H_{2}O(l) \rightarrow 2NaOH(aq);$ Accept Na₂O(s) + H₂O(l) \rightarrow 2Na⁺ (aq) + 2OH -(aq). $P_4O_{40}(s) + 6H_2O(l) \rightarrow 4H_4PO_4(aq)$; [2] Accept $P_2O_5(s) + 3H_2O(1) \rightarrow 2H_3PO_4(aq)$. Accept $P_4O_{10}(s) + 6H_2O(l) \rightarrow 4H^+(aq) + 4H_2PO_4^-(aq)$. Ignore state symbols. Topic Chem 3 Q# 46/ IB Chem/2015/s/TZ1/P2 Section A/Higher Level/ 1s²2s²2p⁶3s²3p⁶3d¹⁰4s¹ / 1s²2s²2p⁶3s²3p⁶4s¹3d¹⁰; (a) [1] (b) 63Cu lower boiling point/melting point/density/greater rate of diffusion than 85Cu; Accept converse argument. Do not accept "different mass".

Chemical:

(properties identical because) same electron configuration/arrangement of electrons:

Accept "same number of protons and electrons".

Do not accept "same number of electrons" OR "same valence (electrons)" OR "same atomic number" only.

Topic Chem 4 Q# 47/ IB Chem/2019/w/TZ0/P2/Higher Level/Q7./

6. f	f	iii	octahedral AND 90° «180° for axial» ✓		Accept square-based bi-pyramid.	1
pic (Cher	m 4	Q# 48/ IB Chem/2019/w/TZ0/P2/Higher Level/Q2./	:->		
1. a	3		Ö — Ö Ö — Ö ✓ Coordinate bond may be represented by an arrow. Do not accept delocalized structure for ozone.		2025 7.255 5.59	2
1. b	o		resonance «structures» OR delocalization of «the double/pi bond» electrons ✓ 121 «pm» < length < 148 «pm» ✓	Accept any lea	ngth between these two values.	2
l. c	:		any value from 110°–119° ✓			1
l. d	i		«bond» in O₂ stronger than in O₃ ✓ ozone absorbs lower frequency/energy «radiation than oxygen» OR ozone absorbs longer wavelength «radiation than oxygen» ✓	Accept ozone	«layer» absorbs a range of frequencies.	2
. е		i	steps 1 AND 3 ✓			1
1.	e	II	ALTERNATIVE 1: for oxygen: $E = \frac{498\ 000\ J\ mol^{-1}}{6.02 \times 10^{23}\ mol^{-1}} = 8.27 \times 10^{-19}\ \text{«J»} \checkmark$ $\lambda = \frac{6.63 \times 10^{-34}\ J\ s \times 3.00 \times 10^{8}\ m\ s^{-1}}{8.27 \times 10^{-19}\ J} = 9.2.40 \times 10^{-7}\ \text{«m»} \checkmark$ $ALTERNATIVE\ 2:$ for ozone: similar calculation using 200 < bond enthalpy < 400 for ozone, such that the sum of the s	th as	Award [2] for correct final answer.	2
1. 1	f		$\bullet NO + O_3 \rightarrow \bullet NO_2 + O_2 \checkmark$ $\bullet NO_2 + O_3 \rightarrow \bullet NO + 2O_2 \checkmark$	GI	Accept $\bullet NO_2 \rightarrow \bullet NO + \bullet O$ AND $\bullet O + O_3 \rightarrow 2O_2$ for M2.	2



Topic Chem 4 Q# 49/ IB Chem/2018/w/TZ 0/P2/Higher Level/

3. b	•	i	Structure I – follows octet rule: [:O—Br→O:] / [:O—Br—O:] Structure II – does not follow octet rule: [:O—Br=O] OR OR OR OR OR OR OR	Accept dots, crosses or lines to represent electron pairs.	2
3. b	b	ii	«structure I» formal charge on Br = +2 OR «structure II» formal charge on Br = 0/+1 structure II is preferred AND it produces formal charge closer to 0 ✓	Ignore any reference to formal charge on oxygen.	2
3. с	,		Geometry: trigonal/pyramidal ✓ Reason: three bonds AND one lone pair OR four electron domains ✓ O-Br-O angle: 107° ✓	Accept "charge centres" for "electron domains". Accept answers in the range 104–109°.	3

Topic Chem 4 Q# 50/ IB Chem/2018/s/TZ 2/P2/Higher Level/

v	quesu	UII	Q# 50/ IB Chem/2018/s/TZ 2/P2/Higher Level/	Notes	Total
7.	a	i	polar bonds «between H and group 16 element» OR difference in electronegativities «between H and group 16 element» ✓ uneven distribution of charge/electron cloud OR non-linear/bent/V-shaped/angular shape «due to lone pairs» OR polar bonds/dipoles do not cancel out ✓	M2: Do not accept "net/overall dipole moment" without further explanation. Accept "non-symmetrical «shape/distribution of charge»".	2
7.	а	ii	number of electrons increases ✓ London/dispersion/instantaneous induced dipole-induced dipole forces increase ✓	M1: Accept "M,/A, increases" or "molecules become larger in size/mass/surface area".	2
7.	b		Electron domain geometry: tetrahedral ✓ Molecular geometry: bent/V-shaped/angular ✓	Both marks can be awarded for clear diagrams. Electron domain geometry requires a 3-D diagram showing the tetrahedral arrangement.	2



9 10			1					
7.	C	i	Structure: O atom labelled (1) O atom labelled (2)	0	II ≪+»1 −1	*	Award [1] for any two correctly filled cells.	2
7.	С	jii T	structure I AND no formal OR structure I AND no charge	. .	between aton	ns» ✓		11
7.	d		O ₃ has bond between sing OR O ₃ has bond order of 1.5 oR bond in O ₃ is weaker/long O ₃ requires longer wavele	AND O ₂ has	s bond order	M1: Do not accept "ozone has one single and one double bond".	2	
7.	е		CO ₂ «non-polar» «weak» induced dipole forces beth SiO ₂ network/lattice/3D/gi	veen moled	eules 🗸	s/instantaneous induced dipole-	M1: The concept of "between" is essential.	2

Topic Chem 4 Q# 51/ IB Chem/2018/s/TZ 1/P2/Higher Level/

3. d	equal C-C bond «lengths/strengths» OR	Accept "all C-C-C bond angles are equal".	
	regular hexagon		
	OR		1
	«all» C-C have bond order of 1.5		
	OR	//	
	«all» C−C intermediate between single and double bonds ✓	7	

Topic Chem 4 Q# 52/ IB Chem/2018/s/TZ 1/P2/Higher Level/

2.	a		electrostatic attraction AND oppositely charged ions ✓		1
2.	b		multiply relative intensity by «m/z» value of isotope OR find the frequency of each isotope ✓ sum of the values of products/multiplication «from each isotope» OR find/calculate the weighted average ✓	Award [1 max] for stating "m/z values of isotopes AND relative abundance/intensity" but not stating these need to be multiplied.	2
2.	С		«promoted» electrons fall back to lower energy level ✓ energy difference between levels is different ✓	Accept "Na and Ca have different nuclear charge" for M2.	2
2.	d	i	Any two of: stronger metallic bonding ✓ smaller ionic/atomic radius ✓ two electrons per atom are delocalized OR greater ionic charge ✓ greater atomic mass ✓	Do not accept just "heavier" or "more massive" without reference to atomic mass.	2
2.	d	ii	delocalized/mobile electrons «free to move» ✓		1



е е		1		
		x x x x x x x x x x x x x x x x x x x		2
L.S.		Number of electron removed general increase only one discontinuity between "IE2" and "IE3" ✓		
f		pH > 7 √	Accept any specific pH value or range of values above 7 and below 14.	1
g	i	sigma (σ): overlap «of atomic orbitals» along the axial/internuclear axis OR head-on/end-to-end overlap «of atomic orbitals» ✓ pi (π): overlap «of p-orbitals» above and below the internuclear axis OR sideways overlap «of p-orbitals» ✓	Award marks for suitable diagrams.	2
g	ii	sigma (σ): 3 AND pi (π): 2 √		1
oic C	hem 4	Q# 53/ IB Chem/2018/s/TZ 1/P2/Higher Level/		·
b		Electron geometry Molecular geometry	Note: Urea's structure is more complex	

1.	b			Electron geometry	Molecular geometry	Note: Urea's structure is more complex than that predicted from VSEPR theory.	
			Nitrogen	tetrahedral ✓	trigonal pyramidal 🗸	than that predicted from VSEPR theory.	3
			Carbon	trigonal planar ✓	togonal planar	-	
1.	e	i	urea has more hyd	ectron density/greater London/	dispersion √	Accept "urea has larger size/greater van der Waals forces". Do not accept "urea has greater intermolecular forces/IMF".	1
1.	e	H	H H H H H H H H H H H H H H H H H H H	+	THING)	Award [1] for each correct interaction. If lone pairs are shown on N or O, then the lone pair on N or one of the lone pairs on O MUST be involved in the H-bond. Penalize solid line to represent H-bonding only once.	2
1.	i		OR C-N bond partial d OR delocalization «of e	en atoms can be donated to/shouble bond character electrons occurs across molecu	le»		1

4.	а	i	triple bond in nitrogen «molecule» AND single bond in hydrazine ✓	Accept bond enthalpy values from data booklet (158 and 945 kJ mol ⁻¹).	
			triple bond is stronger than single bond OR	1820	2
			more shared «pairs of» electrons make bond stronger/attract nuclei more ✔		
	а	ii	hydrogen bonding «between molecules, dinitrogen tetraoxide does not» ✓		1
	b	i	lone pair on p orbital «of O atom» overlaps/delocalizes with pi electrons «from double bond» ✓ both O–O bonds have equal bond length	Accept "p/pi/T electrons are delocalized/not localized".	
			OR		
			both O=O bonds have same/1.5 bond order		3
			OR		3
			both O−O are intermediate between O−O AND O=O ✓		
			The state of the second of the second production of the second se		
			both O–O bonds have equal bond energy ✓		
4	b	ii	ALTERNATIVE 1:	Accept any combination of lines, dots or crosses to represent electrons.	
			(O O O O	Do not accept structure that represents 1.5 bonds.	
			FC: -1 AND +1 AND 0 ✓	Do not penalize missing lone pairs if already penalized in 3(b).	
			ALTERNATIVE 2:	If resonance structure is incorrect, no ECF.	2
			10 00	Any one of the structures with correct formal charges for [2 max].	
			FC: 0 AND +1 AND −1 ✓		
opi	c Che	em 4	Q# 55/ IB Chem/2017/s/TZ 2/P2/Higher Level/		
3	b		Lewis structure:	Penalize missing lone pairs once only between this question and 4(b)(ii).	
				Accept any combination of lines, dots or crosses to represent electrons.	
			Molecular geometry: trigonal/triangular pyramidal ✓	Do not apply ECF.	3
			Bond angles:	Do not accept answer equal to or less than 90°.	
			< 109.5° ✓	Literature value is 100.1°.	
opi	c Che	em 4	• Q# 56/ IB Chem/2017/s/TZ 2/P2/Higher Level/	5 5 4	
1	c	ii	«molten» Na₂O has mobile ions/charged particles AND conducts electricity ✓ «molten» P₄O₁o does not have mobile ions/charged particles AND does not	Do not award marks without concept of mobile charges being present.	
			conducts electricity/is poor conductor of electricity ✓	Award [1 max] if type of bonding or electrical conductivity correctly identified in each compound.	



2

identified in each compound.

electrons.

Do not accept answers based on

Award [1 max] if reference made to solution.

Topic Chem 4 Q# 57/ IB Chem/2017/s/TZ 1/P2/Higher Level/

8.	а	$NO \cdot (g) + O_3(g) \rightarrow NO_{2^*}(g) + O_2(g) \checkmark$	Allow representation of radicals without • if consistent throughout.	
		$NO_{2^*}(g) + O_*(g) \rightarrow NO_*(g) + O_2(g)$ OR	70 F.F	2
		$NO_{2^{\bullet}}(g) + O_{3}(g) \rightarrow NO_{1}(g) + 2O_{2}(g) \checkmark$		2
8.	b	«loss of ozone» allows UV radiation to penetrate atmosphere/reach earth ✓		
		UV radiation causes skin cancer		2
		OR		
		UV radiation causes tissue damage ✓		

Topic Chem 4 Q# 58/ IB Chem/2017/s/TZ 1/P2/Higher Level/

5.	а	107° ✓	Accept 100° to <109.5°. Literature value = 105.8°	1
5.	b	tetrahedral ✓ sp³ ✓	No ECF allowed.	2

Topic Chem 4 Q# 59/ IB Chem/2017/s/TZ 1/P2/Higher Level/

2.	a		electrostatic attraction ✓ between «a lattice of» metal/positive ions/cations AND «a sea of» delocalized electrons ✓	Accept "mobile electrons". Do not accept "metal atoms/nuclei".	2
2.	d	ii	vanadium has smaller ionic radius «leading to stronger metallic bonding» 🗸	Accept vanadium has «one» more valence electron«s» «leading to stronger metallic bonding». Accept "atomic" for "ionic".	1
2.	d	iv	titanium atoms/ions distort the regular arrangement of atoms/ions OR titanium atoms/ions are a different size to aluminium «atoms/ions» ✓ prevent layers sliding over each other ✓	Accept diagram showing different sizes of atoms/ions.	2
2.	g	i	ionic OR «electrostatic» attraction between oppositely charged ions ✓		1
2.	g	ii	«simple» molecular structure OR weak«er» intermolecular bonds OR weak«er» bonds between molecules ✓	Accept specific examples of weak bonds such as London/dispersion and van der Waals. Do not accept "covalent".	1

Topic Chem 4 Q# 60/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

4. h	«3-D/giant» regularly repeating arrangement «of ions» OR	Accept "giant" for M1 unless "giant covalent" stated.	
	lattice «of ions» ✓	73	2
	electrostatic attraction between oppositely charged ions	Do not accept "ionic" without description.	- 2
	OR		
	electrostatic attraction between Mg ²⁺ and O ^{2−} ions ✓		

Topic Chem 4 Q# 61/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

Questio	n Answers	Notes	Total
2. d		Accept single negative charges on two O atoms singly bonded to C. Do not accept resonance structures. Allow any combination of dots/crosses or lines to represent electron pairs.	1
2. е	electrons delocalized «across the O–C–O system» OR resonance occurs ✓	Accept delocalized π-bond(s). No ECF from (d).	
	122 «pm» < C−O < 143 «pm» ✓	Accept any answer in range 123 «pm» to 142 «pm». Accept "bond intermediate between single and double bond" or "bond order 1.5".	2



Topic Chem 4 Q# 62/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

5.	b	i	Sigma (σ):	OR O				
			Pi (π):					2
i.	b	ii		Number of sigma (a) bonds	Number of pi (π) bonds		Award [1] for two or three correct answers. Award [2] for all four correct.	
			Propane	10	0	1		2

Topic Chem 4 Q# 63/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

(Question	Answers	Notes	Total
1.	d	ethane-1,2-diol can hydrogen bond to other molecules «and ethene cannot» OR ethane-1,2-diol has «significantly» greater van der Waals forces ✓	Accept converse arguments. Award [0] if answer implies covalent bonds are broken.	2
		hydrogen bonding is «significantly» stronger than other intermolecular forces ✓		

Topic Chem 4 Q# 64/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

Q	uesti	ion		Answers		Notes	Total
5.	а	i	Ozone Ozone Sulfur hexafluoride	o / F: F: F: F:	cture	Lines, x's or dots may be used to represent electron pairs. Charges may be included in Lewis structures of ozone but are not required.	2
	а	ii		Electron domain geometry	Molecular geometry	Award [1 max] for either both electron domain geometries correct OR for either both molecular geometries correct.	
			Ozone	trigonal/triangular planar	v-shaped/bent/angular ✓	A V	2
			Sulfur hexaffuoride	octahedral/square bipyramidal	octahedral/square bipyramidal ✓	10	
	а	iii	sulfur hexafluoride/SF	6	AND DESCRIPTION OF THE PERSON		1
	а	iv	Ozone: Accept any ang and Sulfur hexafluoride: 90		but less than 120	Experimental value of bond angle in O3 is 117'.	1

Que		ion			Answ	ers		Notes	Total
6.	a	v	o.	Ö,	;o ⁰ .o;	-		Double-headed arrow not necessary for mark. Lines, x's or dots may be used to represent electron pairs.	1
	b	i		Lewis (electron dot) structure	FC of O on LHS	FC of central N	FC of N on RHS		
			A	:ö=n=ä:	0	+1	-1	Award [2] for all nine FCs correct, [1] for	,
			В	:ö—n≡n:	-1	+1	0	six to eight FCs correct.	-
			С	:o≡n—ÿ:	+1	+l	-2		
	b	ii	smal	llest FC difference for A	or B, so eithe	r is preferred	1	Reason required for M1.	
				ever B is preferred as oxy gh FC per se ignores elec			ve than nitrogen, even	OWTTE	2

(Question		Answers	Notes	Total
6.	d	i	O ₂ has a double bond ✓ O ₃ has intermediate bonds between double and single bonds OR O ₃ has a bond order of 1½ ✓ bond in O ₂ is stronger therefore I needs more energy ✓	Do not award mark for I on its own with no justification.	3
	đ	ii	C-Cl (bond) breaks since weakest bond \checkmark $CCl_{2}F_{2} \xrightarrow{Ono} \cdot CClF_{2} + Cl \cdot \checkmark$ $Cl_{2} + O_{3} \rightarrow ClO_{2} + O_{2} \checkmark$ $Cl_{2} + O_{3} \rightarrow Cl_{2} + Cl_{3} \checkmark$ $Cl_{3} + O_{4} \rightarrow O_{2} + Cl_{4} \checkmark$ $Cl_{4} + O_{5} \rightarrow Cl_{4} + 2O_{3} \checkmark$	Allow representation of radicals without * as long as consistent throughout.	5

Topic Chem 4 Q# 65/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

(Question		Answers	Notes	Total
3.	c	i	ALTERNATIVE 1: σ-bond from N to N AND from N to O ✓ π-bond from N to N ✓ delocalized π-bond/π-electrons extending over the oxygen and both nitrogenss ✓ ALTERNATIVE 2: both have 2 σ-bonds efrom N to N and from N to Os AND π-bond from N to N ✓ one structure has second π-bond from N to N and the other has π-bond from N to O ✓ delocalized π-bond/π-electrons ✓	Award [1 max] if candidate has identified both/either structure having 2 σ-bonds and 2 π-bonds.	3
3.	С	ii	more than one possible position for a multiple/π-/pi- bond ✔	Accept "more than one possible Lewis structure". Accept reference to delocalisation if M3 not awarded in c (i). Accept reference to fractional bond orders.	1

Topic Chem 4 Q# 66/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

-	Juest	ion	Answers	Notes	lotal
1.	а	i	H—P—H	Accept structures using dots and/or crosses to indicate bonds and/or lone pair.	1
1.	a	ii	sp³ ✓	Do not allow ECF from a (i).	1
1.	а	iii	Lewis base AND has a lone pair of electrons ¢to donates ✓		1
1.	а	iv	non-polar AND P and H have the same electronegativity ✓	Accept "similar electronegativities". Accept "polar" if there is a reference to a small difference in electronegativity and apply ECF in 1 a (v).	1
1.	а	v	PH ₃ has London 《 dispersion 》 forces ✓ NH ₃ forms H-bonds ✓ H-bonds are stronger OR London forces are weaker ✓	Accept van der Waals' forces, dispersion forces and instantaneous dipole – induced dipole forces. Accept "dipole-dipole forces" as molecule is polar. H-bonds in NH ₃ (only) must be mentioned to score [2]. Do not award M2 or M3 if: implies covalent bond is the H-bond implies covalent bonds break. Accept "dipole-dipole forces are weaker".	2 max
1.	а	vi	Weak: only partially dissociated/ionized €in dilute aqueous solution Bransted-Lowry base: an acceptor of protons/H*/hydrogen ions	Accept reaction with water is reversible/an equilibrium. Accept "water is partially dissociated to the weak bases".	2
1.	b	i	P₄ is a molecule comprising 4P atoms AND 4P is four/separate <ps 4p="" <4ps="" <4s="" <ps="" and="" atoms="" bonded="" non-bonded="" or="" p₄="" represents="" separate="" td="" together="" ✓<=""><td></td><td>1</td></ps>		1

Topic Chem 4 Q# 67/ IB CHEM/2015/w/TZ0/P2 Section B/Higher Level/

(c) (i)		NF ₄ ⁺	N ₂ H ₂	N_2H_4
	Hybridization	sp³;	sp²;	sp³;

[3]

Do not penalize if it is not superscript.


 sigma bond: overlapping of orbitals end to end/head on / overlapping of orbitals along internuclear axis;
 Accept "axial overlapping of orbitals".

pi bond: side-ways/parallel overlapping of <u>p orbitals</u> (above and below internuclear axis);

[2]

Accept "overlapping of <u>p orbitals</u> above and below the internuclear axis". Accept suitable labelled diagrams.

(iii)



[4]

Accept any combination of lines, dots or crosses to represent electron pairs.

Penalize missing lone pairs on fluorine atoms once only.

For SF4, lone pair on S required for the mark.

Do not allow ECF for a wrong Lewis structure.

Accept saw horse/distorted tetrahedron instead of see-saw.

(d) (i) CH₃CH₂CH₃ < CH₃CHO < CH₃CH₂OH < CH₃COOH ;; Award [2] for correct order.

[2]

Award [1] for CH₃COOH > CH₃CH₂OH > CH₃CHO > CH₃CH₂CH₃ as compounds are not listed in order of increasing boiling point.

Award [1] if one error in the order.

(ii) CH₃CH₂CH₃ /London/dispersion/instantaneous induced dipole-induced dipole forces

CH₃CHO dipole-dipole forces (and London/dispersion forces)

CH₃CH₂OH H-bonding (and dipole-dipole and London/dispersion forces)

CH₃COOH H-bonding (and dipole-dipole and London/dispersion forces);; Award [2] for all four correct.

Award [1] for two or three correct.

H-bonding strongest / London/dispersion forces weakest / dipole-dipole stronger than London/dispersion / dipole-dipole weaker than H-bonding;

Accept van der Waals' forces for London/dispersion forces.

CH₃COOH forms more/stronger H-bonds than CH₃CH₂OH / CH₃COOH is more polar than CH₃CH₂OH;

Accept CH₃COOH has more electrons/higher molar mass than CH₃CH₂OH.

[4]

(e) ionization and (bombardment) by high energy/fast moving electrons/electron gun (to form positive ions);

acceleration and passing through electric field/potential difference/oppositely charged plates;

deflection and passing through magnetic field/electromagnet;

Award [1] for naming 3 processes (ionization, acceleration, deflection) in the correct order with incorrect details.



Topic Chem 4 Q# 68/ IB Chem/2015/w/TZ0/P2 Section A/Higher Level/

 (Li → Cs) atomic/ionic radius increases; force of attraction between metal ions and <u>delocalized</u> electrons decreases; Accept metallic bonding gets weaker.

 $(F \rightarrow I)$ London/dispersion/instantaneous induced dipole-induced dipole forces increase;

Accept vdW/van der Waals' forces for London/dispersion forces.

with increasing number of electrons/molar mass/surface area/size of electron cloud;

[3 max]

[2]

Do not accept "with increasing size" or "with increasing mass" only.

Topic Chem 4 Q# 69/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(c) (i)

Penalize lone pairs missing on Br and F once only.

Accept any combination of lines, dots or crosses to represent electron pairs.

Shape:	Shape:
trigonal pyramidal;	
Accept triangular pyramidal.	see-saw/K-shaped; Accept distorted tetrahedral.
Accept triangular pyramidal.	Accept distorted tetranedral.
No ECF for shape if Lewis stru	cture is incorrect.
Bond angle:	Bond angle:
Any value 99 < 109 ;	Any two for [1] of:
Literature value = 101".	(Feg-S-Feg): Any value 100 <120
	Literature value = 103°
	(F _{ax} -S-F _{ax}): Any value 175 *<185 *
	Literature value = 179°/180°
	(Fax-S-Feq): Any value 85° < 95°
	Literature value = 89°

(iii) P–Br and S–F bonds are polar / bonds in both molecules are polar; non-symmetrical distribution of electron cloud / polar bonds/dipoles do not cancel because of non-symmetrical shape;
 M2 may also be scored with a suitable diagram showing the vectorial addition of the individual S–F dipole moments to show a net dipole moment centred along the axis between the F_{eq}–S–F_{eq} bond.



EITHER (d) (i) OR (11) (iii)

(electrostatic) attraction between (positively charged) nuclei and a pair of

formed as a result of electron sharing (between the carbon and hydrogen nuclei);

sigma bond formed by overlap of atomic orbitals; s orbital from H and p/sp2 from carbon;

[2]

 α : sp³ and β : sp²; Accept if numbers are given as subscripts.

[1]

σ bond: end-on / axial overlap of two orbitals;

 π bond:

sideways overlap of two (parallel) p orbitals;

[2]

Accept suitable diagrams for both marks.

(iv) 11σ and 3π ;

[1]

Topic Chem 4 Q# 70/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

chromium(III) oxide; (i) Do not award the mark for chromium oxide. [1]

(ii) (electrostatic) attraction between positive and negative ions/oppositely charged ions/Cr3+ and O2-; formed as a result of electron transfer from chromium atoms to oxygen atoms / OWTTE: Ignore reference to number of electrons transferred or charges of ion for M2.

[2]

(iii) ions are not free to move (when solid) / ions in rigid lattice / OWTTE;

[1]

[2]

Topic Chem 4 Q# 71/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(f) Sigma bonds:



result from head-on/end-on overlap of orbitals / OWTTE;

Accept axial overlap of orbitals.

Accept "symmetric orbital" with respect to same plane / OWTTE.

Pi bonds



result from sideways overlap of orbitals / OWTTE;

Accept "antisymmetric orbitals" with respect to (defining) plane (containing at

least one atom) / OWTTE.



Topic Chem 4 Q# 72/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(g) Allotropes:

Any three allotropes for [1] from:

diamond

graphite

fullerene

graphene;

Allow (carbon) nanotubes for graphene.

Accept C₆₀ /C₇₀ /buckminsterfullerene/bucky balls for fullerene.

Structures:

Any three for [3] from:

Diamond:

tetrahedral arrangement of (carbon) atoms/each carbon bonded to four others / sp3 and 3D/covalent network structure;

Graphite

each carbon bonded to three others (in a trigonal planar arrangement) / sp² and 2D / layers of (carbon) atoms;

Fullerene:

each (carbon) atom bonded to three others (in a trigonal arrangement) / sp² and joined in a ball/cage/sphere/connected hexagons and pentagons;

Accept "trigonal planar" for "each carbon atom bonded to three others" part in M4. Graphene:

each carbon bonded to three others (in a trigonal arrangement) / sp² and 2D structure:

[4]

Topic Chem 4 Q# 73/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(e) Cl has 7 valence electrons and is in group 7; Accept "group 17" as suggested by IUPAC.

Ct has 3 occupied (electron) shells/energy levels and so is in period 3;

[2]

[3]

[2]

[2]

(f)	Molecule	Shape	Bond angle	Polarity
36	scl ₂	bent/angular/ v-shaped	< 109.5° Accept 100° – 108°. Literature value is 103°.	polar
	SCIF ₅	Octahedral Accept square bipyramidal.	90° (180°)	polar

Do not accept ECF for bond angles and polarities from incorrect shapes.

Award [3] for all six correct.

Award [2] for four or five correct.

Award [1] for two or three correct.

Topic Chem 4 Q# 74/ IB Chem/2015/s/TZ1/P2 Section A/Higher Level/

(b) (i) Na₂O ionic and P₄O₁₀ covalent (within molecule);

 Na_2O in giant/3D/network/lattice structure with strong (ionic) bonds **and** P_4O_{10} has a (simple) molecular structure/weak intermolecular forces (between molecules);

Award [1] for stating that bonds require more energy to break in Na₂O than in P₄O₁₀.

ii)		Phosphorus(V) oxide	Sodium oxide
	Solid state	no	no
	Molten state	no	yes

Award [2] for four correct.

Award [1] for two or three correct.



Topic Chem 4 Q# 75/IB Chem/2015/s/TZ1/P2 Section A/Higher Level/

(c) electrostatic attraction;

between (a lattice of) cations/positive ions **and** delocalized/sea of electrons; Do not award any mark for only stating "metallic bonding". [2]

Topic Chem 4 Q# 76/ IB Chem/2015/s/TZ1/P2 Section A/Higher Level/

(c) hydrogen bonding;

[1]

$$(d) \qquad : \overset{\times}{\circ} \overset{\times}{\circ} \overset{\times}{\circ} : \\ H \times \cdot \overset{\circ}{\circ} \cdot \times \overset{\times}{\circ} \overset{\times}{\circ} \times \overset{\times}{\circ} \times \overset{\circ}{\circ} : \\ H \xrightarrow{} \vdots \overset{\times}{\circ} \overset{\times}{\circ} \overset{\times}{\circ} : \\ \vdots \overset{\times}{\circ} \overset{\times}{\circ} : \overset{\times}$$

[1]

Mark cannot be scored if lone pairs are missing on oxygens.

Accept any combination of lines, dots or crosses to represent electron pairs.

(e) Acid:

one double and one single bond / one shorter and one longer bond; Accept "two double and two single".

Conjugate base:

two 1.5 bonds / both bonds same length;

Accept "four / all".

electrons delocalized / resonance forms;

Award marks for suitable diagrams.

[3]

Topic Chem 5 Q#77/IB Chem/2019/w/TZ0/P2/Higher Level/Q7./

6.	a	ii	$\Delta H^{0} = \Sigma \Delta H^{0}_{t} \text{ (products)} - \Sigma \Delta H^{0}_{t} \text{ (reactants)} \checkmark$ $\Delta H^{0} = 2(-241.8 \text{ «kJ mol}^{-1}\text{»}) - 4(-92.3 \text{ «kJ mol}^{-1}\text{»}) = -114.4 \text{ «kJ»} \checkmark$	Award [2] for correct final answer.	2
6.	d		enthalpy of solution = lattice enthalpy + enthalpies of hydration «of Cu²+ and Cl⁻» ✓	Accept enthalpy cycle.	
			«+2824 kJ mol⁻¹ − 2161 kJ mol⁻¹ − 2(359 kJ mol⁻¹) =» −55 «kJ mol⁻¹» ✓	Award [2] for correct final answer.	2

opic Chem 5 Q# 78/ IB Chem/2019/w/TZ0/P2/Higher Level/Q4./

3.	С	i	$2C_3H_8O(l) + 9O_2(g) \rightarrow 6CO_2(g) + 8H_2O(g)$ OR $C_3H_8O(l) + 4.5O_2(g) \rightarrow 3CO_2(g) + 4H_2O(g)$		1
3.	С	ii	bonds broken: 7(C-H) + C-O + O-H + 2(C-C) + 4.5(O=O) OR 7(414 «kJ mol⁻¹») + 358 «kJ mol⁻¹» + 463 «kJ mol⁻¹» + 2(346 «kJ mol⁻¹») + 4.5(498 «kJ mol⁻¹») / 6652 «kJ» ✓ bonds formed: 6(C=O) + 8(O-H) OR 6(804 «kJ mol⁻¹») + 8(463 «kJ mol⁻¹») / 8528 «kJ» ✓ «ΔH = bonds broken - bonds formed = 6652 - 8528 =» -1876 «kJ mol⁻¹» ✓	Award [3] for correct final answer.	3

Topic Chem 5 Q# 79/ IB Chem/2018/w/TZ 0/P2/Higher Level/

7.	b	α(-286 kJ) + (-1411 kJ) =» -1697 «kJ» ✓	1
7.	С	α -1697 kJ + 1561 kJ =» -136 «kJ» OR $\alpha \Delta H^{\Theta} = \Delta H_{f}^{\Theta}$ (products) - ΔH_{f}^{Θ} (reactants) = -84 kJ - 52 kJ =» -136 «kJ» ✓	1



7.	d	Accurate:	
		no approximations were made in the cycle	
	1 1	OR	
	1 1	values are specific to the compounds	
	1 1	OR	
	1 1	Hess's law is a statement of conservation of energy	
	1 1	OR	
	1 1	method is based on a law	
		OR	
		data in table has small uncertainties ✓	
		Approximate:	2
		values were experimentally determined/had uncertainties	
	1 1	OR	
	1 1	each value has been determined to only three/four significant figures	
	1 1	OR	
	1 1	different sources have «slightly» different values for enthalpy of combustion	
	1 1	OR	
		law is valid until disproved	
	1 1	OR	
		law of conservation of energy is now conservation of mass-energy	
		OR	
		small difference between two guite large terms «leads to high percentage uncertainty» ✓	

Topic Chem 5 Q# 80/ IB Chem/2018/w/TZ 0/P2/Higher Level/

5.	b	negative AND fewer moles/molecules «of gas» in the products ✓		1
5.	С	$\Delta G^{\circ} = \kappa - RT \ln K_c = w - 8.31 \text{ J K}^{-1} \text{ mol}^{-1} \times 1000 \text{ K} \times \ln 280$ OR $\Delta G^{\circ} = -4.7 \times 10^4 \text{ «J »} \checkmark$	Award [2] for correct final answer.	2
		«∆G ^O =» -47 «kJ» √		
5.	d	ΔG° < 0/spontaneous AND ΔS° < 0/unfavourable \checkmark exothermic AND ΔH° «must be» negative/favourable \checkmark		2

Topic Chem 5 Q# 81/ IB Chem/2018/w/TZ 0/P2/Higher Level/

4.	d	Fe ²⁺ AND smaller size/radius	M1 not needed for M2.	
		OR STATE OF THE ST		
		Fe ²⁺ AND higher charge density √	4 5 4	2
3		stronger interaction with «polar» water molecules ✓		



Topic Chem 5 Q# 82/ IB Chem/2018/w/TZ 0/P2/Higher Level/

3 851		\$4.500 to to to to total table to	E	3
1. b	•	ALTERNATIVE 1: $q = \&80.0 \text{ g} \times 4.18 \text{ J g}^{-1} \text{ K}^{-1} \times 7.5 \text{ K} = \&2.5 \times 10^{3} \text{ «J»/2.5 «kJ»} \checkmark$ $ &\text{ wper mol of } \text{ CuSO}_{4} = \frac{-2.5 \text{ kJ}}{0.0160 \text{ mol}} = -1.6 \times 10^{2} \text{ kJ mol}^{-1} \text{ w} $ $ &\text{ wfor the reaction} \&\Delta H = -1.6 \times 10^{2} \text{ «kJ»} \checkmark $ $ &\text{ALTERNATIVE 2:} $ $ &q = \&80.0 \text{ g} \times 4.18 \text{ J g}^{-1} \text{ K}^{-1} \times 7.5 \text{ K} = \&2.5 \times 10^{3} \text{ «J»/2.5 «kJ»} \checkmark $ $ &\text{ wncu} = \frac{0.872}{63.55} = 0.0137 \text{ mol} \text{ w} $ $ &\text{ wper mol of } \text{ CuSO}_{4} = \frac{-2.5 \text{ kJ}}{0.0137 \text{ mol}} = -1.8 \times 10^{2} \text{ kJ mol}^{-1} \text{ w} $ $ &\text{ wfor the reaction} \&\Delta H = -1.8 \times 10^{2} \text{ kJ w} \checkmark $	Award [2] for correct final answer.	2
1. b	ii	density «of solution» is 1.00 g cm ⁻³ OR specific heat capacity «of solution» is 4.18 J g ⁻¹ K ⁻¹ /that of «pure» water OR reaction goes to completion OR iron/CuSO₄ does not react with other substances ✓	The mark for "reaction goes to completion" can only be awarded if 0.0160 mol was used in part (b)(i). Do not accept "heat loss".	1

Topic Chem 5 Q# 83/ IB Chem/2018/s/TZ 1/P2/Higher Level/

3.	С	i	ΔH^{\odot} = bonds broken − bonds formed \checkmark « ΔH^{\odot} = 3(C≡C) − 8(C := C) _{benzene} / 3 × 839 − 6 × 507 / 2517 − 3042 =» −525 «kJ» \checkmark	Award [2] for correct final answer. Award [1 max] for "+525 «kJ»". Award [1 max] for: « $\Delta H^0 = 3(C \equiv C) - 3(C - C) - 3(C = C) / 3 \times 839 - 3 \times 346 - 3 \times 614 / 2517 - 2880 = n - 363 «kJ».$	2
3.	С	ii	$\Delta H^{\Theta} = \Sigma \Delta H_{t}$ (products) $-\Sigma \Delta H_{t}$ (reactants) \checkmark $\ll \Delta H^{\Theta} = 49 \text{ kJ} - 3 \times 228 \text{ kJ} = 8 -635 \text{ kJ} $	Award [2] for correct final answer. Award [1 max] for "+635 «kJ»".	2
3.	c	III	ΔH _f values are specific to the compound OR bond enthalpy values are averages «from many different compounds» ✓ condensation from gas to liquid is exothermic ✓	Accept "benzene is in two different states «one liquid the other gas»" for M2.	2
3.	С	iv	«∆S° = 173 - 3 × 201 = » -430 «J K ⁻¹ » ✓	4 6 6	1
3.	С	v	T = «25 + 273 =» 298 «K» \checkmark ΔG^{\oplus} « = −635 kJ − 298 K × (−0.430 kJ K ⁻¹)» = −507 kJ \checkmark ΔG^{\oplus} < 0 AND spontaneous \checkmark	$\Delta G^0 < 0$ may be inferred from the calculation.	3



opic C	Chei	m 5	Q#84/ IB Chem/2017/s/TZ 2/P2/Higher Level/		v.
9. a		ī	$\Delta H = 177.0 - \frac{189.2}{2} - 285.5 \text{ «kJ»} \checkmark$ $\text{«}\Delta H = \text{»} - 203.1 \text{ «kJ»} \checkmark$	Accept other methods for correct manipulation of the three equations. Award [2] for correct final answer.	2
a	1	ii	203.1 «kJ» = 0.850 «kg» × 4.18 «kJ kg ⁻¹ K ⁻¹ » × ΔT «K» OR «ΔT =» 57.2 «K» ✓ «Tfinal = (57.2 + 21.8)°C =» 79.0 «°C» / 352.0 «K» ✓ If 200.0 kJ was used: 200.0 «kJ» = 0.850 «kg» × 4.18 «kJ kg ⁻¹ K ⁻¹ » × ΔT «K» OR «ΔT =» 56.3 «K» ✓ «Tfinal = (56.3 + 21.8)°C =» 78.1 «°C» / 351.1 «K»✓	Award [2] for correct final answer. Accept two, three or four significant figures. Unit, if specified, must be consistent with value stated.	2
b)	i	«∆S° =» 270 «J K⁻¹ mol⁻¹» − 267 «J K⁻¹ mol⁻¹» − 131 «J K⁻¹ mol⁻¹» ✓ «∆S° =» −128 «J K⁻¹ mol⁻¹» ✓	Award [2] for correct final answer.	2
b)	ii	«non spontaneous if» $\Delta G^{\circ} = \Delta H^{\circ} - T\Delta S^{\circ} > 0$ OR $\Delta H^{\circ} > T\Delta S^{\circ} \checkmark$ « T above» $\frac{-124.4 \text{ «kJ mol}^{-1} \text{»}}{-0.128 \text{ «kJ K}^{-1} \text{mol}^{-1} \text{»}} = 972 \text{ «K»} \checkmark$	Award [2] for correct final answer. Accept 699 °C. Do not award M2 for any negative T value.	2
opic C	hei	m 5	Q# 85/ IB Chem/2017/s/TZ 1/P2/Higher Level/		
4. c	\rightarrow	i	$\Sigma S^{O}_{RHS} = 313.4 \text{ «J K}^{-1} \text{»}$ AND $\Sigma S^{O}_{LHS} = \text{ «(4 × 197.6)} + 29.9 \text{ J K}^{-1} = \text{» 820.3 «J K}^{-1} \text{»} \checkmark$ $\Delta S^{O} \text{ «= } \Sigma S^{O}_{RHS} = \Sigma S^{O}_{LHS} = 313.4 - 820.3 \text{»} = -508.9 \text{ «J K}^{-1} \text{»} \checkmark$	Award [2] for correct final answer.	2
4. c		ii	ΔH ^O «= -833.0 - 4 × (-110.5)» = -191 «kJ» ✓		1
4. c		iii	«when» ΔG = 0 «forward and backward reactions are equally favourable» \checkmark «when ΔG = 0, T = $\frac{\Delta H}{\Delta S}$ », T = « $\frac{191 \text{ kJ}}{0.5069 \text{ kJ K}^{-1}}$ =» 377 «K» \checkmark «temperature =» 104 «°C» \checkmark	Award [3] for correct final answer. Use of -500 J K ⁻¹ and -200 kJ gives 127 °C. Award [2 max] for T < 104 «°C». Accept ∆G < 0 and T > 104 «°C».	3

tetracarbonylnickel is highly flammable «auto-ignition temperature of 60 °C» ✓ Topic Chem 5 Q# 86/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

Ni(CO)4 decomposition deposits nickel in the lungs

tetracarbonylnickel is toxic/poisonous OR

CO is toxic/poisonous

OR

1.	a	iii	Bonds broken: 2C≡O + 3(H−H) / 2(1077 kJ mol⁻¹) + 3(436 kJ mol⁻¹) / 3462 «kJ» ✓ Bonds formed: 2(C−O) + 2(O−H) + 4(C−H) + (C−C) / 2(358 kJ mol⁻¹) + 2(463 kJ mol⁻¹) + 4(414 kJ mol⁻¹) + 346 kJ mol⁻¹ / 3644 «kJ» ✓		3
			«Enthalpy change = bonds broken – bonds formed = 3462 kJ – 3644 kJ =» –182 «kJ» ✓	Award [3] for correct final answer. Award [2 max] for «+»182 «kJ».	
1.	b	i	$\alpha \Delta H = \Sigma \Delta H_r$ products − $\Sigma \Delta H_r$ reactants = −454.8 kJ mol ⁻¹ − 2(−110.5 kJ mol ⁻¹) =» −233.8 α kJ» \checkmark		1
1.	b	II	in (a)(iii) gas is formed and in (b)(i) liquid is formed OR products are in different states OR conversion of gas to liquid is exothermic OR conversion of liquid to gas is endothermic OR enthalpy of vapourisation needs to be taken into account ✓	Accept product is «now» a liquid. Accept answers referring to bond enthalpies being means/averages.	1

1

4. d


(Question		Answers	Notes	Total
1.	b	iii	«∆S is negative because five mols of» gases becomes «one mol of» liquid OR increase in complexity of product «compared to reactants» OR product more ordered «than reactants» ✓	Accept "fewer moles of gas" but not "fewer molecules".	1
1.	b	iv	$\Delta S = \left(\frac{-620.1}{1000}\right) \text{ «kJ K}^{-1} \text{» } \checkmark$ $\Delta G = -233.8 \text{ kJ} - \left(298 \text{ K} \left(\frac{-620.1}{1000}\right) \text{ kJ K}^{-1}\right) = -49.0 \text{ «kJ} \text{» } \checkmark$	Award [2] for correct final answer. Award [1 max] for $\alpha+3185\times10^3$. If -244.0 kJ used, answer is: $\Delta G = -244.0$ kJ $-(298 \text{ K} \left(\frac{-620.1}{1000}\right) \text{ kJ K}^{-1}) = -59.2$ «kJ» Award [2] for correct final answer.	2
1.	b	v	increasing T makes ∆G larger/more positive/less negative OR -T∆S will increase ✓		1

Topic Chem 5 Q# 87/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

c	i	$CH_4(g) + 5O_2(g) \rightarrow CO_2(g) + 2H_2O(g) + 2O_3(g)$	1
c	ii	$\Delta H^{\circ} = \langle [(-393.5) + (2)(-241.8) + (2)(+142.3)] - [(-74.0)] = \rangle - 660.8 \text{ (kJ mol}^{-1}) \checkmark$	1
c	iii	standard enthalpy change of formation ΔH_{ρ}^{0} of an element (in most stable form) is always zero \checkmark	1
c	iv	$\Delta S^{\Theta} = \langle [(+213.8) + (2)(+188.8) + (2)(+237.6)] - [(+186) + (5)(+205.0)] = \rangle$ $-144.4 \langle J K^{-1} \text{ mol}^{-1} \rangle \checkmark$	1
c	v	$\Delta G^{\circ} = \left\langle \Delta H^{\circ} - T \Delta S^{\circ} = (-660.8) - (298) \left(\frac{-144.4}{1000} \right) = \right\rangle - 617.8 \text{dkJ} \text{mol}^{-1} \rangle \checkmark$	1
c	vi	spontaneous since negative ΔG° ✓	1

a double bond ✓ intermediate bonds between double and single bonds a bond order of 1½ ✓	Do not award mark for I on its own with no justification.	3
n O₂ is stronger therefore I needs more energy ✓	210	
cbond: breaks since weakest bond \checkmark $CCIF_2 + CI \cdot \checkmark$ $CO_3 \rightarrow CIO \cdot + O_2 \checkmark$ $CO_4 \rightarrow O_2 + CI \cdot \checkmark$	Allow representation of radicals without - as long as consistent throughout.	5
	0 ₃ → ClO++0 ₂ ✓	$O_3 \rightarrow ClO_4 + O_2 \checkmark$ + $O_3 \rightarrow O_3 + Cl_4 \checkmark$

Topic Chem 5 Q# 88/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/



Topic Chem 5 Q# 89/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

9	Question	Answers	Notes	Total
5.	a	only water/H₂O produced ⟨so non-polluting⟩ ✓		1
	b	Bond breaking: (1)(H-H)+(4)(C-H)+(1)(C=C) OR (1)(436)+(4)(414)+(1)(614) = 2706 ⟨kJ mol ⁻¹ ⟩ ✓		
		Bond formation: (6)(C-H)+(1)(C-C) OR (6)(414)+(1)(346) = 2830 (kJ mol ⁻¹) \checkmark (+2706-2830) = -124(kJ mol ⁻¹) \checkmark	Award [2 max] for +124 (kJ mol ⁻¹). Award [3] for correct final answer.	3

Topic Chem 5 Q# 90/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

0	Questi	ion	Answers	Notes	Total
3.	b		Activation energy / E _s / 63 • NJ • Reactants Enthalpy or reaction / 2H / -664 • NJ • Progress of reaction products lower than reactants AND enthalpy of reaction correctly marked and labelled with name or value activation energy correctly marked and labelled with name or value	Accept other clear ways of indicating energy/ enthalpy changes.	2
3.	b	ii	Reactants Catalysed Products Progress of reaction lower dotted curve, between same reactants and products levels, labelled "Catalysed" ✓		1

Topic Chem 5 Q# 91/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

1.	d	i	temperature rise € = $\frac{750 \times 1.00}{0.2000 \times 1.00}$ > = 3750 €°C/K > ✓	Do not accept -3750.	1

Topic Chem 5 Q# 92/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

4.	a	i	same charge AND same/similar ionic radius ✓	TATE OF THE PARTY	1
4.	а	ii	enthalpy of hydration ← -1483 + 2 (-359)»2201 ←kJ mol** ✓ enthalpy of solution ← 2170 - 2201»31 ←kJ mol** ✓	Award [2] for correct final answer. Award [1 max] for +31 ckJmot's. Award [1 max] for ±4371.	2

Topic Chem 5 Q# 93/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

2. a	iii	ΔH° = -220.1 - (-110.5) ✓ ΔH° = -109.6 « kJ» ✓	Award [2] for correct final answer. Award [1] for −330.6, or +109.6 ck.lp.	2
2. a	iv	$\Delta G^{\circ} = -109.6 - (298 \times \Delta S^{\circ}) = +11.7 \text{ (kJ)} \checkmark$ $\Delta S^{\circ} = -\frac{(11.7 + 109.6) \times 10^{3}}{298} = -407 \text{ (J K}^{-1}) \checkmark$	Award [2] for correct final answer. Award [2] for -470 (I) (result from given values). Do not penalize wrong value for T if already done in (a)(ii). Award [1 max] for -0.407 (k) K ¹ Award [1 max] for -138.9 (J) K ¹ Award [1 max] for -138.9 (J) K ¹ Award [1 max] for -138.9 (L) K ¹ (L)	2



Topic Chem 5 Q# 94/ IB CHEM/2015/w/TZ0/P2 Section B/Higher Level/

(a) possible to get iron and 3:1 ratio of the two equations;

The ratio may be shown in combining the equations/calculating overall ΔG /=29 (kJ).

possible to get iron and resultant ΔG is negative/-29 (kJ);

"Possible to get iron" needs only to be stated once.

Do not accept "possible to get iron" without any reason.

Do not accept "it is not possible to get iron because ΔG is positive".

(b) (i)
$$\Delta G^{\Theta} = [-1582 + 2 \times 10] - [-742];$$

$$\Delta G^{\Theta} = -820 \text{ (kJ mol}^{-1}\text{)};$$

Award [2] for the correct final answer.

(ii)
$$(\Delta S^{\Theta} = [50.9 + 2(34.8)] - [87.5 + 2(28.3)]) = -23.6(JK^{-1}mol^{-1});$$

(iii) $\Delta G^{\Theta} = [-825.2 - (298 \times (-0.0236))];$

$$\Delta G^{\Theta} = -818 \text{ (kJ mol}^{-1}\text{)}/-8.18 \times 10^{5} \text{ (J mol}^{-1}\text{)};$$

Award [2] for the correct final answer.

Award [0] if b(i) is repeated here.

Topic Chem 5 Q# 95/ IB Chem/2015/w/TZ0/P2 Section A/Higher Level/

 (a) C₃H₈(g)+5O₂(g) → 3CO₂(g)+4H₂O(g); Ignore state symbols. [1]

[2]

[2]

[1]

[2]

(b) bonds broken: 2(C-C)/694 + 8(C-H)/3304 + 5(O=O)/2490 / 6488 (kJ);

bonds made: 6(C=O)/4476 + 8(O-H)/3712 / 8188 (kJ);

(6488 – 8188 =) – 1700 (kJ mol⁻¹); Ignore signs in M1 and M2.

Award [3] for the correct final answer.

Award [2] for +1700 (kJ mol-1).

Accept values from 2016 data booklet to give 6494 (kJ) for M1, 8528 (kJ) for M2, and -2034 (kJ) for M3.

Topic Chem 5 Q# 96/ IB Chem/2015/s/TZ2/P2 Section A/Higher Level/

3. (a)
$$(-239.0 - [-110.5] =) -128.5 (kJmol^{-1})$$
;

[1]

[3]

(b)
$$(-166.0 - [-137.2] =) -28.8 (kJmol^{-1});$$

[1]

(C)

$$\left(\Delta G^{\Theta} = -28.8 = -128.5 - \left[\frac{298 \times \Delta S^{\Theta}}{1000}\right]\right)$$

$$(\Delta S^{\oplus} =) - 335 (J \text{ mol}^{-1} \text{ K}^{-1});$$

[1]

(d)
$$\Delta S^{\Theta} = \sum S^{\Theta}_{products} - \sum S^{\Theta}_{reactants} / -335 = 126.8 - 197.6 - 2S^{\Theta}_{H_2};$$

 $S^{\Theta}_{H_1} = (+)132 (Jmol^{-1} K^{-1});$

[2]


Award [2] for correct final answer.

Award [1 max] for $S_{H_h}^{\Theta} = (+)264 (Jmol^{-1} K^{-1})$.



Topic Chem 5 Q# 97/ IB Chem/2015/s/TZ2/P2 Section A/Higher Level/

(a) (i)



drawing best-fit straight lines to show volume;

There should be approximately the same number of points above and below for both lines.

27.0 (cm3);

[2] Accept any value in the range 26.0 to 28.0 (cm3) if consistent with student's

annotation on the graph.

Accept ECF for volumes in the range 27.0-30.0 cm3 if it corresponds to maximum temperature of line drawn.

Volume should be given to one decimal place.

(ii)
$$([HCl]) = \frac{1.00 \times 0.0250}{0.0270}$$

= 0.926 moldm⁻³;

Volume of 26.0 gives [HCl] = 0.962 moldm⁻³. Volume of 28.0 gives [HCl] = 0.893 moldm⁻³

Award [2] for correct final answer with units.

Award [1 max] for correct concentration without units.

Accept M, mol L-1, mol/dm3 as units.



[2]

[1]

[2]

Any accepted value must be consistent with student's annotation on the graph but do not accept $\Delta T < 5.1$.

Accept (+)5.6(°C/K) (ie, taking into account heat loss and using T when volume = 0.0 cm³).

(ii)
$$Q = (m \times c \times \Delta T = (25.0 + 27.0) \times 4.18 \times 5.2 = 1130.272 J =) 1.13 (kJ);$$

 $n = (1.00 \times 0.0250 =) 0.0250 (mol);$

$$\Delta H = (-\frac{Q}{n} = -45210.88 \,\text{Jmol}^{-1} =) -45 \,(\text{kJmol}^{-1});$$
 [3]

Award [3] for correct final answer.

Award [2] for +45 (kJ mol -1).

Apply ECF for M3 even if both m and ΔT are incorrect in M1.

Accept use of $c = 4.2 \text{ Jg}^{-1} \text{K}^{-1}$.

(iii)
$$\left(\left| \frac{-45 - (-58)}{(-58)} \right| \times 100 = \right) 22 (\%);$$
 [1]

Answer must be given to two significant figures. Ignore sign.

(iv) heat losses;

better (thermal) insulation / using a polystyrene cup / putting a lid on the beaker;

Accept other suitable methods for better thermal insulation but do not accept just "use a calorimeter" without reference to insulation.

Topic Chem 5 Q# 98/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(g) atomization of chlorine = $\frac{1}{2}$ bond enthalpy $\frac{1}{2}$ 243 / 121.5 (kJ mol⁻¹);

correct values for ionization Na (+496 kJ mol⁻¹) and electron affinity Cl (-349 kJ mol⁻¹) and lattice enthalpy of NaCl (+790 kJ mol⁻¹ / +769 kJ mol⁻¹);

Born-Haber energy cycle;

Accept lines or arrows in energy cycle.



 $\Delta H_{\star}^{\Theta}(NaCl(s)) = -413.5 / -413 / -414 (kJ mol^{-1});$

Accept -392.5 / -392 / -393 if +769 used for lattice enthalpy. Award [4] for correct final answer.



Topic Chem 5 Q# 99/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

energy needed to break (1 mol of) a bond in the gaseous state/phase; (averaged over) similar compounds; [2] Do not accept "similar bonds" instead of "similar compounds". Concept of "similar" is important for M2. (ii) CH₃CH₂OH+3O₂ → 2CO₂ + 3H₂O₂; Bonds broken: 347 + (5×413) + 358 + 464 + (3×498) / 4728(kJ) / C-C+5C-H+C-O+O-H+3O=O; Bonds made: $(4 \times 746) + (6 \times 464) = 5768 (kJ) / 4C = O + 6O = H;$ $\Delta H = (4728 - 5768 =) - 1040 (kJmol^{-1}) / bonds broken - bonds formed;$ [4] Award [4] for correct final answer. Award [3] for (+)1040 (kJ mol-1). (C) heat loss (to the surroundings); [1] $3C(s) + 3H_2(g) + O_2(g) \rightarrow CH_3CH_2COOH(l)$; (f) (i) $\Delta H_{r}^{\Theta} = \sum \Delta H_{s}^{\Theta} (reactants) - \sum \Delta H_{s}^{\Theta} (products);$ Accept any suitable energy cycle. $\sum \Delta H_{s}^{\Theta}$ (reactants) = $3 \times (-394) + 3 \times (-286) / -2040$ (kJ mol⁻¹); $(\Delta H_{\star}^{\Theta} = [3 \times (-394) + 3 \times (-286)] - (-1527) =) - 513 \text{ (kJ mol}^{-1});$ OR $CH_3CH_2COOH(1) + 3.5O_2(g) \rightarrow 3CO_2(g) + 3H_2O(g);$ $\Delta H_{c}^{\Theta} = \sum \Delta H_{c}^{\Theta} (products) - \sum \Delta H_{c}^{\Theta} (reactants);$ $\sum \Delta H_{\star}^{\Theta}$ (products) = $3 \times (-394) + 3 \times (-286) / -2040$ (kJ mol⁻¹); $(\Delta H_{\star}^{\Theta} = [3 \times (-394) + 3 \times (-286)] - (-1527) =) - 513 \text{ (kJ mol}^{-1});$ [4] Ignore state symbols. Award [4] for correct final answer. (ii) negative; reduction in the number of gaseous molecules; [2] Topic Chem 6 Q# 100/ IB Chem/2019/w/TZ0/P2/Higher Level/Q7./ 6. a iii Proportion of molecules Potential energy $4HCl(g) + O_2(g)$ 2 2Cl₂(g) + 2H₂O(g) Estato Es Kinetic energy Progress of reaction Ea (cat) to the left of Ea V peak lower AND E_{a (cat)} smaller √ a «catalyst provides an» alternative pathway 🗸 «with» lower E_a 2

OR

6.

higher proportion of/more particles with «kinetic» E ≥ E_{a(cat)} «than E_{a»} ✓

Topic Chem 6 Q# 101/ IB Chem/2019/w/TZ0/P2/Higher Level/Q4./

3.	b	i	Rate = $k [C_3H_7Cl] [OH^-] \checkmark$		
			«[OH⁻] held constant and» [C₃H ₇ Ct] triples <i>AND</i> rate triples «so first order wrt C₃H ₇ Ct» \checkmark	3	
			[C₃H₁Ct] doubles AND [OH⁻] doubles AND rate quadruples «so first order wrt OH⁻» ✓		

Topic **Chem 6 Q# 102/** IB Chem/2018/w/TZ 0/P2/Higher Level/Q10

10.	a	B: reactant ✓ D: intermediate ✓	2
10.	b	rate = k[A][B] √	1
10.	С	1.80 «mol dm ⁻³ s ⁻¹ » √	1

Topic Chem 6 Q# 103/ IB Chem/2018/w/TZ 0/P2/Higher Level/

1.	C	i	initial concentration is zero AND concentration increases with time ✓ decreasing gradient as reaction proceeds ✓		2
1.	С	ii	«draw a» tangent to the curve at time = 0 ✓ «rate equals» gradient/slope «of the tangent» ✓	Accept suitable diagram.	2
1.	c	III.	piece has smaller surface area ✓ lower frequency of collisions OR fewer collisions per second/unit time ✓	Accept "chance/probability" instead of "frequency". Do not accept just "fewer collisions".	2



Topic Chem 6 Q# 104/ IB Chem/2018/s/TZ 1/P2/Higher Level/

	a		Any two of:	Do not accept "disappearance of	1
			loss of mass «of reaction mixture/CO₂» ✓	calcium carbonate".	
			«increase in» volume of gas produced √	Do not accept "gas bubbles".	
			MARK MARK CONTRACTOR OF THE CO	Do not accept "colour change" or	2
			change of conductivity ✓	"indicator".	1
			change of pH ✓		
			change in temperature ✓		
	b	i	reaction is fast at high concentration AND may be difficult to measure accurately OR so many bubbles of CO ₂ produced that inhibit contact of HCl (aq) with CaCO ₃ (s) OR insufficient change in conductivity/pH at high concentrations OR calcium carbonate has been used up/is limiting reagent/ there is not enough calcium carbonate «to react with the high concentration of HCl» OR HCl is in excess		1
			so many bubbles of CO₂ produced that inhibit contact of HCl (aq) with CaCO₃ (s) ✓		
t	b	ii	22 20 18 16 19 10 11 12 10 10 10 10 10 10 10 10 10 10 10 10 10		1
b	•	iii	«directly» proportional ✓	Accept "first order" or "linear". Do not accept "rate increases as	1
				concentration increases" or "positive correlation".	
b	0	iv	rate = k [H ⁺] ✓	Accept "rate = k [HCl]".	1
b	0	v	0.02 ✓		
			s ⁻¹ √		2
+	\neg	-	20.5 × 10 ⁻³ «mol dm ⁻³ s ⁻¹ »	Accept any answer in the range	
C	-				1



4.	d	ALTERNATIVE 1: carry out reaction at several temperatures ✓		
		plot ¹ / _T against log rate constant ✓		
		$E_a = -$ gradient $\times R \checkmark$	Accept "gradient = $\frac{-E_*}{R}$ " for M3.	
		ALTERNATIVE 2:		
		carry out reaction at two temperatures ✓	Award both M2 and M3 for the formula $\ln \frac{rate_1}{rate_2} = \frac{E_*}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right).$	3
		determine two rate constants	rate ₂ R (I ₂ I ₁)	
		OR	20 52 60 525 6600 An 150	
		determine the temperature coefficient of the rate \checkmark	Accept any variation of the formula, such as $\frac{\text{rate}_1}{\text{rate}_2} = e^{-\frac{E_1}{R}(\frac{1}{2_1} - \frac{1}{\Gamma_2})}$.	
		use the formula $\ln \frac{k_1}{k_2} = \frac{E_*}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right) \checkmark$	2	

Topic Chem 6 Q# 105/ IB Chem/2017/s/TZ 2/P2/Higher Level/

5.	a	i	Q# 105/ IB Chem/2017/s/TZ 2/P2/Higher Level/ concentration of acid decreases	Accept "less frequency/chance/rate/	1
			OR surface area of magnesium decreases✓	probability/likelihood of collisions". Do not accept just "less acid" or "less magnesium". Do not accept "concentration of reagents decrease".	1
5	а	ii	Time /s curve starting from origin with steeper gradient AND reaching same maximum volume ✓		1
5	b	i	«rate =» k [NO₂]² ✓	Accept rate = k [NO ₂] ² [CO] ⁰ .	1
5	b	ii	«step» I AND CO does not appear in the rate law expression OR «step» I AND only «2 molecules of» NO₂ appears in rate expression ✓	Do not allow ECF from (i).	1
5	b	iii	«IR or UV-vis» spectroscopy OR colorimetry OR colour change «over time» ✓	Accept GC/gas chromatography.	1
5	b	iv	«E _{a(rev)} = 226 + 132 =» 358 «kJ» ✓	Do not accept -358.	1
5	b	v	Rate (*ON)	Curve must go through origin.	1
5	С		activation energy is independent of temperature ✓	Accept "no relationship".	1
5	d		$2NO_2(g) + H_2O(l) \rightarrow HNO_3(aq) + HNO_2(aq)$ OR $4NO_2(g) + 2H_2O(l) + O_2(g) \rightarrow 4HNO_3(aq) \checkmark$	Accept ionized forms of the acids.	1


Topic Chem 6 Q# 106/ IB Chem/2017/s/TZ 1/P2/Higher Level/

15			10	10
1. a	į	use a colorimeter/monitor the change in colour OR take samples AND quench AND titrate «with thiosulfate» ✓	Accept change in pH. Accept change in conductivity. Accept other suitable methods. Method must imply "change".	.1
1 a		0.8 0.7 0.6 0.5 0.7 0.6 0.7 0.7 0.6 0.7 0.7 0.8 0.9 0.9 0.1 0.1 0.1 0.1 0.1 0.1	M2 is independent of M1. Accept range from 0.0070 to 0.0080.	2
1 Ь	i	Relationship: rate of reaction is «directly» proportional to [H ⁺] OR rate of reaction α [H ⁺] ✓ Order of reaction with respect to [H ⁺]: first ✓	Accept "doubling the concentration doubles the rate". Do not accept "rate increases as concentration increases".	2
C		zero order ✓ rate of reaction is the same for all concentrations of iodine ✓	Accept "all graphs have same/similar gradient".	2
d		slow rate of reaction which gradually increases ✓ as H ⁺ ions are produced «to catalyse the reaction» OR reaction is autocatalytic ✓	M1 should mention "rate of reaction".	2

Topic Chem 6 Q# 107/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

-	Question	Answers	Notes	Total
3.	a	H ₂ O AND (I) ✓	Do not accept H ₂ O (aq).	1
3.	b	SO₂(g) is an irritant/causes breathing problems OR SO₂(g) is poisonous/toxic ✓	Accept $SO_2(g)$ is acidic but do not accept "causes acid rain". Accept $SO_2(g)$ is harmful. Accept $SO_2(g)$ has a foul/pungent smell.	1
3.	С	$\begin{split} &n(HCl) = \frac{10.0}{1000} dm^3 \times 2.00 mol dm^{-3} = 0.0200 / 2.00 \times 10^{-2} \text{«mol»} \\ &\textit{AND} \\ &n(Na_2S_2O_3) = \frac{50}{1000} dm^3 \times 0.150 mol \times dm^{-3} = 0.00750 / 7.50 \times 10^{-3} \text{«mol»} \checkmark \end{split}$	Accept answers based on volume of solutions required for complete reaction.	
		0.0200 «mol» > 0.0150 «mol» OR 2.00 × 10 ⁻² «mol» > 2 × 7.50 × 10 ⁻³ «mol» OR $\frac{1}{2}$ × 2.00 × 10 ⁻² «mol» > 7.50 × 10 ⁻³ «mol» ✓	Award [2] for second marking point. Do not award M2 unless factor of 2 (or half) is used.	2





Question	Answers	Notes	Total
. e i	first order ✓ «because» [Na₂S₂O₃] is «directly» proportional to rate of reaction 1 1 1 1	Do not accept "linear" for M2.	2
. e ii	rate = k [Na ₂ S ₂ O ₃][HCI] ✓		1
	45 - 40 - 35 - 22.5 Signal of the state of	Award [2] for correct final answer. Accept value based on candidate's graph. Award M2 as ECF from M1. Award [1 max] for methods involving taking mean of appropriate pairs of 1/t values. Award [0] for taking mean of pairs of time values. Award [2] for answers between 42.4 and 46.4 «s».	2

Question	Answers	Notes	Total
3. g i	Kinetic energy correctly labelled axes peak of T ₂ curve lower AND to the right of T ₁ curve ** ** ** ** ** ** ** ** **	Accept "probability «density» / number of particles / N / fraction" on y-axis. Accept "kinetic E/KE/E _x " but not just "Energy/E" on x-axis.	2

Question		tion	Answers	Notes	Total
3.	g	II	greater proportion of molecules have $E \ge E_a$ or $E > E_a$ OR greater area under curve to the right of the $E_a \checkmark$ greater frequency of collisions «between molecules» OR more collisions per unit time/second \checkmark	Accept more molecules have energy greater than E₂. Do not accept just "particles have greater kinetic energy". Accept "rate/chance/probability/likelihood" instead of "frequency". Accept suitably shaded/annotated diagram. Do not accept just "more collisions".	2
3.	h		shorter reaction time so larger «%» error in timing/seeing when mark disappears ✓	Accept cooling of reaction mixture during course of reaction.	1

Topic Chem 6 Q# 108/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

e	A is indicative of frequency of collisions and probability that collisions have proper orientations ✓	1
f	$k = \left\langle \exp\left[\frac{(-87.0 \times 1000)}{(8.31 \times 298)} + \ln(2.10 \times 10^{11})\right] = \right\rangle 1.2 \times 10^{-4} \checkmark$	2
	S _N 2 implies second-order so mol ⁻¹ dm ³ s ⁻⁴ ✓	

Topic Chem 6 Q# 109/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

3.	a	î	$2NO(g) + 2H_2(g) \rightarrow N_2(g) + 2H_2O(g)$		1
3.	a	ii	rate = k[NO] ² [H ₂] ✓		1
3.	€independently > OR		cindependentlys OR test the effect of varying [NO] con rates, whilst keeping [H ₂] constant AND test effect of varying [H ₂] con rates, whilst keeping [NO] constant ✓ rate proportional to [NO] ² OR	Remember to refer back to a (ii) for ECF.	3
			rate proportional to [H₂] OR doubling [H₂] doubles rate ✓	If only one species in rate expression, third mark can be awarded for zero order discussion.	
3.	a iv no AND different mechanisms could give the same rate expression OR no AND mechanisms can only be disproved OR no AND just suggest it is consistent with the mechanism given OR no AND does not give information about what occurs after RDS ✓		OR no AND mechanisms can only be disproved OR no AND just suggest it is consistent with the mechanism given OR		1
3.	a	٧	change of pressure ∢at constant volume and temperatures with time OR change of volume ∢at constant pressure and temperatures with time ✓	Accept other methods where rate can be monitored with time.	1

Qu	Question		Answers	Notes	Total
3.	b	in .	Second curve at a higher temperature is correctly drawn (maximum lower and to right of original) ✓	31!!	1
3.	b	iv	greater proportion of molecules have E ≥ E, or E > E, OR greater area under curve to the right of the E, ✓ greater frequency of collisions €between molecules OR more collisions per unit time/second ✓	Do not accept just particles have greater kinetic energy. Do not accept just "more collisions".	2



(c) (i) Order with respect to O₂: first (order);Order with respect to NO: second (order);

[2]

(ii) (rate =) $k [NO]^2 [O_2]$;

[1]

(iii) 172.6 / 173; mol⁻² dm⁶ s⁻¹;

[2]

- $\label{eq:accept M-2 s-1/mol-2 L2 s-1} Accept M^{-2} \, s^{-1} / mol^{-2} \, L^2 \, s^{-1}.$ (iv) NO + NO \rightleftharpoons N₂O₂ (fast);
 - $N_2O_2 + O_2 \rightarrow 2NO_2$ (slow); second step is rate determining step;

OR

 $NO + O_2 \rightleftharpoons O + NO_2$ (fast);

O + NO → NO₂ (slow);

second step is rate determining step;

OR

NO + O₂ ⇌NO₃ (fast);

NO₃ + NO → 2NO₂ (slow);

second step is rate determining step;

Accept single arrow instead of reversible sign.

Accept fast/slow next to the equations.

Do not accept the following mechanisms:

$$NO + NO \rightleftharpoons NO_2 + N \text{ (fast)}$$

$$N + O_2 \rightarrow NO_2$$
 (slow)

OR

$$NO + NO + O_2 \rightleftharpoons N_2O_4$$
 (slow)

$$N_2O_4 \rightarrow 2NO_2$$
 (fast)

[3]

Topic Chem 6 Q# 111/ IB Chem/2015/w/TZ0/P2 Section A/Higher Level/

(f) gradient =
$$-\frac{E_a}{R}/-1233$$
 (K);

Accept value from -1200 to -1260 (K).

$$E_a = (1233 \times 8.31 =) 1.02 \times 10^4 \text{ Jmol}^{-1} / 10.2 \text{ kJ mol}^{-1}$$

correct E, value;

correct units;

Allow value in range 9.97 – 10.5 kJ mol-1.

Answer must be given to three significant figures.

Award [3] for correct final answer and units.

Accept J or kJ instead of J mol-1 or kJ mol-1.

[3]



Topic Chem 6 Q# 112/ IB Chem/2015/s/TZ2/P2 Section A/Higher Level/

change in concentration of reactant/product with time / rate of change of concentration;

[1]

Accept "increase" instead of "change" for product and "decrease" instead of "change" for reactant.

Accept "mass/amount/volume" instead of "concentration". Do not accept substance.

- (b) surface area decreases:
 - frequency/probability of collisions decreases; Accept number of collisions per unit time decreases.

[2]

step $2/X + NO \rightarrow Y + H_2O/slow$; (C) (i)

[1]

(ii) invalid / unlikely as order most likely one (with respect to hydrogen); rate = $k[NO]^2[H_2] / H_2$ only involved once in the formation of the intermediate before the slow step / OWTTE; Award M2 only if M1 is correct.

[2]

Topic Chem 6 Q# 113/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

[NaOH] / [OHT] is 1/first order and [C, H, Br] is 1/first order; (C) rate = $k[OH^{-}][C_{\star}H_{\circ}Br] / rate = k[NaOH][C_{\star}H_{\circ}Br]$; Square brackets must be used for M2.

[2]

(ii) $= 0.0016/1.6 \times 10^{-3}$; 0.25×0.25 mol-1 dm3 s-1;

[2]

Accept M-1 s-1. Ignore order of units. Must use experiment 3 data.

(iii) bimolecular/2: Accept dimolecular. [1]



Topic Chem 6 Q# 114/ IB Chem/2015/s/TZ1/P2 Section A/Higher Level/

(a) change in concentration of reactant/product with time / rate of change of concentration;

[1]

Accept "increase" instead of "change" for product and "decrease" instead of "change" for reactant.

Accept "mass/amount/volume" instead of "concentration". Do not accept substance.

(b) collision frequency;

two particles must collide;

particles must have sufficient energy to overcome the activation energy/ $E \ge E_a$;

Concept of activation energy must be mentioned.

appropriate collision geometry/orientation;

[3 max]

(c) (i) increases yield;

(equilibrium shifts to the right/products as) more gaseous moles in reactants/on left / fewer gaseous moles in products/on right;

[2]

(ii) Eqm $[O_2] = 2.6 \text{ (mol dm}^{-3})$;

Eqm[SO₂] = $1.2 \text{(moldm}^{-3})$;

$$K_c = \frac{[SO_3]^2}{[SO_2]^2[O_2]}$$

 $K_c = 0.17$;

[4]

Award [4] for correct final answer. Ignore units.

(iii) (K_c) decreases;

[1]

(d) catalyst increases rate of reaction / equilibrium reached faster / increases yield of product per unit time;

reduces costs / reduces energy needed; Do not accept just "increases the yield". [2]

Topic Chem 7 Q# 115/ IB Chem/2018/w/TZ 0/P2/Higher Level/

	anconon	The state of the s	110163	1000
5.	e	wreaction quotient/Q =» $\frac{[SO_3]^2}{[SO_2]^2[O_2]} I \frac{0.500^2}{0.200^2 \times 0.300} /20.8$ \checkmark	7	
		reaction quotient/Q/20.8/answer < $K_c/280$ OR mixture needs more product for the number to equal K_c		3
		reaction proceeds to the right/products ✓	Do not award M3 without valid reasoning.	

Topic Chem 7 Q# 116/ IB Chem/2018/s/TZ 1/P2/Higher Level/

1.	d	i	$K_{c} = \frac{\left[(H_{2}N)_{2}CO\right] \times \left[H_{2}O\right]}{\left[NH_{3} \right]^{2} \times \left[CO_{2} \right]} \checkmark$	1
1.	d	II	K_{C} decreases AND reaction is exothermic OR K_{C} decreases AND ΔH is negative OR K_{C} decreases AND reverse/endothermic reaction is favoured $$	1



d	iii	In K « = $\frac{-\Delta G^{\Phi}}{RT}$ = $\frac{-50 \times 10^{3} \text{ J}}{8.31 \text{ K}^{-1} \text{ mol}^{-1} \times 298 \text{ K}}$ » = -20 \checkmark	Accept range of 20-20.2 for M1. Award [2] for correct final answer.	
	1 1	«K _c =» 2 × 10 ⁻⁹ OR		2
		1.69 × 10 ⁻⁹		
	1	OR		
		10 ⁻⁹ ✓		

Topic Chem 7 Q# 117/ IB Chem/2017/s/TZ 2/P2/Higher Level/

100			3 3	K.
4 d	i	$\ll Q_c = \frac{0.10}{0.52^2} = \approx 0.37 \checkmark$ reaction proceeds to the left/NO ₂ (g) \ll until $Q = K_c \approx OR$ reverse reaction \ll are reverse reaction \ll are reverse reaction \ll are reverse reaction \ll and \ll are represented as \ll and \ll and \ll are represented as \ll and \ll are represented as \ll and \ll are res	Do not award M2 without a calculation for M1 but remember to apply ECF.	2
4 d	ii ii	ΔG = 0 ✓ reaction at equilibrium OR rate of forward and reverse reaction is the same OR constant macroscopic properties ✓		2

Topic Chem 7 Q# 118/ IB Chem/2017/s/TZ 2/P2/Higher Level/

3.	а	i	$\ll K_{ch} = \frac{[PCl_3][Cl_2]}{[PCl_8]} \checkmark$		1
3	а	ii	decrease in temperature ✓	Do not accept "temperature change".	
			endothermic «reaction» AND «equilibrium» shifts to the left/reactants OR endothermic «reaction» AND Kc decreases OR	Accept "ΔH positive" in place of "endothermic".	2
			endothermic «reaction» AND concentration of PCl ₅ increased/concentration of PCl ₃ and Cl ₂ decreased OR «equilibrium» shifts in the exothermic direction ✓	Accept "products" instead of "PCl $_3$ and Cl $_2$ ".	

Topic Chem 7 Q# 119/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

Question		Answers	Notes	Total
1. a	i	$\ll K_c = n \frac{[HOCH_2CH_2OH]}{[CO]^2 \times [H_2]^3} \checkmark$	1 1 1	1
1. a	ii	Position of equilibrium: moves to right OR favours product ✓ K _c : no change OR is a constant at constant temperature ✓	1:1	2

Topic Chem 7 Q# 120/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

ь				
	Change	Shift	Reason	
	Increase in temperature	LHS	since (forward) exothermic reaction/ $\Delta H < 0$ \(\square\$	
	Increase in pressure	RHS	since fewer (gaseous) molecules on RHS ✓	
	Addition of a catalyst to the mixture	No change	since affects rate of forward and reverse reactions equally 🗸	

Topic Chem 7 Q# 121/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

(Question		Answers	Notes	Total
2.	a	i	«K _c -» [COCl ₂]		1
2.	a	ii	T <-800 + 273> - 873K ✓ ΔG°8.31 × 873 × ln (0.200) OR ΔG° - <+> 11676 <j> ✓ ΔG° - <+> 11.7 < kJ> ✓</j>	Accept 11.5 to 12.0. Award final mark only if correct sig fig. Award [3] for correct final answer.	3

Topic Chem 7 Q# 122/ IB CHEM/2015/w/TZ0/P2 Section B/Higher Level/

(a) (i) Increasing the pressure, at constant temperature:

decreases

more (gas) molecules/moles on the right / fewer (gas) molecules/moles on the left;

Increasing the temperature, at constant pressure:

increases:

(forward) reaction is endothermic;

[4]

(ii) equilibrium reached faster;

no change in the concentration of reactants/products/yield (at equilibrium) / position of equilibrium is not affected;

rates of forward and reverse reactions increase (equally);

reduces activation energy;

no change in Kc;

[3 max]

(iii) $(K_c =) \frac{[H_2]^3 [CO]}{[CH_4][H_2O]}$;

[1]

(iv) (increasing) temperature and (K_c) increases;Award [0] if both temperature and pressure stated.

[1]

(b) $CO(g) + H_2O(g) \rightleftharpoons H_2(g) + CO_2(g)$ (1.00 - x) (1.00 - x) x x

$$K_c = 10.0 = \frac{x^2}{(1.00 - x)^2} / \sqrt{10.0} = \frac{x}{(1.00 - x)};$$

x = 0.760 / (1.00 - x) = 0.240;

 $[CO] = 0.240 \text{ (moldm}^{-3})$ and $[H_2O] = 0.240 \text{ (moldm}^{-3})$ and $[H_2] = 0.760 \text{ (moldm}^{-3})$

and $[CO_2] = 0.760 \text{ (mol dm}^{-3});$

[3]

Topic Chem 7 Q# 123/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(d) (i)
$$(K_c =) \frac{[NH_3(g)]^2}{[N_2(g)] \times [H_2(g)]^3}$$
;

[1]

Ignore state symbols.

Concentrations must be represented by square brackets.

 (ii) moles at equilibrium: nitrogen 0.27, hydrogen 0.81 / concentrations at equilibrium: nitrogen 0.27 (mol dm⁻³), hydrogen 0.81 (mol dm⁻³) (and ammonia 1.46 mol dm⁻³);

$$K_{c} = 15$$
;

[2]

Actual calculation gives $K_c = 14.86$.

Award [2] for correct final answer.

Award [1 max] if
$$K_c \left(= \frac{1.46^2}{3^3 \times 1} \right) = 0.079$$



Topic Chem 7 Q# 124/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(a) rates of forward and reverse reactions are equal / opposing changes occur at equal rates; the concentrations of all reactants and products remain constant / macroscopic properties remain constant; closed/isolated system; [2 max] Accept "the same" for "equal" in M1 and for "constant" in M2. The volume of the container is increased: position of equilibrium shifts to the left/reactants and fewer moles of gas on the right hand side/pressure decreases / OWTTE; Ammonia is removed from the equilibrium mixture: position of equilibrium shifts to the right/products and [NH3] decreases so [N2] and [H2] must also decrease to keep Kc constant position of equilibrium shifts to the right/products and rate of reverse reaction decreases / OWTTE: [2] Award [1 max] if both predicted changes are correct. Do not accept "to increase [NH₃]" or reference to LCP without explanation. (iii) minimum energy needed (by reactants/colliding particles) to react/start/ initiate a reaction; [1] Accept "energy difference between reactants and transition state". (b) rate increases; more effective/successful collisions per unit time / greater proportion of collisions effective:

alternative pathway and a lower activation energy

lowers activation energy so that more particles have enough energy to react;

Do not accept just "lowers/reduces the activation energy".

Accept "provides a surface for reacting/reactants/reaction".

[2 max]

(c) (i) slower rate / OWTTE; uneconomic / OWTTE;

[2]

(ii) high cost for building/maintaining plant / high energy cost of compressor / OWTTE;

[1]

Do not accept "high pressure is expensive" without justification. Accept high pressure requires high energy.



Topic Chem 7 Q# 125/ IB Chem/2015/s/TZ1/P2 Section A/Higher Level/

(c) (i) increases yield;

(equilibrium shifts to the right/products as) more <u>gaseous</u> moles in reactants/on left / fewer <u>gaseous</u> moles in products/on right;

[2]

(ii) Eqm $[O_2] = 2.6 \text{ (mol dm}^{-3});$

Eqm[SO₂] = 1.2(moldm⁻³);

$$K_c = \frac{[SO_3]^2}{[SO_2]^2[O_2]};$$

 $K_c = 0.17$;

[4]

Award [4] for correct final answer. Ignore units.

(iii) (K_c) decreases;

[1]


 (d) catalyst increases rate of reaction / equilibrium reached faster / increases yield of product per unit time;

reduces costs / reduces energy needed; Do not accept just "increases the yield". [2]

Topic Chem 8 Q# 126/ IB Chem/2019/w/TZ0/P2/Higher Level/Q7./

6.	f	iv	Any two of:	
			ligand/chloride ion Lewis base AND donates e-pair ✓	
			not Brønsted–Lowry base AND does not accept proton/H⁺ ✓	2 max
			Lewis definition extends/broader than Brønsted-Lowry definition ✓	

Topic Chem 8 Q# 127/ IB Chem/2019/w/TZ0/P2/Higher Level/Q6./





_		T		
5.	b	ii	ALTERNATIVE 1:	
			H ⁺ (aq) + CH₃COO ⁻ (aq) → CH₃COOH(aq) ✓	
			added acid neutralised by ethanoate ions	
			OR	
			«weak» CH₃COOH (aq)/ethanoic acid replaces H+ (aq)	
			OR	
			CH₃COOH/CH₃COO⁻ ratio virtually/mostly unchanged ✓	2
			ALTERNATIVE 2:	
			$CH_3COOH(aq) \rightleftharpoons H^+(aq) + CH_3COO^-(aq) \checkmark$	
			equilibrium shifts to the ethanoic acid side	
			OR	
			CH₃COOH/CH₃COO⁻ ratio virtually/mostly unchanged ✓	

Topic Chem 8 Q# 128/ IB Chem/2019/w/TZ0/P2/Higher Level/Q5.,

4.	a	i	C ₆ H ₆ O ₇ AND C ₆ H ₇ O ₇ ⁻ OR H ₂ O AND H ₃ O ⁺ ✓		1
4.	а	ii	weak acid <i>AND</i> partially dissociated OR weak acid <i>AND</i> equilibrium lies to left OR weak acid <i>AND</i> K₃ < 1 ✓		1
4.	a	iii	Effect on [H+] Effect on K _s + increases ✓ increases ✓		2
4.	а	iv	« ΔG^0 = − RT In K = −8.31 J K ⁻¹ mol ⁻¹ × 298 K × In(5.01 × 10 ⁻⁴) ÷ 1000 =» 18.8 «kJ mol ⁻¹ » \checkmark		1
4.	a	v	non-spontaneous AND ∆G ^o positive ✓		1
4.	b		Any two of: «electrical» conductivity AND HCl greater ✓ pH AND citric acid higher ✓ titrate with strong base AND pH at equivalence higher for citric acid ✓ add reactive metal/carbonate/hydrogen carbonate AND stronger effervescence/faster reaction with HCl ✓ titration AND volume of alkali for complete neutralisation greater for citric acid ✓	Accept "add universal indicator AND HCI more red/pink" for M2. Accept any acid reaction AND HCI greater rise in temperature. Accept specific examples throughout. Do not accept "smell" or "taste".	2 max
		Ŕ	titrate with strong base AND more than one equivalence point for complete neutralisation of citric acid ✓ titrate with strong base AND buffer zone with citric acid ✓		




Topic Chem 8 Q# 129/ IB Chem/2018/w/TZ 0/P2/Higher Level/

6.	а	i	Butanoic acid: $ CH_3CH_2CH_2COOH(aq) + H_2O(l) \rightleftharpoons CH_3CH_2CH_2COO^-(aq) + H_3O^+(aq) \checkmark $ $ Ethylamine: \\ CH_3CH_2NH_2(aq) + H_2O(l) \rightleftharpoons CH_3CH_2NH_3^+(aq) + OH^-(aq) \checkmark $		2
6.	a	ii	CH ₃ CH ₂ CH ₂ —C:— Diagram showing: dotted line along O–C–O AND negative charge	Accept correct diagrams with pi clouds.	1
6.	a	iii	-1 ✓		1
6.	b	i	$\alpha = \frac{1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}}{0.00192 \text{ mol dm}^{-3}} \text{ w} = 5.21 \times 10^{-12} \text{ emol dm}^{-3} \text{w} \checkmark$		1
	r	_	1	ALOS ATTIMONE PIR STERRE SAMI	f

6.	ь	ii	$\alpha p K_b = 3.35, K_b = 10^{-3.35} = 4.5 \times 10^{-4} \text{w}$	Award [3] for correct final answer.	
			$\alpha C_2 H_5 N H_2 + H_2 O \rightleftharpoons C_2 H_5 N H_3^+ + O H^- w$		
			$K_0 = \frac{[OH^-][CH_3CH_2NH_3^+]}{[OH^-]}$		
			$K_b = \frac{[CH_3CH_3NH_2]}{[CH_3CH_3NH_2]}$		
			OR		
			$\alpha K_b = 3.5 \times 10^{-4} = \frac{[OH^{-}][GH_3CH_3NH_3^{+}]}{0.250}$		
			OR		3
			$\alpha K_b = 34.5 \times 10^{-4} = \frac{x^2}{0.250}$		3
			«x = [OH¬] =» 0.011 «mol dm ⁻³ » √		
			$\alpha pH = -log \frac{1.00 \times 10^{-14}}{0.011} = 12.04$		
			OR		
			«pH = 14.00 − (−log 0.011)=» 12.04 ✓	A Comment	





Topic Chem 8 Q# 130/ IB Chem/2018/s/TZ 1/P2/Higher Level/

5.	a	slower rate with ethanoic acid OR smaller temperature rise with ethanoic acid ✓	Accept experimental observations such as "slower bubbling" or "feels less warm".	
		[H ⁺] lower OR ethanoic acid is weak OR ethanoic acid is partially dissociated ✓		2
5.	b	Any one of: corrosion of materials/metals/carbonate materials ✓ destruction of plant/aquatic life ✓ «indirect» effect on human health ✓	Accept "lowering pH of oceans/lakes/waterways".	1
5.	С	Brønsted-Lowry base: $NH_3 + H^+ \rightarrow NH_4^+ \checkmark$ Lewis base: $NH_3 + BF_3 \rightarrow H_3NBF_3 \checkmark$	Accept "AlCl3 as an example of Lewis acid". Accept other valid equations such as $Cu^{2+} + 4NH_3 \rightarrow [Cu(NH_3)_4]^{2+}$.	2
5.	d	[H ⁺] «= $\sqrt{K_* \times [C_5H_{10}O_2]}$ = $\sqrt{9.333 \times 10^{-6} \times 0.010}$ » = 3.055 × 10 ⁻⁴ «mol dm ⁻³ » \checkmark «pH =» 3.51 \checkmark	Accept "pH = 3.52". Award [2] for correct final answer. Accept other calculation methods.	2



	Š.,				
5.	e		$(CH_3)_3CCOOH\ (aq) + OH^-(aq) \rightarrow (CH_3)_3CCOO^-\ (aq) + H_2O\ (l)$ OR $(CH_3)_3CCOOH\ (aq) + OH^-\ (aq) \rightleftharpoons (CH_3)_3CCOO^-\ (aq) + H_2O\ (l)$ AND addition of alkali causes equilibrium to move to right \checkmark	Accept "HA" for the acid. Award [1 max] for correct explanations of buffering with addition of acid AND base without equilibrium equations.	2
			$(CH_3)_3CCOO^ (aq) + H^+$ $(aq) \rightarrow (CH_3)_3CCOOH$ (aq) OR $(CH_3)_3CCOO^ (aq) + H^+$ $(aq) \rightleftharpoons (CH_3)_3CCOOH$ (aq) AND addition of acid causes equilibrium to move to right \checkmark		2
pi	ic Ch	em 8	Q# 131/ IB Chem/2018/s/TZ 1/P2/Higher Level/		
2.	f		pH > 7 ✓	Accept any specific pH value or range of values above 7 and below 14.	1
ppi	ic Ch	em 8	Q# 132/ IB Chem/2017/s/TZ 2/P2/Higher Level/		
1.	a	i	H ₂ O/water ✓	Accept "hydroxide ion/OH".	1
	а	ii	Acid Base HOCI AND OCIT OR H₂O AND OH™ ✓		1
	b	i	«0.100 mol dm ⁻³ × 0.0250 dm ³ » = 0.00250 «mol» ✓		1
	b	ii	«M = 0.510 g =» 204 «g mol⁻¹» ✓		1
	b	iii	«1.00 × 10 ⁻¹⁴ = [H [†]] × 0.100» 1.00 × 10 ⁻¹³ «mol dm ⁻³ » ✓		1
	b	iv	weak AND pH at equivalence greater than 7 OR weak acid AND forms a buffer region ✓		1
	b	v	calorimetry OR measurement of heat/temperature OR conductivity measurement ✓	Accept "indicator" but not "universal indicator".	1
	b	vi	«pK ₃ = pH at half-equivalence =» 5.0		1
	C		$K_8 = 10^{-4.35} / 4.48683 \times 10^{-5} \checkmark$ $[H_3O^+] = \sqrt{4.46683 \times 10^{-5} \times 1.60 \times 10^{-9}} / \sqrt{7.1469 \times 10^{-9}} / 2.6734 \times 10^{-4} \text{ emol}$ $dm^{-3}\text{ w} \checkmark$ $pH = \text{emolog} \sqrt{7.1469 \times 10^{-8}} = \text{w} 3.57 \checkmark$	Award [3] for correct final answer to two decimal places. If quadratic equation used, then: $[H_3O^+]=2.459 \times 10^{-4}$ «mol dm ⁻³ » and pH = 3.61	3



Topic Chem 8 Q# 133/ IB Chem/2017/s/TZ 1/P2/Higher Level/

5.	а		107° ✓	Accept 100° to <109.5°. Literature value = 105.8°	1
5.	b		tetrahedral ✓ sp³ ✓	No ECF allowed.	2
5.	С		removes/reacts with OH⁻ ✓ moves to the right/products «to replace OH⁻ ions» ✓	Accept ionic equation for M1.	2
5.	d	i	$K_b = 10^{-5.77} / 1.698 \times 10^{-6}$ OR $K_b = \frac{[N_2 H_5^*] \times [OH^-]}{[N_2 H_4]} \checkmark$ $[OH^-]^2 \ll 1.698 \times 10^{-6} \times 0.0100 \approx 1.698 \times 10^{-8}$ OR $[OH^-] \ll \sqrt{1.698 \times 10^{-6}} \approx 1.303 \times 10^{-4} \pmod{dm^{-3}} \checkmark$ $PH \ll -\log_{10} \frac{1 \times 10^{-14}}{1.3 \times 10^{-4}} \approx 10.1 \checkmark$	Award [3] for correct final answer. Give appropriate credit for other methods containing errors that do not yield correct final answer.	3
5.	d	ii	methyl red OR bromocresol green OR bromophenol blue OR methyl orange ✓		1
5.	e		bubbles OR gas OR magnesium disappears ✓ 2NH4+ (aq) + Mg (s) → Mg2+ (aq) + 2NH3 (aq) + H2 (g) ✓	Do not accept "hydrogen" without reference to observed changes. Accept "smell of ammonia". Accept $2H^*$ (aq) $+Mg$ (s) $\to Mg^{2+}$ (aq) +Hz (g) Equation must be ionic.	2
5.	f		bonds broken: E(N-N) + 4E(N-H) OR 158 «kJ moi⁻¹» + 4 × 391 «kJ moi⁻¹» / 1722 «kJ» ✓ bonds formed: E(N≡N) + 2E(H-H) OR 945 «kJ moi⁻¹» + 2 × 438 «kJ moi⁻¹» / 1817 «kJ» ✓ «△H - bonds broken - bonds formed - 1722 - 1817 -» -95 «kJ» ✓	Award [3] for correct final answer. Award [2 max] for +95 «kJ».	3

Topic Chem 8 Q# 134/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

2.	a		Weak acid: partially dissociated/ionized «in solution/water» AND Strong acid: «assumed to be almost» completely/100 % dissociated/ ionized «in solution/water» ✓	Accept answers relating to pH, conductivity, reactivity if solutions of equal concentrations stated.	1
Question		ion	Answers	Notes	Total
2.	b		«log scale» reduces a wide range of numbers to a small range <i>OR</i> simple/easy to use <i>OR</i> converts exponential expressions into a linear scale/simple numbers ✓	Do not accept "easy for calculations".	1
2.	С	i	phenolphthalein OR phenol red ✓		1



Topic Chem 8 Q# 135/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

Q	uestion	Answers	Notes	Total
7.	a	Calculation: ALTERNATIVE 1: $[H^*] = (K_a \times [HA])^{1/2} / (1.6 \times 10^{-4} \times 0.0100)^{1/2} / 1.3 \times 10^{-3} \text{ emol dm}^{-3} \text{ w} \text$	Award [2] for correct final answer. Do not accept partial dissociation.	3
		OR [H'] = [HCOO ⁻] ✓	7 - 20	

Question	Answers	Notes	Total
b i	pH 7 — Volume V _n / at neutralization Volume of strong base added correct shape of graph ✓ pH at half neutralization/equivalence ✓	M1: must show buffer region at pH < 7 and equivalence at pH > 7. Accept graph starting from where two axes meet as pH scale is not specified.	2
b ii	ALTERNATIVE 1: HCOOH ⇒ HCOO⁻ + H⁺ ✓ H⁺ ions consumed in reaction with OH⁻ are produced again as equilibrium moves to the right «so [H⁺] remains almost unchanged» ✓ ALTERNATIVE 2: HCOOH + OH⁻ ⇒ HCOO⁻ + H₂O ✓ added OH⁻ are neutralized by HCOOH OR strong base replaced by weak base ✓	Accept HA or any other weak acid in equations. Equilibrium sign must be included in equation for M1.	2

Topic Chem 8 Q# 136/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

(Question	Answer	Notes	Total
2.	а	radical / unpaired electron ✓	MAN A STATE OF THE	1
(Question	Answers	Notes	Total
	d	$2NO_2(g) + H_2O(l) \rightarrow HNO_3(aq) + HNO_2(aq)$	Ignore state symbols.	1

Topic Chem 8 Q# 137/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

а	$HCOO^{-}(aq) + H_2O(l) \rightleftharpoons OH^{-}(aq) + HCOOH(aq)$	Equilibrium sign must be given for mark.	1
ь	K _a = 1.8×10 ⁻⁴ ✓	114 11 114 114 114	1
c	$K_b = \left(\frac{K_w}{K_k} = \frac{1.0 \times 10^{-14}}{1.8 \times 10^{-4}} = \right) 5.6 \times 10^{-11} \checkmark$	MGJII	1
d	$K_b = \frac{x^2}{0.12} = 5.6 \times 10^{-11}$ $[OH^-(aq)] = 2.6 \times 10^{-6} \langle moldm^{-2} \rangle$ Assumption: $0.12 - x \sim 0.12$	Award [2] for correct final answer of [OH (aq)]. Accept any other reasonable assumption.	3
e	pOH = <-log(2.6×10 ⁻⁶) =>5.59 ✓ pH = (14.00 - 5.59 =>8.41 ✓	Award [2] for correct final answer.	2



Topic Chem 8 Q# 138/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

	1 -	1		•	
2.	d		$K_b \approx \frac{[OH^-]^2}{[C_eH_bNH_b]} = 10^{-9.13} / 7.413 \times 10^{-10} \checkmark$ $[OH^-] = \sqrt{0.0100 \times 10^{-9.13}} = 2.72 \times 10^{-6} \checkmark$ $[H^+] = \frac{1 \times 10^{-14}}{2.72 \times 10^{-6}} = 3.67 \times 10^{-9}$ OR $pOH = 5.57 \checkmark$	Accept other approaches to the calculation. Award [4] for correct final answer. Accept any answer from 8.4 to 8.5.	4
		1	pH = -log[H*] = 8.44 ✓		

Topic Chem 8 Q# 139/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/Q1(a)

1.	а	iii	Lewis base AND has a lone pair of electrons €to donate> ✓		1
1.	а	vi	Weak: only partially dissociated/ionized €in dilute aqueous solutions ✓ Brønsted–Lowry base: an acceptor of protons/H*/hydrogen ions ✓	Accept reaction with water is reversible/an equilibrium. Accept "water is partially dissociated @by the weak bases".	2
1.	b	i	P ₄ is a molecule comprising 4P atoms AND 4P is four/separate P atoms OR P ₄ represents 4P atoms bonded together AND 4P represents 49 separate/non-bonded P atoms		1

Questio	Answers	Notes	Total
1. b	ii can act as both a ∉Brønsted–Lowrys acid and a ∢Brønsted–Lowrys base OR can accept and/or donate a hydrogen ion/proton/H" ✓ HPO,2- AND H,PO,✓		2

Topic Chem 8 Q# 140/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

4.	a	iii	hydrochloric acid shifts equilibrium to left	Accept "hydrochloric acid reacts with «basic»		1
-135	1	100	OR	chloride" OR "hydrochloric acid suppresses	1	-
1			hydrochloric acid prevents the basic chloride forming/precipitating ✓	salt hydrolysis".		

Topic Chem 8 Q# 141/ IB CHEM/2015/w/TZ0/P2 Section B/Higher Level/

(d) (i) Pb < Ni < Fe < Zn;;

[2]

Award [2] for the correct order.

Award [1] for Zn > Fe > Ni > Pb as metals not listed in order of increasing reactivity.

Award [1] if one error in the order.

(ii) Pb²⁺/ lead(II) (ions);Do not accept Pb/lead.

[1]

(e) Dilute sodium chloride:

$${\rm H_2O}({\rm I}) \rightarrow \frac{1}{2}{\rm O_2}({\rm g}) + 2{\rm H^+}({\rm aq}) + 2{\rm e^-}/2{\rm OH^-}({\rm aq}) \rightarrow \frac{1}{2}{\rm O_2}({\rm g}) + {\rm H_2O}({\rm I}) + 2{\rm e^-};$$

Concentrated sodium chloride:

$$Cl^{-}(aq) \rightarrow \frac{1}{2}Cl_{2}(g) + e^{-};$$
 [2]

Award [1 max] if equations are given the wrong way round.

Accept e instead of e-.

Ignore state symbols and equilibrium sign.

Accept correct half-equations balanced with different coefficients.



Topic Chem 8 Q# 142/ IB CHEM/2015/w/TZ0/P2 Section B/Higher Level/

8. (a) (i) base 1: 7.0;

base II: 5.5;

[2]

(ii) base I: strong base and base II: weak base;

[1]

[1]

(iii) NaOH and KOH;

Accept LiOH. (iv) (from the graph: initial pH = 11, pOH = 3) $[OH^{-}] = 10^{-3} (mol dm^{-3})$:

$$K_{b} = \frac{10^{-3} \times 10^{-3}}{0.100 - x} / \frac{10^{-3} \times 10^{-3}}{0.100};$$

$$K_{b} = 1.0 \times 10^{-5};$$

OR

at half-equivalence point, $pH = 9.0 = pK_a$ for base II;

$$pK_n = 14.0 - 9.0 = 5.0$$
;

pH range, 3.8 is too low).

Award M1 and M2 for stating: "at half-equivalence point $pOH = 5 = pK_b$ for base II".

$$K_{\rm b} = 1.0 \times 10^{-5}$$
; [3]

Accept other valid methods of working.

Accept a slightly different value of pK_b and so K_b if the student uses the pH at the end point, rather than the initial or half-neutralisation values.

- (v) methyl red;
 Do not accept bromocresol green (since from table 16 data booklet
- (b) (i) resists change in pH; on addition of small amounts of acid and base; [2]
 - (ii) (after mixing) $[NH_3] = [NH_4^+] / nNH_3 = nNH_4^+ / n = 4.00 \times 10^{-3} \text{ mol};$

$$[OH^{-}] = K_b \frac{[base]}{[salt]}$$
 / pOH = pK_b = 4.75;

 $p_{\Pi} = 9.23,$ $A_{\text{total}} = 66 \text{ mag s}^{-1} \text{ if an analytic substantial}$

Award [1 max] if no working shown.
Accept other valid methods of working.

Reaction A: base and accepts a proton/H+;

Reaction B: acid and donates a proton/H+;

Accept donates a pair of electrons.

[2]

[3]

Award [1] if acid and base identified correctly without a reason.

	Acid		Base
Conjugate acid-base pair 1/2	H ₂ CO ₃	and	HCO₃ ⁻
Conjugate acid-base pair 2/1	HCO ₃ -	and	CO ₃ ²⁻ ,
Conjugate acid-base pair 1/2	H ₃ O ⁺	and	H ₂ O;

[2 max]



(C)

(i)

(d) (i) strong acid: (assumed to be) completely/100% dissociated/ionized and weak acid: partially dissociated/ionized;

[1]

 (ii) Similarity: bubbling/effervescence/gas / heat/increase in temperature / solid dissolves;
 Difference: strong acid more vigorous / faster reaction / greater temperature

[2]

increase;
Accept converse statements for weak acid.

(iii) 10⁴(: 1) / 10⁻¹:10⁻⁵ / 1:10⁻⁴; Do not accept inverse ratio, 1:10⁴. [1]

Topic Chem 8 Q# 143/ IB Chem/2015/w/TZ0/P2 Section A/Higher Level/

(b) (NH₃) Lewis base and (Cu²⁺) Lewis acid;

NH₃/ligand donates an electron pair (to Cu²⁺) / Cu²⁺ accepts an electron pair (from NH₃);

[3]

forming coordinate/dative (covalent) bond;

Accept coordination bond.


Do not accept just covalent bond.

Topic Chem 8 Q# 144/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(e) (i) electron pair donor, Accept lone pair donor. [1]

(ii) proton acceptor and partially/slightly ionized;Accept "proton acceptor and partially/slightly dissociated".

[1]



[2]

Award [1 max] for two correct acids OR two correct conjugate bases.

(f)
$$K_b = \frac{[NH_4^+][OH^-]}{[NH_3]} = 1.8 \times 10^{-5} / 10^{-4.75};$$

 $[NH_4^+] = [OH^-]$ and $[NH_3] \approx 1.00 \times 10^{-1} (mol \ dm^{-3})$;


$$[OH^{-}] = (\sqrt{1.8 \times 10^{-6}}) = (1.3 \times 10^{-3}) / pOH = 2.89;$$

 $pH = (14.0 - 2.89) = (11.1);$

[4]

Award [4] for correct final answer.





Ignore state symbols.

Accept reversible arrows.

addition of base:

Award [1 max] if correct equations are given without reference to addition of acid or alkali.

 $CH_3CH_3COOH(aq) + OH^-(aq) \rightarrow CH_3CH_3COO^-(aq) + H_3O(l) / addition of OH^-$

removes H⁺ and more propanoic acid dissociates/ionizes;



[2]

(c)
$$K_a = \frac{[H^+(aq)][CH_3CH_2COO^-(aq)]}{[CH_3CH_2COOH(aq)]} / pH = pK_a + log(\frac{[base]}{[acid]});$$

 $K_a = 1.3 \times 10^{-5} / 10^{-4.87}$ and $[H^+] = 1.3 \times 10^{-4} / 10^{-3.87} (moldm^{-3}) / log(\frac{[CH_3CH_2COO^-]}{[CH_3CH_2COOH]} = 3.87 - 4.87 = -1;$
 $([CH_3CH_2COOH] = \frac{7.41}{74.09} =)0.100 / 1.00 \times 10^{-1} (moldm^{-3});$
 $([CH_3CH_2COONa] =)0.010 / 1.0 \times 10^{-2} (moldm^{-3});$

[4]

Award [4] for correct final answer.

Accept corresponding use of [H₃O]⁺ for [H⁺], [acid] for [CH₃CH₂COOH], and [base] or [salt] for [CH₃CH₂COO⁻] throughout.

Topic Chem 8 Q# 146/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

 (a) (i) weak acids dissociate only partially and strong acids (are assumed to) dissociate fully;

[1]

 (ii) measuring electrical conductivity and strong acids have greater electrical conductivity/weak acids have lower electrical conductivity;
 Do not accept conductivity for electrical conductivity.
 Accept explanation in terms of lightbulb in circuit.

measure pH/use universal indicator and pH higher for weak acid/pH lower for strong acid;

conduct titration with a strong base **and** equivalence point higher for weak acid / buffer region for weak acid;

adding a reactive metal/carbonate/hydrogen carbonate and stronger effervescence/faster reaction with strong acids;

Accept converse argument.

Accept correct example.

adding a strong base and strong acid would increase more in temperature/ weak acids increase less in temperature;

Accept correct example.

[3 max]

Award [1 max] for three suitable tests without correct results.

Accept specific examples with given strong acid and weak acid.

Accept "addition of AgNO₃ (aq) and white precipitate with HCl (aq)".

Do not accept "smell".



(b)	$\frac{1.40}{60.06} = 0.0233 \text{ (mol)}$ and $\frac{0.0233}{0.500} = 0.0466 \text{ (moldm}^{-3}\text{)};$	
	$(pK_a = 4.76)K_a = 1.7 \times 10^{-5}$;	
	$[H^+] = \sqrt{K_a[HA]} = 8.9 \times 10^{-4}$;	
	Accept 9.0×10-4.	
	pH = 3.05;	[4]
	Award [4] for correct final answer.	
	Accept alternative methods.	

(c) (i) Equivalence point: pH of 9.5;Accept values between 9 and 10.

$$pK_a = pH$$
 at half equivalence point;
 $pK_a = 5.4$; [3]
Accept any value between 5.2 and 5.6.
Award [2] for M2 and M3 if correct pK_a given without explanation.

- (ii) phenolphthalein; [1]
- (iii) HIn (aq) ⇒ H+ (aq) + In- (aq) and HIn and In- have different colours; Ignore state symbols.
 equilibrium shifts depending on addition of H+ and OH-/ more HIn in acid/low pH / more In- in alkali/high pH;
 [2]
- (d) (i) CH₂ClCOO-; [1]
 - (ii) stronger because pK_a of chloroethanoic acid is $< pK_a$ of ethanoic acid; [1]
 - (iii) Concentration of acid: $\frac{0.030}{0.300} = 0.10 \text{ (mol dm}^{-3}\text{)};$ Concentration of base/salt: $\frac{0.020}{0.300} = 0.067 \text{ (mol dm}^{-3}\text{)};$ $[H^{+}] = \frac{K_{a} \times [HA]}{[A^{-}]} / \frac{1.3 \times 10^{-3} \times 0.10}{0.067} / 1.9 \times 10^{-3} \text{ (mol dm}^{-3}\text{)};$

pH = 2.72;

Award [4] for correct final answer.

Accept 2.69, 2.70 or 2.7.

Alternative for M3 and M4 if Henderson-Hasselbalch equation used:

M3:
$$pH = pKa + log \frac{[base]}{[acid]} / 2.87 + log \left(\frac{0.067}{0.10}\right)$$

M4: $pH = 2.70$.

Award [1 max] for $n_{\rm acid}$ (= $100 \times 0.50 + 1000$) = $0.050 \, \rm mol$ and $n_{\rm base}$ (= $200 \times 0.10 + 1000$) = $0.020 \, \rm mol$.



[4]

Topic Chem 8 Q# 147/ IB Chem/2015/s/TZ1/P2 Section A/Higher Level/

(c) Ammonium chloride:

Accept any value in the range: 3 < pH < 7;

 $NH_4^+(aq) \rightleftharpoons NH_3(aq) + H^+(aq)$;

Sodium methanoate:

7 < pH < 11;

 $HCOO^{-}(aq) + H_2O(I) \rightleftharpoons HCOOH(aq) + OH^{-}(aq);$

[4]

Award [1 max] for both M1 and M3 combined if stated "pH < 7/acidic for ammonium chloride and pH > 7/alkaline/basic for sodium methanoate".

Accept alternative suitable equations.

Award [1 max] for two correct explanations, such as "salt of weak acid and strong base" or "salt of weak base and strong acid", without equations. Penalize missing equilibrium sign once only.

Ignore state symbols.

Topic Chem 9 Q# 148/ IB Chem/2019/w/TZ0/P2/Higher Level/Q7./

6.	С	i	Wires: «delocalized» electrons «flow» ✓		2
			Electrolyte: «mobile» ions «flow» ✓		
6.	С	ii	$2Cl^- \rightarrow Cl_2(g) + 2e^ OR$ $Cl^- \rightarrow \frac{1}{2}Cl_2(g) + e^- \checkmark$	Accept e for e⁻.	1
6.	С	iii	«electrode» 3 AND oxygen/O₂ ✓	Accept chlorine/Cl ₂ .	1
6.	С	iv	$2H_2O(l) \rightarrow 4H^+(aq) + O_2(g) + 4e^-\checkmark$	Accept 2Cl ⁻ (aq) \rightarrow Cl ₂ (g) + 2e ⁻ . Accept 4OH ⁻ \rightarrow 2H ₂ O + O ₂ + 4e ⁻	1
6.	е	i	E ⁰ = «+0.52 − 0.15 = +» 0.37 «V» ✓		1
6.	е	ii	spontaneous AND E ^o positive ✓	1	1
6.	е	iii	$\Delta G^0 = \text{$\sim$-nFE} = -1 \text{ mol } \times 96500 \text{ C Mol}^{-1} \times 0.37 \text{ \vee=} \text{\sim} -36000 \text{ J/} -36 \text{ kJ} \checkmark$	Accept "-18 kJ mol-1 «per mole of Cu+»." Do not accept values of n other than 1. Apply SF in this question. Accept J/kJ or J mol-1/kJ mol-1 for units.	1
6.	е	iv	2 mol (aq) → 1 mol (aq) AND decreases ✓	Accept "solid formed from aqueous solution AND decreases". Do not accept 2 mol → 1 mol without (aq).	1
6.	е	V	$\Delta G^{\circ} < 0$ AND $\Delta S^{\circ} < 0$ AND $\Delta H^{\circ} < 0$ OR $\Delta G^{\circ} + T\Delta S^{\circ} < 0$ AND $\Delta H^{\circ} < 0$ \checkmark		1
6.	е	vi	T∆S more negative «reducing spontaneity» AND stability increases ✓	Accept calculation showing non-spontaneity at 433 K.	1



Topic Chem 9 Q# 149/ IB Chem/2019/w/TZ0/P2/Higher Level/Q3./

10.74	100	*	F AND 1047 15 21 21 21 22 PAGE 105 AND 105	842	
2.	a	i	4:1 ✓		1
2.	a	ii	$n_{8_20_3^{9-}} = \text{ $<0.0258 dm}^3 \times 0.010 \text{ mol dm}^{-3} = \text{ $>2.58 \times 10^{-4} \text{ $<$mol $>$}$} $ $ \text{ $<\frac{2.58 \times 10^{-4} \text{ mol}}{4} = \text{ $>6.45 \times 10^{-5} \text{ $<$mol $>$}$} $	Award [2] for correct final answer.	2
2.	а	iii	«difference in moles per dm³ = $(6.45 \times 10^{-5} - 5.03 \times 10^{-5}) \times \frac{1000}{300.0}$ =» 4.73×10^{-5} «mol dm⁻³» ✓ «convert to mg per dm³: 4.73×10^{-5} mol dm⁻³ × 32.00 g mol⁻¹ × 1000 mg g⁻¹ = » 1.51 «ppm/mg dm⁻³» ✓	Award [2] for correct final answer.	2
		+			+

Topic Chem 9 Q# 150/ IB Chem/2018/w/TZ 0/P2/Higher Level/

- 1				,, , , , , , , , , , , , , , , , , , , ,		
	6.	a	iii	-1 ✓	1	

Topic Chem 9 Q# 151/ IB Chem/2018/w/TZ 0/P2/Higher Level/

3.	d	i	$BrO_3^-(aq) + 6e^- + 6H^+(aq) \rightarrow Br^-(aq) + 3H_2O(l)$ correct reactants and products \checkmark balanced equation \checkmark	Accept reversible arrows.	2
3.	d	ii	$BrO_3^-(aq) + 8Fe^{2^+}(aq) + 6H^+(aq) \rightarrow Br^-(aq) + 3H_2O(l) + 6Fe^{3^+}(aq)$		1
3.	d	iii	$E^{\circ}_{\text{reaction}} = $		2

Topic Chem 9 Q# 152/ IB Chem/2018/w/TZ 0/P2/Higher Level/Q1

obi	CCnem	19 Q# 152/ IB Chem/2018/w/12 U/P2/Higher Level/Q1	inou.	1 USMI
1.	d	Anode (positive electrode): $2H_2O(l) \rightarrow O_2(g) + 4H^+(aq) + 4e^- \checkmark$	Accept " $4OH^{-}(aq) \rightarrow O_{2}(g) + 2H_{2}O(l) + 4e^{-s}$ $OR "Fe^{2s}(aq) \rightarrow Fe^{3s}(aq) + e^{-s} \text{ for } M1.$	
		Cathode (negative electrode);		
		$2H_2O(l) + 2e^- \rightarrow H_2(g) + 2OH^-(aq)$	Accept "Fe2+ (aq) $+2e^- \rightarrow Fe$ (s)" OR	
		OR	" $SO_4^{2-}(aq) + 4H^+(aq) + 2e^- \rightarrow 2H_2SO_3(aq) + H_2O(l)$ "	2
		$2H^+(aq) + 2e^- \rightarrow H_2(g) \checkmark$	for M2.	2
		VALL V	10	
		SMACHIN	(3111	



Tonic Chem 9 O# 153/ IB Chem/2018/s/T7 1/P2/Higher Level/

OP	ar saon	1 9 Q# 153/ IB Chem/2018/s/TZ 1/P2/Higher Level/	HOLLS	1 Other
3.	a	salt bridge ✓ movement of ions OR	Do not accept "to complete circuit" unless ion movement is mentioned for	2
S.	b	balance charge ✓ Positive electrode (cathode):	M2. Award [1 max] if correct equations given at wrong electrodes.	
		$Ag^{+}(aq) + e^{-} \rightarrow Ag(s) \checkmark$ Negative electrode (anode): $Mg(s) \rightarrow Mg^{2+}(aq) + 2e^{-} \checkmark$		2
	С	in external wire from left to right ✓		1
	d	«E = +0.80 V − (-2.37 V) = +» 3.17 «V» ✓		1
	е	«moles of silver = $\frac{0.10 \text{ g}}{107.87 \text{ g mol}^{-1}}$ » moles of magnesium = $\frac{0.5 \times 0.10 \text{ «g»}}{107.87 \text{ «g mol}^{-1}}$ ✓ «loss in mass of magnesium = $\frac{24.31 \text{ g mol} \times 0.5 \times 0.10 \text{ g}}{107.87 \text{ g mol}^{-1}}$ = » 0.011 «g» ✓	Award [2] for correct final answer.	2

4 3	a	iii	N ₂ H ₄ : −2 AND N ₂ O ₄ : +4 ✓		1
4 2	a	îv	N₂H₄ AND oxidized/oxidation state increases OR N₂H₄ AND loses hydrogen OR N₂H₄ AND reduces/removes oxygen from N₂O₄ ✓	Accept "N ₂ H ₄ AND gives electrons «to N ₂ O ₄ »".	1

Topic Chem 9 Q# 155/ IB Chem/2017/s/TZ 2/P2/Higher Level/

2.	a	i	$\operatorname{Sn^{2+}}(\operatorname{aq}) \to \operatorname{Sn^{4+}}(\operatorname{aq}) + 2\operatorname{e}^{-} \checkmark$	Accept equilibrium sign. Accept $Sn^{2+}(aq) = 2e \rightarrow Sn^{4+}(aq)$	1
2	b	i	«E [©] =» +0.85 «V» ✓	Accept 0.85 V.	1
2	b	ii	ΔG° «= $-nFE^{\circ}$ » = -2 «mol e ⁻ » × 96500 «C mol ⁻ !» × 0.85 «V» \checkmark « ΔG° =» -164 «kJ» \checkmark	Accept "«+»164 «kJ»" as question states energy released. Award [1 max] for "+" or "-" 82 «kJ». Do not accept answer in J.	2
2	c		Zn/zinc is a stronger reducing agent than Fe/iron OR Zn/zinc is oxidized instead of Fe/iron OR Zn/zinc is the sacrificial anode ✓	Accept "Zn is more reactive than Fe". Accept "Zn oxide layer limits further corrosion". Do not accept "Zn layer limits further corrosion".	1.

Topic Chem 9 Q# 156/ IB Chem/2017/s/TZ 1/P2/Higher Level/

		PAGE 1007 TS 14 MINE 10 10 VOIC TS		
4.	a	2NiS (s) + 3O ₂ (g) → 2NiO (s) + 2SO ₂ (g) ✓		1
4.	b	formation of «gaseous» pentacarbonyliron is slower OR «gaseous» complexes form at different rates OR gases have different rates of diffusion «due to difference in masses» OR difference in thermal stability of «gaseous» complexes OR difference in boiling points of «gaseous» complexes OR difference in solubility of «gaseous» complexes OR difference in surface affinity «onto solid absorbent» OR difference in chemical properties of «gaseous» complexes ✓	Accept any other valid answer.	1

Topic Chem 9 Q# 157/ IB Chem/2017/s/TZ 1/P2/Higher Level/

3.	a		V ₂ O ₆ +5 ✓ VO ²⁺ : +4 ✓	Do not penalize incorrect notation twice.	2
3.	b	i	H ₂ SO ₃ (aq) OR Pb (s) ✓		1
3.	b	ii	Zn (s) √		1
3.	С	i	$VO^{2+}(aq) + V^{2+}(aq) + 2H^{+}(aq) \rightarrow 2V^{3+}(aq) + H_2O(l) \checkmark$	Accept equilibrium sign.	1
3.	c	ii	E° «= +0.34 V = (-0.26 V)» = +0.60 «V» \checkmark ΔG° = «-nFE $^{\circ}$ = -9.65 × 10 ⁴ C mol ⁻¹ × 0.60 J C ⁻¹ =» = 57 900 «J mol ⁻¹ » / -57.9 «kJ mol ⁻¹ » \checkmark spontaneous as ΔG° is negative \checkmark	Do not award M3 as a stand-alone answer. Accept "spontaneous" for M3 if answer given for M2 is negative. Accept "spontaneous as E " is positive" for M3.	3

Topic Chem 9 Q# 158/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

4.	•	iii	Anode (positive electrode): oxygen/O₂ OR hydogen ion/proton/H* AND oxygen/O₂ ✓ Cathode (negative electrode): hydrogen/H₂ OR hydroxide «ion»/OH* AND hydrogen/H₃ ✓	Award [1 max] if correct products given at wrong electrodes.	2
4.	j		Any two of: «inert» Pt electrode OR platinum black conductor ✓ 1 mol dm ⁻³ H ⁺ (aq) ✓ H ₂ (g) at 100 kPa ✓	Accept 1 atm $H_2(g)$. Accept 1 bar $H_2(g)$ Accept a labelled diagram. Ignore temperature if it is specified.	2 max
4.	k	i	$Mg(s) + Cu^{2+}(aq) \rightarrow Mg^{2+}(aq) + Cu(s) \checkmark$		1
4.	k	ii	α+0.34V - (-2.37V) = +»2.71 αV» ✓		1
4.	k	III	cell potential increases ✓ reaction «in Q4(k)(i)» moves to the right OR potential of the copper half-cell increases/becomes more positive ✓	Accept correct answers based on the Nernst equation.	2

Topic Chem 9 Q# 159/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

4.	k	i	$Mg(s) + Cu^{2+}(aq) \rightarrow Mg^{2+}(aq) + Cu(s) \checkmark$		1
4.	k	ii	«+0.34V - (-2.37V) = +»2.71 «V» ✓		1
4.	k	III	cell potential increases ✓	Accept correct answers based on the Nernst equation.	
			reaction «in Q4(k)(i)» moves to the right OR	100	2
			potential of the copper half-cell increases/becomes more positive ✓		

Topic Chem 9 Q# 160/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

1.	С	Ethene: -2 ✓	Do not accept 2-, 1- respectively.	2
		Ethane-1,2-diol: -1 ✓		2



Topic Chem 9 Q# 161/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

ь	i	NaClO: +1 (for chlorine) and I₂: 0 (for iodine) ✓		1
ь	ii	CIO since chlorine reduced/gains electrons OR CIO since oxidation state of chlorine changes from +1 to -1/decreases OR CIO since it loses oxygen / causes iodide to be oxidized ✓		1
ь	iii	produces chlorine ⟨gas⟩/Cl₂ ⟨on reaction with ClO⁻⟩ which is toxic ✓	OWITE	1
b	iv	oxidation states are not real OR oxidation states are just used for electron book-keeping purposes ✓ average oxidation state of sulfur calculated to be +2 ✓ but the two sulfurs are bonded differently/in different environments in thiosulfate so have different oxidation states ✓	OWTIE	2 max

Question	Answers	Notes	Total
c	Valid: addition of oxygen signifies an oxidation reaction so C is oxidized OR loss of hydrogen signifies an oxidation reaction so C is oxidized OR oxidation state of C changes from −4 to +4/increases ✓ Not valid: loss of electrons might suggest formation of ionic product but not valid since CO₂ is covalent OR loss of electrons might suggest formation of ionic product but not valid since reaction only involves neutral molecules ✓	OWTTE	2

Topic Chem 9 Q# 162/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

4.	ь	1	E° <- 0.34 - 0.15> = 0.19 <\> ✓ ΔG° < nFE° = -2 × 96500 × 0.19> 36670 / -37000 <j> / -37 <kj> ✓</kj></j>	Accept −18335 € Js or −18 €kJs as equation not specified.	2
4.	b	H	yes AND ∆G ^o is negative OR yes AND E ^o for the cell is positive OR yes AND Sn ²⁺ (aq) is a stronger reducing agent than V ²⁺ (aq) OR yes AND E ^o for Sn ⁴⁺ (aq) is more negative than E ^o for VO ²⁺ (aq) OR yes AND VO ²⁺ (aq) is a stronger oxidizing agent than Sn ⁴⁺ (aq) OR yes AND E ^o for VO ²⁺ (aq) is more positive than E ^o for Sn ⁴⁺ (aq) ✓	Do not accept reference to anti-clockwise rule.	1

Topic Chem 9 Q# 163/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

1.	ь	iii	P ₄ : H ₂ PO ₂ ::	0 +1	: V		Do not accept 1 or 1+ for H ₂ PO ₂ ⁻ .	2
1.	b	o iv	iv oxygen gained, so could be oxidation ✓ hydrogen gained, so could be reduction OR negative charge €on product/H,PO, **p/gain of electrons, so could be	Award [1 max] for M1 and M2 if candidate displays knowledge of at least two of these definitions but does not apply them to the reaction.	3			
			reduction ✓ oxidation number increases so must be oxidation ✓	Do not award M3 for "oxidation number changes".				

Topic Chem 9 Q# 164/ IB Chem/2015/w/TZ0/P2 Section A/Higher Level/

 IO_3^- to I_2 : V/+5 to 0; (a) (i)

$$I^-$$
 to $I_2:-1/-1$ to 0;

[2]

Accept change in oxidation number -5 and +1. Penalize incorrect notation such as 5+ or 5 once only.

(ii) Oxidizing agent: IO₃/iodate and Reducing agent. I⁻/iodide;

[1]



Topic Chem 9 Q# 165/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(d) (i) Half equation:

$$I^{-}(aq) + 3H_{2}O(l) \rightarrow IO_{3}^{-}(aq) + 6H^{+}(aq) + 6e^{-};$$

Accept e instead of e.

Accept reversible arrows.

Overall equation:

$$Cr_2O_7^{2-}(aq) + I^-(aq) + 8H^+(aq) \rightarrow 2Cr^{3+}(aq) + IO_3^-(aq) + 4H_2O(l);$$
 [2]

Ignore state symbols.

oxidized and increase (in oxidation number) of 6/from -1/-I to +5/+V; [1]

Topic Chem 9 Q# 166/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

potential (of reduction half-reaction) under standard conditions measured (e) relative to standard hydrogen electrode/SHE / OWTTE; [1] Accept "solute concentration of 1 mol dm⁻³" or "1 bar/1 atm (pressure) for gases" instead of "standard conditions".

Accept voltage/emf for potential.

- (ii) (+)1.08 (V); [1]
- (iii) $2Cr(s) + 3Cu^{2+}(aq) \rightarrow 2Cr^{3+}(aq) + 3Cu(s)$; [1] Ignore state symbols. Do not accept reversible arrows.
- (iv) Negative electrode: chromium/Cr and Positive electrode: copper/Cu; [1] Accept "Cr is the anode and Cu the cathode".
- from chromium/Cr to copper/Cu; [1] Accept "from negative electrode/anode to positive electrode/cathode" if electrodes correctly identified in (iv).
- (vi) negative ions/anions towards the chromium(III) solution and positive ions/cations towards the copper(II) solution / OWTTE; [1]

Topic Chem 9 Q# 167/ IB Chem/2015/s/TZ2/P2 Section A/Higher Level/

6. (a) Positive electrode (anode):

$$2Cl^{-}(l) \rightarrow Cl_{2}(g) + 2e^{-}/Cl^{-}(l) \rightarrow \frac{1}{2}Cl_{2}(g) + e^{-};$$

Negative electrode (cathode):

$$Mg^{2+}(l) + 2e^- \rightarrow Mg(l)$$
;

Accept e instead of e-.

Award [1 max] for correct half-equations given at the wrong electrode.

Penalize use of reversible arrows once only.

correct state symbols in both equations;

[3]

(b) aluminium/Al is less dense (compared to iron/Fe) / Al is more ductile or malleable / aluminium forms a protective oxide layer / At does not corrode / iron/Fe rusts /

Do not accept "Al is lighter" OR "less expensive" OR "Al can be recycled".



[1]

- Ethanol: -2/-II; (ii)
 - Ethanal: -1/-1;

[2]

Do not accept 2-, 1- but penalize once only.

(iii) CH,CH,OH → CH,CHO + 2H+ + 2e-;

[1]

Half-equation required. Do not accept $C_3H_3OH + 2[O] \rightarrow CH_3CHO + H_3O$.

Accept e for e-.

(iv) $3CH_3CH_2OH(aq) + Cr_2O_7^{2-}(aq) + 8H^+(aq) \rightarrow 2Cr^{3+}(aq) + 3CH_3CHO(l) + 7H_2O(l)$

correct reactants and products; correct balancing;

[2]

M2 can only be scored if M1 correct.

Ignore state symbols.

Topic Chem 9 Q# 169/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

Positive electrode (anode): (d)

$$2Br^{-} \rightarrow Br_{2}(g) + 2e^{-} / Br^{-} \rightarrow \frac{1}{2}Br_{2}(g) + e^{-}$$
;

Negative electrode (cathode):

$$Na^+ + e^- \rightarrow Na(l)$$
;

[2]

Award [1 max] for correct equations at incorrect electrodes.

Ignore state symbols.

Accept e instead of e-.

Penalize use of equilibrium signs once only.

Positive electrode (anode):

bromine/Br₂;

Negative electrode (cathode):

hydrogen/H₂;

Allow sodium hydroxide/NaOH/hydroxide/OH formation.



Allow voltage instead of potential.

- (e) bromine/Br₂;

 Do not accept bromide/Br⁻.
- (f) (i) potential of reduction half-reaction under standard conditions measured relative to standard hydrogen electrode/SHE / potential under standard conditions relative to standard hydrogen electrode/SHE; Instead of standard state allow either solute concentration of 1 mol dm⁻³ or 100 kPa/1.00 × 10⁵ Pa for gases. Allow 1 bar for 100 kPa/1.00 × 10⁵ Pa. Allow 1 atm.

direction of electron flow

Salt bridge

Cu (s)

correct diagram including (voltmeter), 4 correct species (state symbols not required) and connecting wires;

Cu2+ (aq)

No credit if wires to electrodes immersed in the solutions. Accept ammeter/meter/lamp instead of voltmeter.

Mg2+ (aq)

labelled salt bridge;

Accept an appropriate salt (name or formula) instead of salt bridge (eg, potassium nitrate).

correctly labelled electrodes as +/cathode and -/anode; flow of electrons from Mg to Cu in external circuit;

- (iii) Random uncertainty: (±) 0.01 (V); Significant figures: 3;
- (iv) repeat readings and take an average / use more precise equipment;

[4]

[1]

[1]

[2]

[1]



Topic Chem 10 **Q# 171/** IB Chem/2019/w/TZ0/P2/Higher Level/Q4./

3.	a	i	«electrophilic» addition ✓	Do not accept "nucleophilic addition" or "free radical addition". Do not accept "halogenation".	1	
3.	а	ii	2-chloropropane ✓	Do not accept halogenation.	1	
3.	а	iii	secondary carbocation/carbonium «ion» is more stable		1	
			OR		1	
			carbocation/carbonium «ion» stabilized by two/more alkyl groups ✓			
3.	а	iv	$CH_3CHCICH_3(I) + OH^-(aq) \rightarrow CH_3CH(OH)CH_3(aq) + CI^-(aq)$ OR		1	
			$CH_3CHClCH_3(l) + NaOH(aq) \rightarrow CH_3CH(OH)CH_3(aq) + NaCl(aq) \checkmark$			
3.	b	i	Rate = k [C₃H₁Cl] [OH⁻] ✓			
			«[OH⁻] held constant and» [C₃H₂Ct] triples AND rate triples «so first order wrt C₃H₂Ct» ✓		3	
			[C₃H₂CI] doubles AND [OH⁻] doubles AND rate quadruples «so first order wrt OH⁻» ✓	-	170	
3.	b	ii	SN2√	Accept 'bimolecular nucleophilic substitution.'	1	
3.	a	i	«electrophilic» addition ✓	Do not accept "nucleophilic addition" or "free radical addition". Do not accept "halogenation".	1	
i.	a	ii	2-chloropropane ✓		1	
3.	а	Ш	secondary carbocation/carbonium «ion» is more stable OR carbocation/carbonium «ion» stabilized by two/more alkyl groups ✓		1	
3.	а	iv	$CH_3CHClCH_3\left(l\right) + OH^-(aq) \rightarrow CH_3CH(OH)CH_3\left(aq\right) + Cl^-(aq)$ OR $CH_3CHClCH_3\left(l\right) + NaOH\left(aq\right) \rightarrow CH_3CH(OH)CH_3\left(aq\right) + NaCl\left(aq\right) \checkmark$		1	
3.	b	1	Rate = k [C ₃ H ₇ Cl] [OH ⁻] \checkmark «[OH ⁻] held constant and» [C ₃ H ₇ Cl] triples <i>AND</i> rate triples «so first order wrt C ₃ H ₇ Cl» \checkmark [C ₃ H ₇ Cl] doubles <i>AND</i> [OH ⁻] doubles <i>AND</i> rate quadruples «so first order wrt OH ⁻ » \checkmark	17	3	
3.	b	ii	SN2 ✓	Accept 'bimolecular nucleophilic substitution.'	1	
3.	b	iii	CH ₃ CH ₃ CH ₃ CH ₃ HO CL CH ₃ HO CH ₃ HO CH ₃ CH ₃ HO CH ₃ HO CH ₃ CH ₃ CH ₃ CH ₃ CH ₃ CH ₃ HO CH ₃	Do not allow arrow originating on H in OH*. Allow curly arrow going from bond between C and Cl to Cl in either reactant or transition state. Do not award M3 if OH–C bond is represented. Accept formation of NaCl instead of Cl*.	4	
			representation of transition state showing negative charge, square brackets and partial bonds ✓			
			formation of CH₃CH(OH)CH₃ AND CI-✓			

SMASHING LLI

3.	d	i	K₂Cr₂O₂/Cr₂O₂²⁻/«potassium» dichromate «(VI)» AND acidified/H⁺ OR «acidified potassium» manganate(VII) / «H⁺ and» KMnO₄ / «H⁺ and» MnO₄⁻ ✓	Accept "H ₂ SO ₄ " or "H ₃ PO ₄ " for "H+". Do not accept HCl. Accept "permanganate" for "manganate(VII)".	1
3.	d	ii	C₃H₅O/propan-2-ol: hydrogen-bonding <i>AND</i> C₃H₅O/propanone: no hydrogen bonding/«only» dipole–dipole/dispersion forces ✓ hydrogen bonding stronger «than dipole–dipole» ✓		2
	е		CH ₃ H CH ₃ H	Continuation bonds must be shown. Methyl groups may be drawn on opposite sides of the chain or head to tail. Ignore square brackets and "n".	1

opi	1 1	n 10 Q# 172/ IB Chem/2018/w/TZ 0/P2/Higher Level/		
Э.	а	polarity/polar «molecule/bond» OR carbon—halogen bond is weaker than C−H bond ✓		1
9.	b	primary AND Br/bromine is attached to a carbon bonded to two hydrogens OR primary AND Br/bromine is attached to a carbon bonded to one C/R/alkyl «group» ✓	Accept "primary AND Br/bromine is attached to the first carbon in the chain".	1
9.	С	HOTCH3CH2 Br CH3CH2 CH3CH2 CH3CH2 CH3CH2 CH3CH2 CH3CH2 Curly arrow going from lone pair/negative charge on O in HOT to C curly arrow showing Br leaving representation of transition state showing negative charge, square brackets and partial bonds formation of organic product CH3CH2CH2OH AND BrT	Do not allow curly arrow originating on H in HO". Accept curly arrow either going from bond between C and Br to Br in 1-bromopropane or in the transition state. Do not penalize if HO and Br are not at 180° to each other. Do not award M3 if OH-C bond is represented.	4
9.	d	«Lewis» base AND donates a pair of electrons 🗸		1
9.	e	Any two of: choose «most» appropriate reaction «for preparing the target compound» design/discover new reactions/reagents apply this knowledge to other areas of chemistry/science «retro-»synthesis «more effective» control/predict «desired» products control rate of reaction «more effectively» satisfy intellectual curiosity predicting how changing reagents/conditions might affect reaction	Accept other reasonable answers.	2 max



opi	Che	≥m 10	0 Q# 173/ IB Chem/2018/w/TZ 0/P2/Higher Level/	<u> </u>	1
В.	a		angle between bonds is 60°/strained/smaller than 109.5° ✓		1
8.	b	1	Any two of: CH3COCH3 CH3CH2CHO CH2=CHCH2OH CH3OCH=CH2 OH	Accept displayed or condensed structural formulas or skeletal formulas. Accept CH(OH)=CHCH3 and CH2=C(OH)CH3.	2
В.	b	ii	no AND only one «axial/methyl/CH ₃ » substituent «at the ring» OR no AND two «axial» substituents required «for cis/trans-isomers	Accept "no AND «O in the ring and» one carbon has two H atoms".	1
pji	c Che	≥m 10	0 Q# 174/ IB Chem/2018/w/TZ 0/P2/Higher Level/		
7. opi	a c Che	em 10	«electrophilic» addition/A _E OR reduction ✓ 0 Q# 175/ IB Chem/2018/w/TZ 0/P2/Higher Level/	Accept "hydrogenation".	1
5.	d		Any two of: butanoic acid forms more/stronger hydrogen bonds ✓ butanoic acid forms stronger London/dispersion forces ✓ butanoic acid forms stronger dipole—dipole interaction/force ✓	Accept "butanoic acid forms dimers". Accept "butanoic acid has larger M _t /hydrocarbon chain/number of electrons" for M2. Accept "butanoic acid has larger «permanent» dipole/more polar" for M3.	2 max
opi	c Che	m 10	0 Q# 176/ IB Chem/2018/w/TZ 0/P2/Higher Level/		
6.	e	i	lithium aluminium hydride/LiAlH₄ ✓	A	1
6.	e	ii	butan-1-ol/1-butanol/CH₃CH₂CH₂CH₂OH ✓	7	1



Topic Chem 10 Q# 177/ IB Chem/2018/s/TZ 1/P2/Higher Level/

PIC			U U# 1777 IB CHEIT/ 2010/3/12 1/72/ Higher Level/	*****	
	a		Any two similarities: heterolytic bond breaking OR chloride ions leave ✓ nucleophilic/OHT substitution ✓ both first order with regard to [halogenoalkane] ✓ One difference: CH3CH2CH2CH2CI is second order/bimolecular/S _N 2 AND (CH3)3CCI is first order/unimolecular/S _N 1 OR CH3CH2CH2CH2CI rate depends on [OHT] AND (CH3)3CCI does not OR CH3CH2CH2CH2CI is one step AND (CH3)3CCI is two steps	Do not accept "produces alcohol" or "produces NaCl". Accept "substitution in 1-chlorobutane and «some» elimination in 2-chloro-2-methylpropane".	3
			OR (CH ₃) ₃ CCl involves an intermediate AND CH ₃ CH ₂ CH ₂ CH ₂ Cl does not OR CH ₃ CH ₂ CH ₂ CH ₂ Cl has inversion of configuration AND (CH ₃) ₃ CCl has c. 50 : 50 retention and inversion √		
	b		C-Br bond weaker than C-Cl bond ✓	Accept "Br - is a better leaving group". Do not accept "bromine is more reactive". Do not accept "C-Br bond is longer than C-Cl" alone.	1
-			*******		
	C	i	butan-1-ol/CH₂CH₂CH₂CH₂OH ✓	Do not accept "butanol" for "butan-1-ol". Accept "1-butanol". Do not penalize for name if correct formula is drawn.	1
7.	С	ii	«reduction with» lithium aluminium hydride/LiAtH₄ ✓	Do not accept "sodium borohydride/NaBH4".	1
7.	c	iii	ester 🗸		1

Topic Chem 10 Q# 178/ IB Chem/2018/s/TZ 1/P2/Higher Level/

3. a	a í	nickel/Ni «catalyst» ✓ high pressure OR heat ✓	Accept these other catalysts: Pt, Pd, Ir, Rh, Co, Ti. Accept "high temperature" or a stated temperature such as "150 °C".	2
3. a	a ii		Ignore square brackets and "n". Connecting line at end of carbons must be shown.	1
3. b	b	ethyne: $C_2H_2 + Cl_2 \rightarrow CHClCHCl \checkmark$ benzene: $C_6H_6 + Cl_2 \rightarrow C_6H_5Cl + HCl \checkmark$	Accept "CzHzClz".	2



Topic Chem 10 Q# 179/ IB Chem/2017/s/TZ 2/P2/Higher Level/

6. a	Initiation: Cl−Cl → Cl• + Cl• ✓	Do not penalize missing electron dot on radicals if consistent throughout.	
	Propagation:	Accept Cl ₂ , HCl and CH ₃ Cl without showing bonds.	3
	Cl-Cl+ ·CH ₄ → Cl-H+ ·CH ₅ ✓ Cl-Cl+ ·CH ₄ → Cl-CH ₄ + Cl- ✓	Do not accept hydrogen radical, H• or H, but apply ECF to other propagation steps.	
6 b	«ΔH ^O =» −82.0 «kJ» − 92.3 «kJ» − (−74.0 «kJ») ✓ «ΔH ^O =» −100.3 «kJ» ✓	Award [2] for correct final answer.	2
6 c	H ₂ C==CHOL OR H CL	Accept "CH2CHCl" or "CHClCH2". Do not accept "C2H3Cl".	1
	H - H - / / / / / / / / / / / / / / / /		

Topic Chem 10 Q# 180/ IB Chem/2017/s/TZ 2/P2/Higher Level/

2	a	v	hydroxyl/OH	Accept "hydroxy/alcohol" for "hydroxyl".	
			OR	Accept amino/amine/NH ₂ .	1
			aldehyde/CHO ✓		40

Topic Chem 10 Q# 181/ IB Chem/2017/s/TZ 1/P2/Higher Level/

7.	a	Any two of: planar «X-ray» ✓ C to C bond lengths all equal OR	5	2 max
		C to C bonds intermediate in length between C−C and C−C ✓ all C−C−C bond angles equal ✓		
7.	b	benzene: «electrophilic» substitution/S _E AND cyclohexene: «electrophilic» addition/A _E ✓	Accept correct equations.	1
7.	С	«concentrated» nitric AND sulfuric acids ✓ †NO₂ ✓	Accept NOz ⁺ .	2
7.	d	curly arrow going from benzene ring to N of *NO2* ✓ carbocation with correct formula and positive charge on ring ✓	Accept mechanism with corresponding Kekulé structures. Do not accept a circle in M2 or M3. Accept first arrow starting either inside the circle or on the circle. M2 may be awarded from correct diagram for M3. M4: Accept C ₆ H ₆ NO ₂ + H ₂ SO ₄ if HSO ₄ ⁻¹ used in M3.	4
7.	e	curly arrow going from C−H bond to benzene ring of cation ✓ formation of organic product AND H ⁺ ✓ Fe/Zn/Sn AND HCI/H ₂ SO ₄ /CH ₃ COOH ✓ NaOH/KOH ✓	Accept other suitable metals and acids. Accept other suitable bases. Award [1 max] for single-step reducing agents (such as H ₂ /Pt, Na ₂ S etc.).	2



Topic Chem 10 Q# 182/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

(Question	Answers	Notes	Total
6.	а	H ₅ C ₂ CH ₃ H ₃ C C ₂ H ₅ Correct isomer ✓ mirror image shown clearly ✓		2
6.	b	S _N 2 would give inversion of configuration «almost 100%» OR S _N 1 would give «approximately» 50% of each so mechanism is a mixture of both mechanisms ✓		2
6.	С	C–I bond «longer, so» weaker «than C–Br bond» OR I is a better leaving group than Br ✓		1

Topic Chem 10 Q# 183/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

Question Answers	Notes Total
Answers a Propane:	Notes Total

(Quest	ion	Answers	Notes	Total
5.	С	i	$C_3H_8 + Br_2 \rightarrow C_3H_7Br + HBr \checkmark$ «sun»light/UV/ hv OR high temperature \checkmark	Do not accept "reflux" for M2.	2
5.	c	ii	$C_3H_6 + Br_2 \rightarrow C_3H_6Br_2 \checkmark$		1
5.	С	iii	Propane: «free radical» substitution / S _R AND Propene: «electrophilic» addition / A _E ✓		1
			CH ₃ H CH ₃ H CH ₃ H CH ₃ H CH ₄ Br Curly arrow going from C=C to H of HBr and curly arrow showing Br leaving ✓ representation of carbocation ✓ curly arrow going from lone pair/negative charge on Br to C*✓	1-bromopropane.	3



Topic Chem 10 Q# 184/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

Topic Chem 10 Q# 185/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

_	_		and the same of th	_
ď	iv	Similarity: both involve shared pair of electrons / both are covalent ✓ Difference: Pr-N: pair of electrons comes from nitrogen / coordinate bond and N-H: one electron comes from each bonded atom ✓		2
d	v	London / dispersion / instantaneous induced dipole-induced dipole ✓ dipole-dipole ✓ hydrogen bonding ✓	Award [2] for all three correct, [1] for any two correct.	2 ma

Topic Chem 10 Q# 186/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

(Questi	on	Answers	Notes	Total
7.	3		I: carboxamide ✓ II: phenyl ✓ III: carboxyl / carboxy ✓ IV: hydroxyl ✓	Award [2] for all four correct, [1] for two or three correct. Do not allow benzene. Do not allow carboxylic/alkanoic acid. Do not allow alcohol or hydroxide.	2 max
	b	iii	⟨(0.5)(40-20-2)-⟩9 ✓	2 2 2 2 2 2	1
(Questi	on	Answers	Notes	Total
7.	d	i	CI NH ₃ and H ₃ N CI NH ₃ CI NH ₃ CI Transplatin	Names of complexes are not required. Complexes may be drawn without tapered bonds.	1
	d	ii	M CI NH3 and H3N CI NH3 M=0 Cisplatin Transplatin Cis: polar and trans: non-polar		2

Topic Chem 10 Q# 187/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

Question	Answers	Notes	Total
a	HOCH3 HOCH3 HOCH3 HOCH3 HOCH3 HOCH3 HOCH3 HOCH3 HOCH3 CH3	Do not allow curly arrow originating on H in HO Accept curly arrow going from bond between C and I to I in 1-iodoethane or in the transition state. Do not allow arrow originating from C to C-I bond. Do not award M3 if OHC bond is represented. Inversion of configuration must be shown to score M4.	4
ь	Rate expression: rate = k[OH^][CH_1CH_1] Molecularity of RDS: bimolecular ✓		2

Question	Answers	Notes	Total
c	polar, protic solvents decrease nucleophilic reactivity due to hydrogen bonding OR polar, protic solvents have a cage of solvent molecules surrounding anionic nucleophile resulting in increased stabilization (so are slower) OR polar, aprotic solvents have no hydrogen bonding so S _N 2 reactions are favoured since nucleophiles do not solvate effectively so have an enhanced/pronounced effect on nucleophilicity of anionic nucleophiles (so are faster) ✓ S _N 1: polar, protic solvents favour S _N 1 reactions since the carbocation (intermediate) is solvated by ion-dipole interactions by the polar solvent ✓		2
d	DMF since aprotic solvent so favours S _N 2 ✓		1

Topic Chem 10 Q# 188/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

5.	a	i			1
5.	a	II	Test:	Accept other correct reagents, such as manganate(VII) or iodine solutions, and descriptions of the corresponding changes observed. Accept "decolourized" for A and "not decolourized/unchanged" for B. Do not accept "clear/transparent" instead of "colourless".	2

Topic Chem 10 Q# 189/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

2.	b	i	primary ✓		1
2.	b	ii	ALTERNATIVE 1: <pre></pre>	Accept specific equations having correct reactants. Do not accept LiAlH4 or NaBH4. Accept Pt or Pd catalyst. Accept equations having correct reactants.	2



- (Quest	ion	Answers	Notes	Total
2.	b	111	HNO ₃ + 2H ₂ SO ₄ ⇒ NO ₂ * + 2HSO ₄ * + H ₃ O* ✓	Accept: $HNO_3 + H_2SO_4 \rightleftharpoons NO_2^* + HSO_4^- + H_2O$ Accept $HNO_3 + H_2SO_4 \rightleftharpoons H_2NO_3^* + HSO_4^-$. Accept equivalent two step reactions in which sulfuric acid first behaves as a strong acid and protonates the nitric acid, before behaving as a dehydrating agent removing water from it.	1
2. b		iv	NO ₂ + H NO ₂ curly arrow going from benzene ring to N of *NO ₂ /NO ₂ * ✓ carbocation with correct formula and positive charge on ring ✓ curly arrow going from C–H bond to benzene ring of cation ✓	Accept mechanism with corresponding Kekulé structures. Do not accept a circle in M2 or M3. Accept first arrow starting either inside the circle or on the circle. M2 may be awarded from correct diagram for M3. M4: Accept C ₈ H ₈ NO ₂ + H ₂ SO ₄ if HSO ₄ ⁻¹ used in M3.	4
			formation of organic product nitrobenzene AND H* ✓		

Question	Answers	Notes	Total
. c i	Name: ethane-1,2-diol ✓ Class: alcoholess ✓	Accept ethan-1,2-diol / 1,2-ethanediol. Do not accept "diol" for Class.	2

Topic Chem 10 Q# 190/ IB CHEM/2015/w/TZ0/P2 Section B/Higher Level/

9. (a) (i) RBr(l)+NaOH(aq) → ROH(aq)+NaBr(aq) / RBr (l) + OH⁻ (aq) → ROH(aq) + Br⁻ (aq); Ignore state symbols.

[1]

(ii) $(1.35 \times 10^{-2} - 7.36 \times 10^{-3} =) 6.14 \times 10^{-3} / 6.1 \times 10^{-3} \text{ (mol)};$

[1]

(iii) (molar mass = $\frac{0.842}{6.14 \times 10^{-3}}$ =)137 (g mol⁻¹); Accept 138.

- [1]
- (iv) $(137 80 = 57 \text{ which corresponds to } C_4H_9$, hence molecular formula) C_4H_9Br ;

[1]

Accept correct structural formula of one of the isomers as molecular formula.

Do not accept ECF from 9a(iii) for an impossible molecular formula, such as C₄H₁₀Br.



(v) CH₃CH₂CH₂CH₂Br and primary;

(CH₃)₂CHCH₂Br and primary;

CH3CHBrCH2CH3 and secondary;

(CH₃)₃CBr and tertiary;

If primary, secondary or tertiary not stated, award [3] for four correct, [2] for

three correct and [1] for two correct structural formulas.

Penalize missing hydrogens once only.

Accept either full or condensed structural formulas.

If C5H11Br was used, accept any correct structural formulas.

CH₃CH₂CH₂CH₂CH₂Br and primary;

(CH₃)₂CHCH₂CH₂Br and primary;

CH₃CH₂CH(CH₃)CH₂Br and primary;

(CH₃)₃CCH₂Br and primary;

CH3CHBrCH2CH2CH3 and secondary;

CH3CH2CHBrCH2CH3 and secondary;

CH3CHBrCH(CH3)2 and secondary;

CH₃CH₂C(CH₃)₂Br and tertiary;

[4]

If primary, secondary or tertiary not stated, award [3] for four correct, [2] for three correct and [1] for two correct structural formulas.

Penalize missing hydrogens once only.

Accept either full or condensed structural formulas.

(b) (i) increase in carbon chain;

[1]

OR

curly arrow going from lone pair/negative charge on NC⁻ to C; curly arrow showing Br leaving;

Accept curly arrow either going from bond between C and Br to Br in 1-bromobutane/1-bromo-2-methylpropane or in the transition state.

representation of transition state showing negative charge, square brackets and partial bonds;

Do not penalize if NC and Br are not at 180° to each other.

Products: CH₃CH₂CH₂CH₂CN/(CH₃)₂CHCH₂CN and Br⁻/KBr,

[4]

Penalize missing hydrogens and/or incorrect bond linkages (eg CN–C instead of NC–C) only once in this question.

Remember to apply ECF from part (a)(v).

(iii) $CH_2CH_2CH_2CN + 2H_2 \rightarrow CH_3CH_2CH_2CH_2CH_2NH_2$

OR

 $(CH_3)_2CHCH_2CN + 2H_2 \rightarrow (CH_3)_2CHCH_2CH_2NH_2$;

Ni / Co / Ru:

Accept Pt/Pd/Cu/Rh.

Accept either chemical symbol or name.



(c)

(11) addition:

-CH(C₂H₅)-CH₂-CH(C₂H₅)-CH₂- / -C(CH₃)₂-CH₂-C(CH₃)₂-CH₂-; [2] Continuation bonds required for the mark.

(d) Initiation:

$$Cl_2 \xrightarrow{UV/hf/hv/heat} 2Cl_{\bullet}$$
;

Reference to UV light or high temperatures must be included.

Propagation:

$$Cl \cdot + C_2H_6 \rightarrow C_2H_5 \cdot + HCl;$$

 $C_2H_8 \cdot + Cl_2 \rightarrow C_2H_8Cl + Cl \cdot ;$

Termination:

$$Cl \cdot + Cl \cdot \rightarrow Cl_2 / C_2H_5 \cdot + Cl \cdot \rightarrow C_2H_5Cl / C_2H_5 \cdot + C_2H_5 \cdot \rightarrow C_4H_{10};$$
 [4]

Accept representation of radical without • (eg, Cl, C2H5) if consistent throughout mechanism.

Accept further chlorination.

Award [3 max] if initiation, propagation and termination are not stated or are incorrectly labelled for equations.

Accept correct description of processes without equations.

Award [3 max] if Br used correctly instead of Cl.

Topic Chem 10 Q# 191/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

Penalize missing hydrogens only once in Question 10.

10. (a) (i)



Accept bromine atoms cis to each other.

2.3-dibromobutane: Do not penalize the incorrect use of spaces, comma or hyphen.

(iii) red/brown/orange/yellow to colourless/decolourized; [1] Do not accept clear.

Do not accept just "decolorized".

(synthesis of) plastics/polymers/organic materials not naturally available / (b) (i) synthetic materials;


wide range of uses/physical properties / versatile;

large industry / many tons of plastics consumed by society / OWTTE; Do not accept "useful" for M2.

Award [1 max] if specific addition polymer and its use is given.

Penalize reference to condensation polymers once only.

(11)



Ignore n.

Brackets are not required for the mark, but continuation bonds are.

Do not penalize if methyl groups are trans to each other.

[1]

[1]

[1]

[2 max]

(c) (i)
$$HO\overline{:}$$
 CH_2CH_3 CH_2CH_3 CH_2CH_3 CH_3CH_3 CH_3CH_3 CH_3 CH_3

curly arrow going from lone pair/negative charge on O in HO⁻ to C; Do not accept curly arrow originating on H in HO-.

curly arrow showing Br leaving;

Accept curly arrow either going from bond between C and Br to Br in 2-bromobutane or in the transition state.

Accept if arrow goes from C-Br bond to/or beyond Br.

representation of transition state showing negative charge, square brackets and partial bonds;

Do not penalize if HO and Br are not at 180° to each other.

Do not award M3 if OH----C bond is represented.

formation of organic product CH3CHOHCH2CH3 and KBr/Br:

[4]

OHT has a negative charge/higher electron density; stronger attraction to the carbon atom with the partial positive charge / OWTTE:

Do not accept just stronger attraction.

Reference to carbon atom needed for M2.

2-methylbutanenitrile; (d) (i)

Accept small errors in spelling.

Accept 2-cyanobutane.

Do not accept butan-2-nitrile.

(ii) hydrogen/H2 and nickel/Ni;

Accept other suitable metal catalysts such as platinum/Pt, palladium/Pd.

[1]

[2]

[1]


(iii)

Accept condensed CH3 branch in structural formula as this was present in

structure given in question.

carbonyl; (e) (i) Accept ketone. [1]

[1]



[1]

Accept condensed or full structural formula.



(f) hydrogen bonding in compound C;

dipole-dipole forces in C / C is more polar;

C has greater molar mass/more dispersion/London/instantaneous induced dipoleinduced dipole forces/van der Waal forces;

[2 max]

Accept converse argument.

Award [1 max] for stronger intermolecular forces.

(g) C₄H₉OH(l)+6O₂(g) → 4CO₂(g)+5H₂O(l); Ignore state symbols.

[1]

 (h) (i) compounds with the same structural formula and different arrangement in space/3D structures;

[1]

Accept molecular formula instead of structural formula.

Do not accept "similar" instead of "same".

(ii) restricted rotation around a (double) bond;

carbon atoms of the C=C/carbon-carbon double bond (in alkene)/carbon atoms of the C-C/carbon-carbon single bond (in cycloalkane) must have two different atoms/groups of atoms / OWTTE;

Do not accept "functional groups" for "groups of atoms" in M2.

[2]

(iii) CH₃ CH₃

and

and

cis

trans;

[2]

Award [1 max] if cis and trans isomers are correctly drawn and identified for alkene other than but-2-ene.

Award [1 max] if student draws and labels one structure correctly but not the other.

Topic Chem 10 Q# 192/ IB Chem/2015/s/TZ2/P2 Section A/Higher Level/

(a) same functional group;

same general formula;

(successive members) differ by CH₂;

similar chemical properties;

gradation in physical properties;

Do not accept "same" instead of "similar", or vice-versa.

[2 max]

(b) Initiation:

Reference to UV light or high temperatures must be included.

Propagation:

$$Br \cdot + C_2H_6 \rightarrow C_2H_5 \cdot + HBr$$
;

$$C_2H_5 \cdot + Br_2 \rightarrow C_2H_5Br + Br \cdot ;$$

Termination:

$$Br \cdot + Br \cdot \rightarrow Br_2 / C_2H_5 \cdot + Br \cdot \rightarrow C_2H_5Br / C_2H_5 \cdot + C_2H_5 \cdot \rightarrow C_4H_{10}$$
;

[4]

Accept representation of radical without • (eg, Br, C₂H₅) if consistent throughout mechanism.

Penalize reference to heterolytic fission once only.

Award [0] to any mechanism involving ions.

Accept further bromination.

Award [3 max] if initiation, propagation and termination are not stated or are incorrectly labelled for equations.

Accept correct description of processes without equations.



Topic Chem 10 Q# 193/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

5. (a) Equation:

 $CH_2CH_2 + H_2O \rightarrow CH_3CH_2OH / C_2H_4 + H_2O \rightarrow C_2H_5OH$;

Conditions:

(concentrated) sulfuric acid/H2SO4;

Do not accept dilute sulfuric acid.

Accept phosphoric acid/H₃PO₄ (on pellets of silicon dioxide) (for industrial preparation).

heat / high temperature;

[3]

Do not accept warm.

Accept high pressure (for industrial preparation) for M3 only if H_3PO_4 is given for M2.

(d) CH₃CH₂OH+CH₃CH₂COOH ⇒ CH₃CH₂OOCCH₂CH₃+H₂O; ethyl propanoate;

[2]

Do not penalize if equilibrium arrow missing.

(e) (i) Repeating unit:

Continuation lines must be shown.

Ignore brackets and n.

Accept condensed formulas such as CH2 and C6H4.

Other product:

H₂O/water;

[2]

(ii) condensation;

[1]

Topic Chem 10 Q# 194/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

(a) CH₄ +Br₂ → CH₃Br + HBr;

[1]

(b) (i) Initiation:

UV/hf/h

Br₂ → 2Br•;

Reference to UV/light or high temperatures must be included.

Propagation:

$$CH_3 \cdot + Br_2 \rightarrow CH_3Br + Br \cdot ;$$

Termination:

Award [1 max] for any one of:

$$Br \cdot + Br \cdot \rightarrow Br_2$$
;

$$CH_3 \cdot + CH_3 \cdot \rightarrow C_5H_6$$
;

[4 max]

Allow representation of radical without • (eg Br, CH₃) if consistent throughout mechanism.

Award [3 max] if initiation, propagation and termination are not stated or are incorrectly labelled for equations.

(ii) methanol/CH₃OH;

- (c) C–I bond is weaker than the C–Br bond so more easily broken;
 C–I bond is longer than the C–Br bond / I larger than Br so bonding electrons not as tightly held / I- is better leaving group than Br-;
- [2]

(d) (i) Positive electrode (anode):

$$2Br^{-} \rightarrow Br_{2}(g) + 2e^{-} / Br^{-} \rightarrow \frac{1}{2}Br_{2}(g) + e^{-};$$

Negative electrode (cathode):

$$Na^+ + e^- \rightarrow Na(l)$$
;

[2]

Award [1 max] for correct equations at incorrect electrodes.

Ignore state symbols.

Accept e instead of e-.

Penalize use of equilibrium signs once only.

(ii) Positive electrode (anode):

bromine/Br2;

Negative electrode (cathode):

hydrogen/H2;

[2]

Allow sodium hydroxide/NaOH/hydroxide/OH formation.

Topic Chem 10 Q# 195/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

7. (a) (i) Ethanal: distill off product as it forms;

Accept distillation.

Ethanoic acid: (heat under) reflux / use excess oxidizing agent;

[2]

(b)

curly arrow going from lone pair/negative charge on O in HO⁻ to C; Do not allow curly arrow originating on H in HO⁻.

curly arrow showing Br leaving:

Accept curly arrow either going from bond between C and Br to Br in bromoethane or in the transition state.

representation of transition state showing negative charge, square brackets and partial bonds;

Do not penalize if HO and Br are not at 180° to each other.

Do not award M3 if OH----C bond is represented.

formation of organic product CH₃CH₂OH and Br⁻; Award [3 max] for correct S_N1 mechanism.

[4]

 (d) (i) chiral/asymmetric carbon / carbon attached to 4 different groups / nonsuper imposable mirror images;

[1]

(ii) enantiomers rotate plane of (plane-) polarized light; in opposite directions (by equal amounts);

[2]

Topic Chem 11 **Q# 196/** IB Chem/2019/w/TZ0/P2/Higher Level/Q4./

3.	d			Accept "all hydrogens belong to methyl groups «which are in identical positions»".	1
----	---	--	--	--	---

Tonic Chem 11 O# 197	/ IB Chem/2019/w/TZ0/P2/Higher Level/Q3./
TODIC CHEILI TT CL TSV	TO CHEIT ZOIS W/TZO/FZ/TIENEL LEVERQS.

2.	b	i	$\ll \frac{100 \times 0.1 \mathrm{cm}^3}{20.1 \mathrm{cm}^3} = 0.5 \%$	1
2.	b	ii	repetition / take several samples «and average» ✓	1

Topic Chem 11 Q# 198/ IB Chem/2018/w/TZ 0/P2/Higher Level/

	-		54.67 (s)(2.50 (s)(4.50 (s)	
8.	c	Chemical shift:		
		3.7-4.8 «ppm» ✓		
			2	
		Splitting pattern:		
		doublet ✓		

Topic Chem 11 Q# 199/ IB Chem/2018/w/TZ 0/P2/Higher Level/

2.	b	«absorption at wavenumber 1700–1750 cm ⁻¹ » C=O/carbonyl ✓	Do not accept "ketone" or "aldehyde".	1
2.	С	«m/z =» 58 ✓		1
2.	d	H O H H C C C H CH₃COCH₃/ H H ✓		1

Topic Chem 11 Q# 200/ IB Chem/2018/w/TZ 0/P2/Higher Level/

l. b	iii	ALTERNATIVE 1: «0.2 °C × 100 = » 3 %/0.03 ✓	Accept values in the range 4.1–5.5 «kJ». Award [2] for correct final answer.	
		«0.03 × 160 kJ» = «±» 5 «kJ» ✓		
		ALTERNATIVE 2:	3	2
		«0.2°C× 100/7.5°C -» 3 %/0.03 √	X	
		«0.03 × 180 kJ» = «±» 5 «kJ» ✓	7	

Topic Chem 11 Q# 201/ IB Chem/2018/s/TZ 2/P2/Higher Level/

1.	g	to reduce random errors	Accept "to ensure reliability".	
		OR		1
		to increase precision ✓	1 1	

Topic Chem 11 Q# 202/ IB Chem/2018/s/TZ 1/P2/Higher Level/

2.	b	multiply relative intensity by «m/z» value of isotope OR find the frequency of each isotope ✓ sum of the values of products/multiplication «from each isotope» OR	Award [1 max] for stating "m/z values of isotopes AND relative abundance/intensity" but not stating these need to be multiplied.	2
		find/calculate the weighted average ✓		
		A Company of the Comp	1	

Topic Chem 11 Q# 203/ IB Chem/2018/s/TZ 1/P2/Higher Level/

4	t circi	60: CON2H4 ⁺ ✓	DECEMBER OF ANALYSIS	
	J .		Accept "molecular ion".	1
	1 1	44: CONH ₂ + ✓		
	1 1			2
	1 1			



1. k		3450 cm ⁻¹ : N−H √ 1700 cm ⁻¹ : C=O √	Do not accept "O-H" for 3450 cm ⁻¹ .	2
1. 1	i	1 🗸		1
1. I	ii	singlet ✓	Accept "no splitting".	1
1. 1	III	acts as internal standard OR acts as reference point ✓ one strong signal OR 12 H atoms in same environment OR signal is well away from other absorptions ✓	Accept "inert" or "readily removed" or "non-toxic" for M1.	2

Topic Chem 11 Q# 204/ IB Chem/2016/w/TZ0/P2 Section A/Higher Level/

1.	f	Ethanedioic acid:	Number of signals	Splitting pattern AND singlet ✓	Accept "none/no splitting" for singlet.	2
		Ethane-1,2-diol:	2⊀	Not required		8945

Topic Chem 11 Q# 205/ IB Chem/2016/s/TZoSP/P2 Section A/Higher Level/

b	iv	A: C-H and B: C=0 ✓	1
ь	v	O–H and N–H ✓	
		frequencies/stretches due to O-H and N-H occur above 3200 (cm ⁻¹) which are not present in IR of bute <	2
c	i	1:1:6 ✓	1

Question		Answers	Notes	Total
7. c	i	H H H H—C—C—C—H H O H		1
c	i	ii CH,OCH,CH,✓	F 20 9"	1
c	i i	Similarity: both have fragment corresponding to $(M_*-15)^*/m/z = 45$ ✓ Difference: X has fragment corresponding to $(M_*-17)^*/m/z = 43$ OR X has fragment corresponding to $(M_*-43)^*/m/z = 17$ OR Y has fragment corresponding to $(M_*-31)^*/m/z = 29$ OR Y has fragment corresponding to $(M_*-29)^*/m/z = 31$ ✓	Allow "both have same molecular ion peak/M" / both have m/z = 60". However in practice the molecular ion peak is of low abundance and difficult to observe for propan-2-ol.	2

Topic Chem 11 Q# 206/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

2.	С	ii	two AND two hydrogen environments in the molecule OR two AND both CH₂ and OH present ✓		1
2.	С	iii	*CH₂OH✔	Accept CH ₃ O*. Accept [•CH ₂ OH]* and [•CH ₃ O]*. Do not accept answers in which the charge is missing.	1
2.	С	iv	oxygen-hydrogen €bonds/O-H €in hydroxyls ✓		1

Topic Chem 11 Q# 207/ IB Chem/2016/s/TZ1/P2 Section A/Higher Level/

5.	b	compound A would absorb	rb at 1620-1680) ∉cm ⁻¹ > √	,		Accept any value in range 1620 – 1680 cm ⁻¹ .	1
5.	С	Signal Chemical shift / ppm	1/2 0.9 – 1.0	AND	2/1 4.5 – 6.0	1	Accept 0.9 to 2.0 for the first signal as the C=C affects the CH ₃ shift (actually 1.7).	
		Splitting pattern	singlet	AND	singlet	*	Accept "none/no splitting" for both splitting patterns. Award [1 max] for the correct deduction (both shift and splitting) of signal 1 or 2.	2

(c) (i) 1.4 (%); Accept 1 (%). [1]

(ii) systematic;

dilute the orange juice;

[2]

Accept other valid suggestions, eg. compare with a standard (showing colour at equivalence) / look at mixture through a yellow filter / add more starch (for a sharper colour change) / filter orange juice (through charcoal). Do not accept repeat titrations or alternative indicator.

N₂H₂ sp²;

(iii) 1.44×10⁻⁵ (mol);

[1]

Topic Chem X 92 Q# 209/ IB CHEM/2015/w/TZ0/P2 Section B/Higher Level/

(c)	(i)		NF ₄	
		Hybridization	sp ³ ;	1

N_2H_4	
sp ³ ;	[3]

Do not penalize if it is not superscript.

 (ii) sigma bond: overlapping of orbitals end to end/head on / overlapping of orbitals along internuclear axis;

Accept "axial overlapping of orbitals".

pi bond: side-ways/parallel overlapping of <u>p orbitals</u> (above and below internuclear axis);

[2]

Accept "overlapping of <u>p orbitals</u> above and below the internuclear axis". Accept suitable labelled diagrams.

(iii)





See saw;

Octahedral;

[4]

Accept any combination of lines, dots or crosses to represent electron pairs.

Penalize missing lone pairs on fluorine atoms once only.

For SF4 lone pair on S required for the mark.

Do not allow ECF for a wrong Lewis structure.

Accept saw horse/distorted tetrahedron instead of see-saw.

(d) (i) CH₃CH₂CH₃ < CH₃CHO < CH₃CH₂OH < CH₃COOH ;;

[2]

Award [2] for correct order.

Award [1] for CH₃COOH > CH₃CH₂OH > CH₃CHO > CH₃CH₂CH₃ as compounds are not listed in order of increasing boiling point.

Award [1] if one error in the order.



(iii) $|\overline{F}|$ $|\overline{F}$

[4]

Accept any combination of lines, dots or crosses to represent electron pairs.

Penalize missing lone pairs on fluorine atoms once only.

For SF₄, lone pair on S required for the mark.

Do not allow ECF for a wrong Lewis structure.

Accept saw horse/distorted tetrahedron instead of see-saw.

(d) (i) CH₃CH₂CH₃ < CH₃CHO < CH₃CH₂OH < CH₃COOH ;; Award [2] for correct order. [2]

Award [1] for CH₃COOH > CH₃CH₂OH > CH₃CHO > CH₃CH₂CH₃ as compounds are not listed in order of increasing boiling point.

Award [1] if one error in the order.

(ii) CH₃CH₂CH₃ /London/dispersion/instantaneous induced dipole-induced dipole forces

CH₃CHO dipole-dipole forces (and London/dispersion forces)
CH₃CH₂OH H-bonding (and dipole-dipole and London/dispersion forces)
CH₃COOH H-bonding (and dipole-dipole and London/dispersion forces);
Award [2] for all four correct.

Award [1] for two or three correct.

H-bonding strongest / London/dispersion forces weakest / dipole-dipole stronger than London/dispersion / dipole-dipole weaker than H-bonding;

Accept van der Waals' forces for London/dispersion forces.

CH₃COOH forms more/stronger H-bonds than CH₃CH₂OH / CH₃COOH is more polar than CH₃CH₂OH;

Accept CH₃COOH has more electrons/higher molar mass than CH₃CH₂OH.

(e) ionization and (bombardment) by high energy/fast moving electrons/electron gun

(to form positive ions); acceleration **and** passing through electric field/potential difference/oppositely charged plates;

deflection and passing through magnetic field/electromagnet;

Award [1] for naming 3 processes (ionization, acceleration, deflection) in the correct order with incorrect details.

[3]

[4]

Topic Chem X 93 Q# 210/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

9. MgCl2 conducts electricity and PCl3 does not; (a) (i) MgCl2 is ionic and PCl3 is covalent/molecular; ions/charged particles can move in MqCl₂ / no free charged particles in PCl₃; [3] Award [1 max] if all three points correct for one substance but not other. (ii) MqCl2: 4≤pH≤6.9; high charge density/high charge and small size of Mg2+ makes [Mg(H2O)8]2+ hydrolyse / polarizes water to produce H*; PCla: $0 \le pH \le 4$; (reacts with water to) form HCl/H₃PO₃; [4] Do not accept H₃PO₄. Na, Mg (oxides): basic (b) (i) Al (oxide): amphoteric Do not accept amphiprotic. Si to Cl (oxides): acidic [2] Award [2] for all three listed sets correct. Award [1] for one or two listed sets correct. Award [1] for stating oxides become more acidic towards right/Cl or more basic towards left/Na. Do not penalize if reference is to Ar instead of Cl. Do not penalize for incorrect formulas of oxides.

(ii) $Na_2O(s)+H_2O(l) \rightarrow 2NaOH(aq)$; $P_4O_{10}(s)+6H_2O(l) \rightarrow 4H_3PO_4(aq)$; [2] Ignore state symbols. $Accept\ P_2O_5(s)+3H_2O(l) \rightarrow 2H_3PO_4(aq)$. Do not award marks if incorrect formulas of the oxides are used.

(c) (i) :Br ·· P·· Br : ; :Br : F: ; :S. :F: ; :

[2]

Penalize lone pairs missing on Br and F once only.

Accept any combination of lines, dots or crosses to represent electron pairs.



	(ii)	PBr ₃	SF ₄										
		Shape: trigonal pyramidal; Accept triangular pyramidal.	Shape: see-saw/K-shaped; Accept distorted tetrahedral.										
		No ECF for shape if Lewis structure is incorrect.											
		Bond angle: Any value 99°<109°; Literature value = 101°.	Bond angle: Any two for [1] of: (F _{eq} -S-F _{eq}): Any value 100°<120° Literature value = 103°										
			(F _{ax} -S-F _{ax}): Any value 175°<185° Literature value = 179°/180°										
			(F _{ax} -S-F _{eq}): Any value 85° < 95°										
		No ECF for angle if shape is inc Do not award mark for correct a		[4]									
	(iii)	non-symmetrical distribution of e cancel because of non-symmetri M2 may also be scored with	a suitable diagra <mark>m sho</mark> wing the vectorial pole moments to sh <mark>ow a</mark> net dipole moment	[2]									
(d)	(i)	electrons;	n (positively charged) nuclei and a pair of sharing (between the carbon and hydrogen										
		OR											
		sigma bond formed by overlap o s orbital from H and p/sp ² from c		[2]									
	(ii)	α : sp ³ and β : sp ² ; Accept if numbers are given as s	subscripts.	[1]									
	(iii)	σ bond: end-on / axial overlap of two orb	itals;										
		π bond: sideways overlap of two (paralle	I) p orbitals;	[2]									
		Accept suitable diagrams for bot	h marks.										
	(iv)	11 σ and 3 π ;		[1]									



Topic Chem 99x99 Q# 211/ IB Chem/2015/w/TZ0/P2 Section A/Higher Level/

- (a) HOOC(CH₂), COOH;
 - Accept COOH(CH₂)₄COOH but not COOH(CH₂)₄HOOC or HOOC(CH₂)₄HOOC. H₂N(CH₂)₈NH₂;

[2]

Accept $NH_2(CH_2)_6NH_2$ but not $H_2N(CH_2)_6H_2N$ or $NH_2(CH_2)_6H_2N$. Accept full or condensed structural formulas.

(b) Type: condensation (polymerization);Structural feature: two functional groups on each monomer;

[2]

Accept this group if correctly identified as part of polymer chain. water/H₂O;

[2]

Topic Chem X 910 Q# 212/ IB CHEM/2015/w/TZ0/P2 Section B/Higher Level/

9. (a) (i) RBr(l)+NaOH(aq) → ROH(aq)+NaBr(aq) / RBr(l)+OH⁻ (aq) → ROH(aq)+Br (aq); Ignore state symbols.

[1]

(ii) $(1.35 \times 10^{-2} - 7.36 \times 10^{-3} =) 6.14 \times 10^{-3} / 6.1 \times 10^{-3} \text{ (mol)};$

[1]

(iii) (molar mass = $\frac{0.842}{6.14 \times 10^{-3}}$ =)137 (g mol⁻¹); Accept 138.

[1]

(iv) (137 – 80 = 57 which corresponds to C₄H₉, hence molecular formula) C₄H₉Br; Accept correct structural formula of one of the isomers as molecular formula. Do not accept ECF from 9a(iii) for an impossible molecular formula, such as C₄H₁₀Br.

[1]

(v) CH₃CH₂CH₂CH₂Br and primary;

(CH₃)₂CHCH₂Br and primary; CH₃CHBrCH₂CH₃ and secondary;

(CH₃)₃CBr and tertiary;

If primary, secondary or tertiary not stated, award [3] for four correct, [2] for three correct and [1] for two correct structural formulas.

Penalize missing hydrogens once only.

Accept either full or condensed structural formulas.

If C5H11Br was used, accept any correct structural formulas.

CH₃CH₂CH₂CH₂CH₂Br and primary;

(CH₃)₂CHCH₂CH₂Br and primary;

CH₃CH₂CH(CH₃)CH₂Br and primary;

(CH₃)₃CCH₂Br and primary;

CH3CHBrCH2CH2CH3 and secondary;

CH3CH2CHBrCH2CH3 and secondary;

CH3CHBrCH(CH3)2 and secondary;

CH₃CH₂C(CH₃)₂Br and tertiary;

[4]

If primary, secondary or tertiary not stated, award [3] for four correct, [2] for three correct and [1] for two correct structural formulas.

Penalize missing hydrogens once only.

Accept either full or condensed structural formulas.



(b) (i) increase in carbon chain;

[1]

OR

curly arrow going from lone pair/negative charge on NC⁻ to C; curly arrow showing Br leaving;

Accept curly arrow either going from bond between C and Br to Br in 1-bromobutane/1-bromo-2-methylpropane or in the transition state.

representation of transition state showing negative charge, square brackets and partial bonds;

Do not penalize if NC and Br are not at 180° to each other.

Products: CH₃CH₂CH₂CH₂CN/(CH₃)₂CHCH₂CN and Br⁻/KBr;

Penalize missing hydrogens and/or incorrect bond linkages (eg CN–C instead of NC–C) only once in this question.

Remember to apply ECF from part (a)(v).

(iii) $CH_3CH_2CH_2CN + 2H_2 \rightarrow CH_3CH_2CH_2CH_2CH_2NH_2$ **OR** $(CH_3)_2CHCH_2CN + 2H_2 \rightarrow (CH_3)_2CHCH_2CH_2NH_2$;

> Ni / Co / Ru; Accept Pt/Pd/Cu/Rh. Accept either chemical symbol or name.

[2]

[4]

$$HO^{-}$$
 HO^{-}
 $HO^{$

curly arrow going from lone pair/negative charge on O in OH⁻ to H on C; Do not allow curly arrow originating on H in OH⁻.

curly arrow going from CH bond to form C=C bond; curly arrow showing Br leaving;

$$H_2C = CHCH_2CH_3 + H_2O + Br^- / H_2C = C(CH_3)_2 + H_2O + Br^-;$$
 [4]

Accept NaBr for Br-.

Remember to apply ECF from part (a)(v).

Do not penalize if mechanism shown with a secondary bromoalkane.

Do not penalize incorrect side-chain.

Penalize missing hydrogens only once in this question.

(ii) addition;

(d) Initiation:

$$Cl_2 \xrightarrow{\text{UV/hf/hw/heat}} 2Cl_{\bullet};$$

Reference to UV light or high temperatures must be included.

Propagation:

$$Cl \cdot + C_2H_8 \rightarrow C_2H_5 \cdot + HCl;$$

 $C_2H_5 \cdot + Cl_2 \rightarrow C_2H_5Cl + Cl \cdot ;$

Termination:

$$Cl \cdot + Cl \cdot \rightarrow Cl_2 / C_2H_5 \cdot + Cl \cdot \rightarrow C_2H_5Cl / C_2H_5 \cdot + C_2H_5 \cdot \rightarrow C_4H_{10};$$
 [4]

Accept representation of radical without • (eg, Cl, C₂H₅) if consistent throughout mechanism.

Accept further chlorination.

Award [3 max] if initiation, propagation and termination are not stated or are incorrectly labelled for equations.

Accept correct description of processes without equations.

Award [3 max] if Br used correctly instead of Cl.



Topic Chem x 910 Q# 213/ IB CHEM/2015/s/TZ1/P2 Section B/Higher Level/

 (a) (i) Ethanal: distill off product as it forms; Accept distillation.

Ethanoic acid: (heat under) reflux / use excess oxidizing agent;

[2]

(ii) Ethanol: -2/-II; Ethanal: -1/-I; Do not accept 2-, 1- but penalize once only.

(iii) $CH_3CH_2OH \rightarrow CH_3CHO + 2H^+ + 2e^-;$ [1] Half-equation required. Do not accept $C_2H_5OH + 2[O] \rightarrow CH_3CHO + H_2O$. Accept e for e^- .

(iv) 3CH₃CH₂OH(aq) + Cr₂O₇²⁻(aq) + 8H⁺(aq) → 2Cr³⁺(aq) + 3CH₃CHO(l) + 7H₂O(l) correct reactants and products; correct balancing;
 M2 can only be scored if M1 correct. Ignore state symbols.

curly arrow going from lone pair/negative charge on O in HO⁻ to C; Do not allow curly arrow originating on H in HO⁻.

curly arrow showing Br leaving;

Accept curly arrow either going from bond between C and Br to Br in bromoethane or in the transition state.

representation of transition state showing negative charge, square brackets and partial bonds;

Do not penalize if HO and Br are not at 180° to each other. Do not award M3 if OH----C bond is represented.

formation of organic product CH₃CH₂OH and Br⁻; Award [3 max] for correct S_N1 mechanism.

[4]

- (c) (i) [NaOH] / [OH⁻] is 1/first order and [C₄H₉Br] is 1/first order; rate = k [OH⁻][C₄H₉Br] / rate = k [NaOH][C₄H₉Br]; [2] Square brackets must be used for M2.
 - (ii) $\left(\frac{1.02 \times 10^{-4}}{0.25 \times 0.25}\right) = 0.0016 / 1.6 \times 10^{-3}$; mol⁻¹ dm³ s⁻¹; Accept M⁻¹ s⁻¹. Ignore order of units. Must use experiment 3 data.
 - (iii) bimolecular/2; [1] Accept dimolecular.



(d)	(i)	chiral/asymmetric carbon / carbon attached to 4 different groups / non- super imposable mirror images;	[1]
	(ii)	enantiomers rotate plane of (plane-) polarized light; in opposite directions (by equal amounts);	[2]
(e)	CH, CH,	CHBrCH ₂ CH ₃ + OH ⁻ \rightarrow CH ₃ CHCHCH ₃ + H ₂ O + Br ⁻ / CHBrCH ₂ CH ₃ + OH ⁻ \rightarrow CH ₂ CHCH ₂ CH ₃ + H ₂ O + Br ⁻ / CHBrCH ₂ CH ₃ + CH ₃ CH ₂ O ⁻ \rightarrow CH ₃ CHCHCH ₃ + CH ₃ CH ₂ OH + Br ⁻ ; ept equations with NaOH. holic sodium hydroxide / ethanolic OH ⁻ ;	[2]
(f)	Sigi resu Acc	ma bonds: o o / O / Ilt from head-on/end-on overlap of orbitals / OWTTE; ept axial overlap of orbitals. ept "symmetric orbital" with respect to same plane / OWTTE.	
	resu Acc	onds:	[2]
(g)	Awa (abu	91 = 79x + 81(1-x); and M1 for any suitable calculation. Indance ⁷⁹ Br =) 54.5 %; and [2] for correct final answer.	[2]



								7			6			5		4			ω			2		<u> </u>		<u>. </u>		
							(223)	Fr	87	132.91	S	55	85.47	Rb	37	39.10	×	19	22.99	Na	11	6.94	Li	ω	1.01	Н	₽	1
							(226)	Ra	88	137.33	Ba	56	87.62	Sr	38	40.08	Ca	20	24.31	Mg	12	9.01	Ве	4				2
	-11-			+			(227)	Ac	\$9 ‡	138.91	La	57 †	88.91	Y	39	44.96	Sc	21				ı			1			ω
232.04	Th	90	140.12	Ce	58		(267)	Rf	104	178.49	Hf	72	91.22	Zr	40	47.87	1	22										4
231.04	Pa	91	140.91	Pr	59		(268)	DЬ	105	180.95	Ta	73	92.91	Nb	41	50.94	٧	23										21
238.03	u	92	144.24	Nd	60		(269)	Sg	106	183.84	8	74	95.96	Mo	42	52.00	Cr	24										6
(237)	Np	93	(145)	Pm	61		(270)	Bh	107	186.21	Re	75	(98)	Tc	43	54.94	Mn	25				В	Relativ		Ele		Atomic	7
(244)	Pu	94	150.36	Sm	62		(269)	Hs	108	190.23	0s	76	101.07	Ru	44	55.85	Fe	26				mass	Relative atomic		Element		Atomic number	8
(243)	Am	95	151.96	Eu	63		(278)	Mt	109	192.22	ī	77	102.91	Rh	45	58.93	00	27										9
(247)	Cm	96	157.25	Gd	64		(281)	Ds	110	195.08	Pt	78	106.42	Pd	46	58.69	N:	28	ij									10
(247)	Bk	97	158.93	Ъ	65	ø	(281)	Rg	111	196.97	Au	79	107.87	Ag	47	63.55	Cu	29										11
(251)	Ωf	98	162.50	Дy	66		(285)	Cn	112	200.59	Hg	80	112.41	Cd	48	65.38	Zn	30	1									12
(252)	Es	99	164.93	Но	67		(286)	Uut	113	204.38	11	81	114.82	In	49	69.72	Ga	31	26.98	A	13	10.81	В	Сī				13
(257)	Fm	100	167.26	달	68		(289)	Uuq	114	207.20	Pb	82	118.71	Sn	50	72.63	Ge	32	28.09	Si	14	12.01	C	6				14
(258)	Md	101	168.93	Tm	69		(288)	Uup	115	208.98	В:	83	121.76	Sb	51	74.92	As	33	30.97	P	15	14.01	Z	7				15
(259)	No	102	173.05	አ	70		(293)	Uuh	116	(209)	Po	84	127.60	Te	52	78.96	Se	34	32.07	S	16	16.00	0	8				16
(262)	Lr	103	174.97	Lu	71		(294)	Uus	117	(210)	At	85	126.90	I	53	79.90	Br	35	35.45	Ω	17	19.00	Ŧ	9				17
			•			1	(294)	Uuo	118	(222)	Rn	86	131.29	Xe	54	83.90	ζ,	36	39.95	Ar	18	20.18	Ne	10	4.00	He	2	18

